



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 11:21 PM UTC

PDB ID : 1QUB / pdb_00001qub
Title : CRYSTAL STRUCTURE OF THE GLYCOSYLATED FIVE-DOMAIN HUMAN BETA2-GLYCOPROTEIN I PURIFIED FROM BLOOD PLASMA
Authors : Bouma, B.; de Groot, P.G.; van den Elsen, J.M.H.; Ravelli, R.B.G.; Schouten, A.; Simmelink, M.J.A.; Derksen, R.H.W.M.; Kroon, J.; Gros, P.
Deposited on : 1999-07-01
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	11% 76% 22% .
2	B	3	33% 67%
3	C	2	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	B	3	X	-	-	-
3	NAG	C	2	X	-	-	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 2607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (human beta2-Glycoprotein I).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2480	1580	410	464	26			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	32	Total	O	0	0
			32	32		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN (human beta2-Glycoprotein I)



- Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	161.17Å 166.49Å 114.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.70 40.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.00-2.70) 99.7 (40.00-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.72Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.249 , 0.269 0.254 , 0.274	Depositor DCC
R_{free} test set	2082 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	55.2	Xtriage
Anisotropy	0.363	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for -k,-h,-l 0.000 for -1/2*h-1/2*k+l,-1/2*h-1/2*k-l,1/2 *h-1/2*k 0.008 for -1/2*h-1/2*k-l,-1/2*h-1/2*k+l,-1/ 2*h+1/2*k 0.008 for -1/2*h+1/2*k-l,1/2*h-1/2*k-l,-1/2 *h-1/2*k 0.000 for -1/2*h+1/2*k+l,1/2*h-1/2*k+l,1/ 2*h+1/2*k	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2607	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	5/2553 (0.2%)	1.27	19/3468 (0.5%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124	PRO	CA-C	6.23	1.57	1.52
1	A	121	ILE	CA-CB	-5.41	1.47	1.54
1	A	55	ILE	CA-CB	5.36	1.60	1.54
1	A	42	MET	SD-CE	5.21	1.92	1.79
1	A	123	CYS	CA-C	-5.12	1.48	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	LEU	CA-C-N	7.37	129.06	119.84
1	A	10	LEU	C-N-CA	7.37	129.06	119.84
1	A	11	PRO	CA-C-N	-6.69	112.84	122.67
1	A	11	PRO	C-N-CA	-6.69	112.84	122.67
1	A	130	THR	N-CA-C	6.68	122.32	111.37
1	A	247	VAL	N-CA-C	-6.31	102.08	108.96
1	A	246	LYS	N-CA-C	-6.10	101.90	110.50
1	A	11	PRO	N-CA-C	5.99	124.81	112.47
1	A	83	TYR	N-CA-C	-5.88	102.18	109.64
1	A	47	CYS	CA-C-N	-5.87	114.51	120.85
1	A	47	CYS	C-N-CA	-5.87	114.51	120.85
1	A	269	ASN	N-CA-C	5.58	118.88	112.57
1	A	128	ILE	N-CA-C	5.53	113.35	107.76
1	A	106	THR	N-CA-C	5.49	118.56	110.59
1	A	222	ASP	N-CA-C	-5.45	106.51	112.72
1	A	138	LYS	CA-C-N	5.29	125.23	119.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	138	LYS	C-N-CA	5.29	125.23	119.78
1	A	85	ASN	N-CA-C	5.23	117.43	110.53
1	A	320	ALA	N-CA-C	-5.02	105.93	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2480	0	2401	50	0
2	B	39	0	34	4	0
3	C	28	0	25	0	0
4	A	28	0	26	1	0
5	A	32	0	0	1	0
All	All	2607	0	2486	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:2:NAG:H4	2:B:3:MAN:O2	1.69	0.92
1:A:2:ARG:HB3	1:A:2:ARG:HH11	1.44	0.82
1:A:186:CYS:CB	1:A:229:CYS:HG	1.95	0.80
1:A:231:LYS:HG3	1:A:232:LEU:HD12	1.65	0.79
1:A:305:LYS:HE2	1:A:305:LYS:H	1.48	0.77
1:A:305:LYS:H	1:A:305:LYS:CE	2.00	0.74
1:A:129:PRO:HG3	1:A:178:LEU:HD13	1.70	0.73
1:A:134:LEU:HD13	1:A:153:PHE:CE1	2.25	0.71
1:A:255:VAL:HB	1:A:278:SER:HB2	1.74	0.68
1:A:43:ARG:HG3	1:A:43:ARG:HH11	1.58	0.68
1:A:283:ASN:C	1:A:283:ASN:HD22	1.99	0.68
1:A:2:ARG:HB3	1:A:2:ARG:NH1	2.11	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:TYR:O	1:A:148:ARG:HB2	1.99	0.62
1:A:265:GLU:O	1:A:268:LYS:HG3	2.03	0.58
1:A:232:LEU:HD12	1:A:232:LEU:N	2.19	0.58
1:A:196:PHE:CZ	1:A:214:GLY:HA3	2.38	0.58
1:A:155:CYS:SG	1:A:161:MET:HB2	2.44	0.57
1:A:135:ARG:HD2	1:A:136:VAL:HG23	1.88	0.55
1:A:186:CYS:HB2	1:A:229:CYS:SG	2.46	0.55
1:A:189:PRO:HD2	1:A:199:TYR:CE2	2.43	0.54
1:A:164:ASN:OD1	1:A:166:THR:N	2.37	0.53
1:A:7:PRO:HD3	1:A:22:TYR:HE1	1.74	0.52
1:A:283:ASN:C	1:A:283:ASN:ND2	2.64	0.52
1:A:292:GLU:HB2	1:A:304:PRO:HG2	1.92	0.51
1:A:186:CYS:HB2	1:A:229:CYS:HG	1.73	0.51
1:A:134:LEU:HD13	1:A:153:PHE:HE1	1.74	0.50
2:B:2:NAG:H4	2:B:3:MAN:HO2	1.74	0.50
1:A:207:TYR:CE2	1:A:208:LYS:HG2	2.46	0.50
1:A:166:THR:HG22	4:A:1641:NAG:H82	1.94	0.50
1:A:167:ILE:HG21	1:A:178:LEU:HD23	1.94	0.50
1:A:153:PHE:CE2	1:A:178:LEU:HD22	2.47	0.49
1:A:79:THR:OG1	1:A:80:THR:N	2.46	0.49
1:A:123:CYS:O	1:A:144:ASN:HB3	2.13	0.49
1:A:186:CYS:CB	1:A:229:CYS:SG	2.99	0.48
1:A:157:PRO:O	1:A:158:GLN:HB2	2.14	0.47
1:A:164:ASN:O	1:A:179:PRO:HB3	2.15	0.47
1:A:285:GLU:N	1:A:285:GLU:OE1	2.48	0.47
1:A:283:ASN:HD21	1:A:285:GLU:HB2	1.80	0.46
1:A:23:GLU:O	1:A:24:PRO:C	2.58	0.45
1:A:292:GLU:HB2	1:A:304:PRO:CG	2.46	0.45
1:A:200:PRO:HA	5:A:679:HOH:O	2.17	0.45
1:A:251:LYS:HA	1:A:264:GLN:OE1	2.17	0.45
1:A:262:LYS:HE2	1:A:264:GLN:HE22	1.80	0.44
2:B:2:NAG:H61	2:B:3:MAN:O5	2.17	0.44
1:A:2:ARG:NH1	1:A:2:ARG:CB	2.81	0.43
1:A:284:LYS:HD2	1:A:284:LYS:HA	1.89	0.43
1:A:43:ARG:HG3	1:A:43:ARG:NH1	2.26	0.43
1:A:207:TYR:O	1:A:208:LYS:HB2	2.17	0.43
1:A:303:VAL:O	1:A:304:PRO:C	2.62	0.43
2:B:2:NAG:C4	2:B:3:MAN:O2	2.55	0.43
1:A:210:LYS:HG2	1:A:228:GLU:HG2	2.00	0.42
1:A:262:LYS:HB3	1:A:264:GLN:NE2	2.35	0.42
1:A:208:LYS:HD3	1:A:228:GLU:OE1	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:PHE:CE1	1:A:214:GLY:HA3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	315/319 (99%)	292 (93%)	20 (6%)	3 (1%)	12 32

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	PRO
1	A	319	ASP
1	A	200	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	280/280 (100%)	269 (96%)	11 (4%)	28 57

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ARG
1	A	11	PRO
1	A	84	PRO
1	A	102	SER
1	A	116	PRO
1	A	239	PRO
1	A	283	ASN
1	A	285	GLU
1	A	305	LYS
1	A	310	HIS
1	A	319	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	85	ASN
1	A	92	ASN
1	A	158	GLN
1	A	159	HIS
1	A	283	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	0.69	0	17,19,21	1.00	1 (5%)
2	NAG	B	2	2	14,14,15	0.83	0	17,19,21	0.81	1 (5%)
2	MAN	B	3	2	11,11,12	1.29	1 (9%)	15,15,17	0.52	0
3	NAG	C	1	3,1	14,14,15	0.64	0	17,19,21	1.26	1 (5%)
3	NAG	C	2	3	14,14,15	1.17	1 (7%)	17,19,21	0.71	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	MAN	B	3	2	1/1/4/5	2/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	1/1/5/7	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	NAG	C1-C2	3.02	1.56	1.52
2	B	3	MAN	C1-C2	2.33	1.57	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C2-N2-C7	-3.21	118.60	122.90
3	C	1	NAG	C4-C3-C2	-2.70	107.07	111.02
2	B	2	NAG	C2-N2-C7	-2.00	120.22	122.90
3	C	2	NAG	C2-N2-C7	-2.00	120.22	122.90

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	3	MAN	C1
3	C	2	NAG	C1

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
2	B	3	MAN	O5-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	3	MAN	C4-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	MAN	4	0
2	B	2	NAG	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1641	1	14,14,15	0.86	1 (7%)	17,19,21	0.91	1 (5%)
4	NAG	A	1431	1	14,14,15	1.03	1 (7%)	17,19,21	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1641	1	-	3/6/23/26	0/1/1/1
4	NAG	A	1431	1	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1641	NAG	C3-C2	2.05	1.56	1.52
4	A	1431	NAG	O5-C5	2.03	1.47	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1641	NAG	C1-C2-N2	-2.26	106.88	110.43

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1641	NAG	O5-C5-C6-O6
4	A	1641	NAG	C4-C5-C6-O6
4	A	1431	NAG	C4-C5-C6-O6
4	A	1641	NAG	C3-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1641	NAG	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	310:HIS	C	318:THR	N	9.78

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/319 (100%)	0.50	34 (10%) 11 9	22, 47, 76, 91	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	310	HIS	7.6
1	A	318	THR	6.3
1	A	326	CYS	5.0
1	A	41	GLY	4.9
1	A	40	GLY	4.7
1	A	204	THR	4.3
1	A	135	ARG	3.8
1	A	155	CYS	3.7
1	A	284	LYS	3.7
1	A	306	CYS	3.5
1	A	308	LYS	3.4
1	A	181	CYS	3.4
1	A	39	ARG	3.2
1	A	202	LYS	3.2
1	A	285	GLU	3.2
1	A	286	LYS	3.0
1	A	319	ASP	2.9
1	A	1	GLY	2.9
1	A	2	ARG	2.8
1	A	177	LYS	2.8
1	A	245	CYS	2.8
1	A	42	MET	2.8
1	A	55	ILE	2.4
1	A	228	GLU	2.4
1	A	206	TYR	2.3
1	A	281	CYS	2.3
1	A	251	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	305	LYS	2.2
1	A	201	ALA	2.2
1	A	287	LYS	2.2
1	A	136	VAL	2.1
1	A	309	GLU	2.1
1	A	77	ARG	2.1
1	A	182	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	B	3	11/12	0.32	0.28	88,90,91,91	0
3	NAG	C	2	14/15	0.40	0.31	89,91,92,92	0
2	NAG	B	2	14/15	0.85	0.16	75,81,83,87	0
3	NAG	C	1	14/15	0.89	0.15	75,79,81,86	0
2	NAG	B	1	14/15	0.94	0.10	48,55,59,67	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	1641	14/15	0.77	0.20	80,87,87,88	0
4	NAG	A	1431	14/15	0.80	0.21	82,86,87,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.