



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 10:12 AM EDT

PDB ID : 2QUT / pdb_00002qut
Title : Dihydroxyacetone phosphate enamine intermediate in fructose-1,6-bisphosphate aldolase from rabbit muscle
Authors : St-Jean, M.; Sygusch, J.
Deposited on : 2007-08-06
Resolution : 1.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	2% 75% 20% • •
1	B	363	3% 75% 21% •
1	C	363	2% 78% 19% •
1	D	363	6% 76% 22% • •

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 13663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-bisphosphate aldolase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	350	Total	C	N	O	S	0	0	0
			2673	1680	475	507	11			
1	B	350	Total	C	N	O	S	0	0	0
			2673	1680	475	507	11			
1	C	353	Total	C	N	O	S	0	0	0
			2691	1690	479	511	11			
1	D	360	Total	C	N	O	S	0	0	0
			2730	1715	484	520	11			

- Molecule 2 is 1,3-DIHYDROXYACETONEPHOSPHATE (CCD ID: 13P) (formula: $C_3H_7O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			9	3	5	1		
2	B	1	Total	C	O	P	0	0
			9	3	5	1		

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	O	P	0	0
			9	3	5	1		
2	D	1	Total	C	O	P	0	0
			9	3	5	1		

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	769	Total	O	0	0
			769	769		
3	B	691	Total	O	0	0
			691	691		
3	C	715	Total	O	0	0
			715	715		
3	D	685	Total	O	0	0
			685	685		

- Molecule 1: Fructose-bisphosphate aldolase A





● Molecule 1: Fructose-bisphosphate aldolase A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.77Å 102.91Å 84.23Å 90.00° 98.48° 90.00°	Depositor
Resolution (Å)	50.00 – 1.88 50.00 – 1.88	Depositor EDS
% Data completeness (in resolution range)	87.1 (50.00-1.88) 94.2 (50.00-1.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.80 (at 1.67Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.159 , 0.191 0.160 , 0.195	Depositor DCC
R_{free} test set	6927 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.558	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 85.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13663	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 13P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2725	0.92	7/3691 (0.2%)
1	B	0.35	0/2725	0.90	8/3691 (0.2%)
1	C	0.36	0/2743	0.90	6/3715 (0.2%)
1	D	0.36	0/2782	0.90	9/3770 (0.2%)
All	All	0.35	0/10975	0.90	30/14867 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	3

There are no bond length outliers.

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	347	GLN	OE1-CD-NE2	-9.27	113.33	122.60
1	D	260	VAL	N-CA-C	9.14	115.87	107.56
1	A	260	VAL	N-CA-C	8.76	115.53	107.56
1	A	232	MET	N-CA-C	-8.41	99.19	110.55
1	C	232	MET	N-CA-C	-7.93	99.84	110.55

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	213	TYR	Sidechain
1	C	213	TYR	Sidechain
1	D	213	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2673	0	2693	50	0
1	B	2673	0	2692	48	0
1	C	2691	0	2708	43	0
1	D	2730	0	2753	52	0
2	A	9	0	5	1	0
2	B	9	0	5	0	0
2	C	9	0	5	0	0
2	D	9	0	5	0	0
3	A	769	0	0	12	0
3	B	691	0	0	6	0
3	C	715	0	0	7	0
3	D	685	0	0	10	0
All	All	13663	0	10866	182	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:193:ASP:HA	1:B:237:HIS:HD2	1.42	0.83
1:B:193:ASP:HA	1:B:237:HIS:CD2	2.24	0.72
1:B:291:LEU:O	1:B:293:LYS:HD3	1.90	0.72
1:A:155:GLU:HG2	3:A:3624:HOH:O	1.91	0.71
1:C:336:LEU:HD11	1:C:347:GLN:HA	1.72	0.70

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/363 (95%)	329 (95%)	14 (4%)	3 (1%)	14	4
1	B	346/363 (95%)	329 (95%)	14 (4%)	3 (1%)	14	4
1	C	349/363 (96%)	335 (96%)	12 (3%)	2 (1%)	21	10
1	D	358/363 (99%)	343 (96%)	11 (3%)	4 (1%)	11	3
All	All	1399/1452 (96%)	1336 (96%)	51 (4%)	12 (1%)	14	4

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	PRO
1	A	188	PRO
1	B	5	PRO
1	B	188	PRO
1	C	5	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/291 (98%)	276 (97%)	8 (3%)	38	22
1	B	284/291 (98%)	279 (98%)	5 (2%)	51	38
1	C	285/291 (98%)	280 (98%)	5 (2%)	51	38
1	D	289/291 (99%)	285 (99%)	4 (1%)	59	48
All	All	1142/1164 (98%)	1120 (98%)	22 (2%)	50	36

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	107	LYS
1	C	318	GLU
1	C	295	TRP
1	D	107	LYS
1	A	318	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	287	ASN
1	D	319	ASN
1	C	4	HIS
1	D	85	GLN
1	C	2	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	13P	A	3001	1	8,8,9	1.20	1 (12%)	10,10,12	1.27	1 (10%)
2	13P	C	3003	1	8,8,9	1.24	1 (12%)	10,10,12	1.33	1 (10%)
2	13P	B	3002	1	8,8,9	1.16	1 (12%)	10,10,12	1.33	1 (10%)
2	13P	D	3004	1	8,8,9	1.22	1 (12%)	10,10,12	1.33	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	13P	A	3001	1	-	1/6/6/8	-
2	13P	C	3003	1	-	1/6/6/8	-
2	13P	B	3002	1	-	1/6/6/8	-
2	13P	D	3004	1	-	1/6/6/8	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3003	13P	C2-C3	-2.90	1.35	1.51
2	D	3004	13P	C2-C3	-2.86	1.35	1.51
2	A	3001	13P	C2-C3	-2.85	1.35	1.51
2	B	3002	13P	C2-C3	-2.78	1.35	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3003	13P	C1-C2-C3	3.84	121.12	113.55
2	D	3004	13P	C1-C2-C3	3.83	121.10	113.55
2	B	3002	13P	C1-C2-C3	3.81	121.06	113.55
2	A	3001	13P	C1-C2-C3	3.58	120.61	113.55

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3003	13P	O1-C1-C2-C3
2	D	3004	13P	O1-C1-C2-C3
2	A	3001	13P	O1-C1-C2-C3
2	B	3002	13P	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3001	13P	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/363 (96%)	-0.12	7 (2%) 65 71	10, 19, 42, 58	2 (0%)
1	B	350/363 (96%)	-0.14	10 (2%) 53 58	9, 19, 43, 62	2 (0%)
1	C	353/363 (97%)	-0.17	6 (1%) 69 74	9, 18, 40, 60	4 (1%)
1	D	360/363 (99%)	0.11	20 (5%) 30 32	11, 23, 60, 106	0
All	All	1413/1452 (97%)	-0.08	43 (3%) 52 57	9, 20, 45, 106	8 (0%)

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	348	ALA	7.5
1	D	356	LEU	5.5
1	A	359	SER	5.1
1	C	359	SER	5.1
1	C	346	GLY	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	13P	D	3004	9/10	0.95	0.07	26,28,29,30	0
2	13P	B	3002	9/10	0.96	0.08	23,23,25,29	0
2	13P	C	3003	9/10	0.97	0.09	21,22,23,23	0
2	13P	A	3001	9/10	0.97	0.07	19,21,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.