



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:45 AM UTC

PDB ID : 1QVN / pdb_00001qvn
Title : Structure of SP4160 Bound to IL-2 V69A
Authors : Thanos, C.D.; Delano, W.L.; Wells, J.A.
Deposited on : 2003-08-28
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	8% 72% 20% 7%
1	B	132	7% 79% 11% 8%
1	C	132	8% 81% 11% 8%
1	D	132	5% 73% 18% 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	123	Total	C	N	O	S	0	0	0
			990	636	160	187	7			
1	B	122	Total	C	N	O	S	0	0	0
			972	627	157	181	7			
1	C	122	Total	C	N	O	S	0	0	0
			972	627	157	181	7			
1	D	123	Total	C	N	O	S	0	0	0
			982	632	158	185	7			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	69	ALA	VAL	engineered mutation	UNP P60568
B	69	ALA	VAL	engineered mutation	UNP P60568
C	69	ALA	VAL	engineered mutation	UNP P60568
D	69	ALA	VAL	engineered mutation	UNP P60568

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	2	Total	Zn	0	0
			2	2		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 2-GUANIDINO-4-METHYL-PENTANOIC ACID [2-(4-{5-[4-(4-ACETYLAMINO-BENZYLOXY)-2,3-DICHLORO-PHENYL]-2-METHYL-2H-PYRAZOL-3-YL}-PIPERIDIN-1-YL)-2-OXO-ETHYL]-AMIDE (CCD ID: FRI) (formula: C₃₃H₄₂Cl₂N₈O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			47	33	2	8	4		
3	B	1	Total	C	Cl	N	O	0	0
			47	33	2	8	4		
3	C	1	Total	C	Cl	N	O	0	0
			47	33	2	8	4		
3	D	1	Total	C	Cl	N	O	0	0
			47	33	2	8	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-2




- Molecule 1: Interleukin-2



- Molecule 1: Interleukin-2



- Molecule 1: Interleukin-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.51Å 85.13Å 122.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.70 15.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.1 (15.00-2.70) 96.4 (15.00-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.02 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.258 , 0.308 0.257 , 0.306	Depositor DCC
R_{free} test set	794 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.657	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4108	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FRI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1004	0.84	1/1355 (0.1%)
1	B	0.38	0/985	0.84	0/1330
1	C	0.38	0/985	0.84	0/1330
1	D	0.37	0/996	0.82	0/1346
All	All	0.38	0/3970	0.84	1/5361 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	GLU	N-CA-C	-5.29	106.60	114.64

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	990	0	1016	15	0
1	B	972	0	988	7	0
1	C	972	0	988	10	0
1	D	982	0	998	15	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
3	A	47	0	41	17	0
3	B	47	0	41	9	0
3	C	47	0	41	5	0
3	D	47	0	41	17	0
All	All	4108	0	4154	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:501:FRI:H17	3:D:501:FRI:C16	1.68	1.21
1:D:35:LYS:HG2	3:D:501:FRI:C28	1.76	1.16
3:B:301:FRI:H17	3:B:301:FRI:C16	1.83	1.08
1:D:35:LYS:CG	3:D:501:FRI:H28	1.85	1.05
3:D:501:FRI:H17	3:D:501:FRI:H16	1.02	1.01
1:A:34:PRO:CB	3:A:201:FRI:H421	1.91	1.01
1:A:34:PRO:HB2	3:A:201:FRI:C42	1.92	0.97
1:A:34:PRO:HB2	3:A:201:FRI:H421	0.98	0.97
1:B:73:ALA:HB2	3:B:301:FRI:CL9	2.01	0.97
3:D:501:FRI:H16	3:D:501:FRI:C17	1.97	0.94
3:D:501:FRI:C16	3:D:501:FRI:C17	2.46	0.93
1:D:35:LYS:HG2	3:D:501:FRI:H28	0.91	0.86
3:B:301:FRI:C16	3:B:301:FRI:C17	2.58	0.82
3:B:301:FRI:C17	3:B:301:FRI:H16	2.12	0.80
3:A:201:FRI:C17	3:A:201:FRI:C16	2.65	0.74
1:D:35:LYS:HA	3:D:501:FRI:C25	2.17	0.74
3:A:201:FRI:H16	3:A:201:FRI:C20	2.22	0.69
3:A:201:FRI:C22	3:A:201:FRI:H17	2.22	0.67
1:D:39:MET:HG2	3:D:501:FRI:CL9	2.32	0.66
3:D:501:FRI:O44	3:D:501:FRI:H18	1.94	0.66
3:D:501:FRI:CL10	3:D:501:FRI:H13	2.33	0.66
1:C:35:LYS:CD	3:C:401:FRI:H16	2.27	0.65
1:A:38:ARG:HG3	3:A:201:FRI:H18	1.79	0.63
3:B:301:FRI:H17	3:B:301:FRI:C22	2.29	0.63
1:D:97:LYS:HD2	1:D:103:PHE:HE1	1.64	0.62
1:C:35:LYS:HD2	3:C:401:FRI:H16	1.81	0.62
1:C:73:ALA:HB2	3:C:401:FRI:CL9	2.37	0.61
3:A:201:FRI:C16	3:A:201:FRI:C20	2.77	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:301:FRI:H17	3:B:301:FRI:H16	1.69	0.60
3:A:201:FRI:C17	3:A:201:FRI:C22	2.77	0.60
1:C:28:ILE:HG12	1:C:39:MET:HE1	1.85	0.58
1:A:38:ARG:CG	3:A:201:FRI:H18	2.33	0.57
1:A:35:LYS:N	3:A:201:FRI:H423	2.19	0.57
1:A:39:MET:HE2	1:A:114:ILE:HD13	1.87	0.56
1:D:35:LYS:HA	3:D:501:FRI:H25	1.88	0.56
3:A:201:FRI:H13	3:A:201:FRI:CL10	2.44	0.55
1:C:35:LYS:CG	3:C:401:FRI:H16	2.38	0.54
1:A:35:LYS:HA	3:A:201:FRI:C18	2.39	0.53
1:A:13:GLN:HG3	1:A:92:ILE:HG23	1.91	0.52
1:D:39:MET:HE1	1:D:114:ILE:HD13	1.92	0.51
1:A:38:ARG:HG3	3:A:201:FRI:C18	2.41	0.50
1:B:47:PRO:HG3	1:B:121:TRP:CZ2	2.47	0.50
1:A:34:PRO:C	3:A:201:FRI:C42	2.86	0.49
1:D:33:ASN:HB3	1:D:34:PRO:HD3	1.94	0.49
1:D:31:TYR:HB3	1:D:35:LYS:HB3	1.95	0.48
1:D:35:LYS:HA	3:D:501:FRI:C28	2.43	0.48
1:D:35:LYS:CA	3:D:501:FRI:H25	2.44	0.47
1:B:42:PHE:CZ	3:B:301:FRI:H292	2.48	0.47
1:B:19:LEU:HD11	1:D:15:GLU:HB3	1.97	0.46
3:D:501:FRI:O44	3:D:501:FRI:C18	2.61	0.46
3:D:501:FRI:C17	3:D:501:FRI:C22	2.91	0.45
3:D:501:FRI:H312	3:D:501:FRI:H291	1.76	0.45
1:B:112:ALA:HB1	1:B:116:GLU:HB3	1.99	0.44
1:A:124:PHE:O	1:A:128:ILE:HG12	2.17	0.44
3:B:301:FRI:H16	3:B:301:FRI:C20	2.48	0.44
3:A:201:FRI:CL10	3:A:201:FRI:C13	3.03	0.44
1:B:42:PHE:CE1	3:B:301:FRI:H292	2.54	0.43
1:C:45:TYR:HB3	1:C:107:TYR:CD1	2.54	0.43
1:D:112:ALA:HB1	1:D:116:GLU:HB3	2.00	0.43
1:C:23:MET:HA	1:C:23:MET:HE3	1.99	0.42
1:C:39:MET:HE3	1:C:39:MET:HB3	1.96	0.42
1:C:35:LYS:HA	3:C:401:FRI:H401	2.02	0.42
1:C:47:PRO:HG3	1:C:121:TRP:CZ2	2.55	0.41
1:A:35:LYS:CA	3:A:201:FRI:H423	2.50	0.41
1:A:112:ALA:HB1	1:A:116:GLU:HB2	2.02	0.41
1:D:62:GLU:C	1:D:65:PRO:HD2	2.46	0.41
1:B:57:GLN:HG2	1:B:97:LYS:HE2	2.03	0.41
1:A:71:ASN:HD22	1:A:71:ASN:HA	1.72	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/132 (90%)	116 (98%)	3 (2%)	0	100	100
1	B	116/132 (88%)	115 (99%)	1 (1%)	0	100	100
1	C	116/132 (88%)	113 (97%)	3 (3%)	0	100	100
1	D	119/132 (90%)	116 (98%)	3 (2%)	0	100	100
All	All	470/528 (89%)	460 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/124 (92%)	97 (85%)	17 (15%)	3	8
1	B	109/124 (88%)	96 (88%)	13 (12%)	5	13
1	C	109/124 (88%)	102 (94%)	7 (6%)	16	38
1	D	111/124 (90%)	94 (85%)	17 (15%)	3	7
All	All	443/496 (89%)	389 (88%)	54 (12%)	5	12

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	19	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	23	MET
1	A	28	ILE
1	A	39	MET
1	A	48	LYS
1	A	57	GLN
1	A	60	GLU
1	A	68	GLU
1	A	99	SER
1	A	105	CYS
1	A	106	GLU
1	A	110	GLU
1	A	114	ILE
1	A	120	ARG
1	A	127	SER
1	A	131	THR
1	B	6	SER
1	B	8	LYS
1	B	9	LYS
1	B	15	GLU
1	B	19	LEU
1	B	23	MET
1	B	35	LYS
1	B	57	GLN
1	B	97	LYS
1	B	110	GLU
1	B	116	GLU
1	B	119	ASN
1	B	132	LEU
1	C	9	LYS
1	C	19	LEU
1	C	39	MET
1	C	57	GLN
1	C	83	ARG
1	C	106	GLU
1	C	132	LEU
1	D	6	SER
1	D	9	LYS
1	D	13	GLN
1	D	18	LEU
1	D	19	LEU
1	D	35	LYS
1	D	53	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	54	LYS
1	D	57	GLN
1	D	64	LYS
1	D	71	ASN
1	D	83	ARG
1	D	89	ILE
1	D	97	LYS
1	D	103	PHE
1	D	105	CYS
1	D	106	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	57	GLN
1	A	71	ASN
1	A	119	ASN
1	B	11	GLN
1	B	57	GLN
1	B	119	ASN
1	B	126	GLN
1	C	22	GLN
1	C	126	GLN
1	D	22	GLN
1	D	71	ASN
1	D	90	ASN
1	D	126	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FRI	B	301	-	50,50,50	2.33	5 (10%)	66,70,70	2.85	15 (22%)
3	FRI	C	401	-	50,50,50	2.26	5 (10%)	66,70,70	2.70	12 (18%)
3	FRI	A	201	-	50,50,50	2.04	6 (12%)	66,70,70	2.26	6 (9%)
3	FRI	D	501	-	50,50,50	2.54	6 (12%)	66,70,70	3.03	13 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FRI	B	301	-	-	16/38/48/48	0/4/4/4
3	FRI	C	401	-	-	8/38/48/48	0/4/4/4
3	FRI	A	201	-	-	12/38/48/48	0/4/4/4
3	FRI	D	501	-	-	10/38/48/48	0/4/4/4

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	501	FRI	C13-C14	-12.01	1.21	1.40
3	D	501	FRI	C13-C23	-11.47	1.21	1.37
3	B	301	FRI	C13-C14	-11.16	1.23	1.40
3	C	401	FRI	C13-C14	-10.75	1.23	1.40
3	B	301	FRI	C13-C23	-10.29	1.23	1.37
3	C	401	FRI	C13-C23	-9.86	1.24	1.37
3	A	201	FRI	C13-C23	-8.82	1.25	1.37
3	A	201	FRI	C13-C14	-8.65	1.27	1.40
3	A	201	FRI	C14-N6	3.48	1.38	1.34
3	C	401	FRI	C34-CL10	3.45	1.80	1.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	FRI	C34-CL10	3.39	1.80	1.72
3	A	201	FRI	C34-CL10	3.38	1.80	1.72
3	D	501	FRI	C35-N1	-3.32	1.34	1.41
3	D	501	FRI	C34-CL10	3.28	1.79	1.72
3	C	401	FRI	C33-CL9	3.24	1.79	1.72
3	A	201	FRI	C33-CL9	3.20	1.79	1.72
3	B	301	FRI	C33-CL9	3.09	1.79	1.72
3	D	501	FRI	C33-CL9	2.83	1.78	1.72
3	C	401	FRI	C35-N1	-2.72	1.36	1.41
3	B	301	FRI	C35-N1	-2.67	1.36	1.41
3	A	201	FRI	C35-N1	-2.62	1.36	1.41
3	D	501	FRI	C20-C33	-2.29	1.37	1.40

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	501	FRI	C14-C13-C23	15.19	132.16	106.24
3	B	301	FRI	C14-C13-C23	14.06	130.22	106.24
3	D	501	FRI	C13-C14-N6	-14.05	95.69	111.11
3	B	301	FRI	C13-C14-N6	-13.41	96.39	111.11
3	C	401	FRI	C14-C13-C23	13.29	128.91	106.24
3	C	401	FRI	C13-C14-N6	-12.95	96.89	111.11
3	A	201	FRI	C14-C13-C23	11.25	125.44	106.24
3	A	201	FRI	C13-C14-N6	-10.25	99.86	111.11
3	D	501	FRI	C11-C14-N6	7.42	131.60	119.28
3	A	201	FRI	C11-C14-N6	6.04	129.32	119.28
3	C	401	FRI	C14-N6-N8	5.30	108.43	105.11
3	B	301	FRI	C14-N6-N8	4.84	108.14	105.11
3	D	501	FRI	C35-N1-C12	-4.30	119.95	127.98
3	D	501	FRI	C14-N6-N8	4.23	107.76	105.11
3	C	401	FRI	C11-C14-C13	4.18	138.07	129.78
3	B	301	FRI	C11-C14-N6	4.04	125.99	119.28
3	B	301	FRI	C11-C14-C13	3.83	137.38	129.78
3	D	501	FRI	C23-N8-N6	-3.46	108.77	112.44
3	C	401	FRI	C11-C14-N6	3.44	125.00	119.28
3	A	201	FRI	C43-N2-C31	3.27	119.35	112.68
3	C	401	FRI	C23-N8-N6	-3.20	109.05	112.44
3	B	301	FRI	C35-N1-C12	-3.16	122.08	127.98
3	B	301	FRI	C23-N8-N6	-3.10	109.16	112.44
3	B	301	FRI	C11-C34-C33	-2.98	118.83	120.86
3	C	401	FRI	C35-N1-C12	-2.82	122.71	127.98
3	D	501	FRI	C34-C11-C14	-2.81	121.30	125.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	401	FRI	O45-C20-C33	2.68	120.08	115.77
3	C	401	FRI	C15-C11-C14	-2.67	114.70	119.25
3	A	201	FRI	C34-C11-C14	-2.65	121.52	125.16
3	C	401	FRI	C11-C34-C33	-2.64	119.06	120.86
3	B	301	FRI	C29-C30-N2	2.63	121.71	117.29
3	A	201	FRI	C35-N1-C12	-2.63	123.08	127.98
3	D	501	FRI	C43-N2-C31	2.55	117.88	112.68
3	D	501	FRI	C16-C22-C28	2.44	121.86	118.23
3	B	301	FRI	C30-C29-N3	2.38	116.77	110.90
3	B	301	FRI	C15-C11-C14	-2.29	115.35	119.25
3	D	501	FRI	C11-C34-CL10	2.25	123.32	120.11
3	B	301	FRI	C40-O45-C20	2.18	122.08	117.76
3	B	301	FRI	C11-C34-CL10	2.12	123.14	120.11
3	B	301	FRI	O45-C20-C33	2.09	119.14	115.77
3	D	501	FRI	C11-C34-C33	-2.09	119.44	120.86
3	C	401	FRI	C43-N2-C31	2.07	116.90	112.68
3	D	501	FRI	C20-C33-C34	2.05	121.22	119.86
3	B	301	FRI	C29-N3-C27	2.04	126.53	121.38
3	C	401	FRI	O45-C20-C17	-2.04	119.50	123.95
3	D	501	FRI	O45-C40-C22	-2.04	103.18	109.16

There are no chirality outliers.

All (46) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	201	FRI	O47-C30-N2-C31
3	A	201	FRI	O47-C30-N2-C43
3	A	201	FRI	C29-C30-N2-C31
3	A	201	FRI	C29-C30-N2-C43
3	A	201	FRI	O44-C12-N1-C35
3	A	201	FRI	C42-C12-N1-C35
3	B	301	FRI	O46-C27-N3-C29
3	B	301	FRI	C26-C27-N3-C29
3	B	301	FRI	N7-C26-C41-C32
3	B	301	FRI	O44-C12-N1-C35
3	B	301	FRI	C42-C12-N1-C35
3	C	401	FRI	O44-C12-N1-C35
3	C	401	FRI	C42-C12-N1-C35
3	D	501	FRI	O44-C12-N1-C35
3	D	501	FRI	C42-C12-N1-C35
3	D	501	FRI	C26-C27-N3-C29
3	C	401	FRI	C33-C20-O45-C40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	501	FRI	O46-C27-N3-C29
3	A	201	FRI	C27-C26-C41-C32
3	A	201	FRI	C25-C35-N1-C12
3	A	201	FRI	C18-C35-N1-C12
3	A	201	FRI	N7-C26-C41-C32
3	B	301	FRI	C37-C32-C41-C26
3	A	201	FRI	C37-C32-C41-C26
3	B	301	FRI	N3-C29-C30-O47
3	C	401	FRI	O46-C27-N3-C29
3	B	301	FRI	C36-C32-C41-C26
3	B	301	FRI	C29-C30-N2-C31
3	D	501	FRI	C29-C30-N2-C43
3	C	401	FRI	C26-C27-N3-C29
3	A	201	FRI	C36-C32-C41-C26
3	D	501	FRI	O47-C30-N2-C43
3	B	301	FRI	C33-C20-O45-C40
3	B	301	FRI	N3-C29-C30-N2
3	B	301	FRI	O47-C30-N2-C31
3	B	301	FRI	O47-C30-N2-C43
3	B	301	FRI	C29-C30-N2-C43
3	C	401	FRI	C22-C40-O45-C20
3	C	401	FRI	C17-C20-O45-C40
3	D	501	FRI	O47-C30-N2-C31
3	B	301	FRI	C27-C26-C41-C32
3	D	501	FRI	C29-C30-N2-C31
3	D	501	FRI	N7-C26-C27-O46
3	D	501	FRI	N7-C26-C27-N3
3	B	301	FRI	C17-C20-O45-C40
3	C	401	FRI	C30-C29-N3-C27


There are no ring outliers.


4 monomers are involved in 48 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	FRI	9	0
3	C	401	FRI	5	0
3	A	201	FRI	17	0
3	D	501	FRI	17	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	123/132 (93%)	0.64	10 (8%) 18 15	45, 64, 100, 106	0
1	B	122/132 (92%)	0.57	9 (7%) 20 18	43, 56, 84, 106	0
1	C	122/132 (92%)	0.51	10 (8%) 17 15	41, 58, 91, 110	0
1	D	123/132 (93%)	0.36	7 (5%) 29 26	41, 56, 82, 108	0
All	All	490/528 (92%)	0.52	36 (7%) 21 18	41, 58, 92, 110	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	5	SER	5.9
1	A	100	GLU	5.2
1	B	100	GLU	4.1
1	D	5	SER	3.6
1	A	99	SER	3.5
1	C	101	THR	3.4
1	B	81	ARG	3.2
1	A	105	CYS	3.2
1	D	105	CYS	3.0
1	D	73	ALA	3.0
1	A	4	SER	3.0
1	C	104	MET	2.9
1	B	4	SER	2.8
1	C	4	SER	2.8
1	B	101	THR	2.8
1	A	101	THR	2.8
1	C	81	ARG	2.7
1	A	31	TYR	2.7
1	C	73	ALA	2.7
1	B	104	MET	2.6
1	C	5	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	74	GLN	2.5
1	D	74	GLN	2.5
1	B	30	ASN	2.4
1	B	130	SER	2.4
1	C	98	GLY	2.4
1	D	103	PHE	2.4
1	D	107	TYR	2.4
1	A	102	THR	2.2
1	D	100	GLU	2.2
1	C	30	ASN	2.2
1	B	48	LYS	2.2
1	A	32	LYS	2.1
1	A	73	ALA	2.0
1	C	132	LEU	2.0
1	A	72	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FRI	D	501	47/47	0.76	0.24	70,87,124,128	0
3	FRI	A	201	47/47	0.77	0.20	73,90,150,161	0
3	FRI	B	301	47/47	0.78	0.22	61,72,135,154	0
3	FRI	C	401	47/47	0.86	0.18	59,69,147,161	0
2	ZN	A	701	1/1	0.90	0.11	107,107,107,107	0
2	ZN	B	601	1/1	0.94	0.08	112,112,112,112	0
2	ZN	B	801	1/1	0.97	0.08	76,76,76,76	0
2	ZN	C	901	1/1	0.97	0.05	75,75,75,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FRI D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FRI A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FRI B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FRI C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.