



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:26 AM UTC

PDB ID : 2QVH / pdb_00002qvh
Title : Crystal structure of O-succinylbenzoate synthase complexed with O-succinylbenzoate (OSB)
Authors : Sugadev, R.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2007-08-08
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	9% 80% 15% 5%
1	B	327	7% 78% 16% • 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5018 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-succinylbenzoate-CoA synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	Se	0	0	0
			2349	1470	436	434	7	2			
1	B	311	Total	C	N	O	S	Se	0	0	0
			2349	1470	436	434	7	2			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP Q47Q21
A	2	SER	-	expression tag	UNP Q47Q21
A	3	LEU	-	expression tag	UNP Q47Q21
A	320	GLU	-	expression tag	UNP Q47Q21
A	321	GLY	-	expression tag	UNP Q47Q21
A	322	HIS	-	expression tag	UNP Q47Q21
A	323	HIS	-	expression tag	UNP Q47Q21
A	324	HIS	-	expression tag	UNP Q47Q21
A	325	HIS	-	expression tag	UNP Q47Q21
A	326	HIS	-	expression tag	UNP Q47Q21
A	327	HIS	-	expression tag	UNP Q47Q21
B	1	MSE	-	expression tag	UNP Q47Q21
B	2	SER	-	expression tag	UNP Q47Q21
B	3	LEU	-	expression tag	UNP Q47Q21
B	320	GLU	-	expression tag	UNP Q47Q21
B	321	GLY	-	expression tag	UNP Q47Q21
B	322	HIS	-	expression tag	UNP Q47Q21
B	323	HIS	-	expression tag	UNP Q47Q21
B	324	HIS	-	expression tag	UNP Q47Q21
B	325	HIS	-	expression tag	UNP Q47Q21
B	326	HIS	-	expression tag	UNP Q47Q21
B	327	HIS	-	expression tag	UNP Q47Q21

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is 2-SUCCINYLBENZOATE (CCD ID: OSB) (formula: $C_{11}H_{10}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 16	C 11	O 5	0	0
3	B	1	Total 16	C 11	O 5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	138	Total O 138 138	0	0
4	B	148	Total O 148 148	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: O-succinylbenzoate-CoA synthase



• Molecule 1: O-succinylbenzoate-CoA synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	44.54Å 52.88Å 61.45Å 70.53° 87.22° 85.56°	Depositor
Resolution (Å)	34.25 – 1.76 34.25 – 1.76	Depositor EDS
% Data completeness (in resolution range)	91.1 (34.25-1.76) 91.3 (34.25-1.76)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11.32 (at 1.76Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.218 , 0.251 0.217 , 0.251	Depositor DCC
R_{free} test set	2875 reflections (5.98%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.390	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5018	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 77.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.3023e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OSB, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2391	0.90	6/3262 (0.2%)
1	B	0.35	0/2391	0.92	11/3262 (0.3%)
All	All	0.35	0/4782	0.91	17/6524 (0.3%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	204	VAL	N-CA-C	8.30	118.86	110.23
1	A	204	VAL	N-CA-C	7.76	118.51	110.36
1	B	137	VAL	N-CA-C	6.59	116.75	110.42
1	B	160	ALA	N-CA-C	6.59	118.54	111.36
1	B	149	ARG	N-CA-C	-6.39	105.46	113.20
1	B	232	THR	N-CA-C	-6.15	101.86	110.35
1	A	229	ALA	N-CA-C	-6.07	103.31	111.87
1	A	232	THR	N-CA-C	-5.82	102.32	110.35
1	B	122	GLY	CA-C-N	5.78	125.47	119.05
1	B	122	GLY	C-N-CA	5.78	125.47	119.05
1	B	229	ALA	N-CA-C	-5.66	103.89	111.87
1	A	149	ARG	N-CA-C	-5.66	106.22	113.01
1	B	59	GLU	N-CA-C	5.46	120.44	113.55
1	A	160	ALA	N-CA-C	5.33	117.17	111.36
1	B	76	VAL	N-CA-C	5.24	113.20	107.60
1	A	136	ASP	N-CA-C	-5.19	103.06	110.59
1	B	136	ASP	N-CA-C	-5.14	102.39	110.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2349	0	2364	38	0
1	B	2349	0	2364	40	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	16	0	8	0	0
3	B	16	0	8	0	0
4	A	138	0	0	10	0
4	B	148	0	0	5	0
All	All	5018	0	4744	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:GLN:HE22	1:A:113:ARG:HH22	1.09	0.95
1:B:126:ARG:NH2	1:B:128:ARG:HH21	1.73	0.86
1:B:126:ARG:HH21	1:B:128:ARG:HH21	1.23	0.83
1:A:105:GLN:NE2	1:A:113:ARG:HH22	1.76	0.83
1:A:56:GLU:HG3	4:A:5643:HOH:O	1.79	0.83
1:A:105:GLN:HE22	1:A:113:ARG:NH2	1.80	0.79
1:A:106:SER:H	1:A:109:ASN:HD22	1.31	0.78
1:B:142:ARG:NH1	1:B:143:MSE:HE3	2.02	0.75
1:A:60:LEU:HB2	4:A:5643:HOH:O	1.88	0.72
1:A:5:GLY:HA3	1:A:25:MSE:HE3	1.72	0.70
1:B:97:LYS:HG2	1:B:128:ARG:HB2	1.75	0.69
1:A:126:ARG:NH2	1:A:128:ARG:HH21	1.91	0.68
1:B:157:GLN:HG2	1:B:180:GLU:HG2	1.76	0.67
1:B:288:ARG:HD3	4:B:5663:HOH:O	1.93	0.66
1:B:143:MSE:CE	1:B:146:LEU:HD12	2.26	0.66
1:B:143:MSE:HE1	1:B:146:LEU:HD12	1.78	0.66
1:B:3:LEU:HD22	1:B:3:LEU:H	1.61	0.65
1:A:106:SER:H	1:A:109:ASN:ND2	1.93	0.64
1:B:142:ARG:HH11	1:B:143:MSE:HE3	1.64	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ARG:HH21	1:A:128:ARG:HH21	1.48	0.61
1:B:106:SER:H	1:B:109:ASN:HD22	1.46	0.61
1:B:28:ARG:HD3	1:B:33:TRP:CE2	2.36	0.60
1:B:126:ARG:NH2	1:B:128:ARG:NH2	2.48	0.60
1:A:82:GLU:O	1:A:86:ARG:HG3	2.02	0.60
1:A:2:SER:OG	1:A:6:ARG:HG2	2.02	0.60
1:A:259:ARG:HG3	4:A:5585:HOH:O	2.01	0.59
1:A:103:ARG:HG2	4:A:5611:HOH:O	2.02	0.59
1:B:25:MSE:HE1	1:B:300:GLN:HG3	1.84	0.58
1:B:157:GLN:HE21	1:B:179:ASP:H	1.49	0.58
1:B:126:ARG:HH22	1:B:252:ALA:H	1.52	0.57
1:B:88:VAL:HG11	1:B:121:LEU:HG	1.85	0.57
1:A:28:ARG:HG2	1:A:33:TRP:CE2	2.39	0.56
1:B:49:ARG:HG3	4:B:5621:HOH:O	2.07	0.55
1:B:3:LEU:HD22	1:B:3:LEU:N	2.22	0.55
1:A:25:MSE:HE2	1:A:303:LEU:HD13	1.88	0.55
1:A:157:GLN:HE21	1:A:179:ASP:H	1.57	0.53
1:B:157:GLN:NE2	1:B:179:ASP:H	2.07	0.53
1:B:282:VAL:HG22	4:B:5659:HOH:O	2.08	0.52
1:A:287:GLN:O	1:A:291:GLU:HG3	2.11	0.51
1:B:168:VAL:O	1:B:172:VAL:HG22	2.11	0.51
1:A:157:GLN:NE2	1:A:179:ASP:H	2.10	0.50
1:B:88:VAL:HG21	1:B:117:VAL:HG13	1.93	0.50
1:B:310:TRP:CD1	4:B:5691:HOH:O	2.65	0.49
1:A:126:ARG:HH22	1:A:252:ALA:H	1.60	0.49
1:A:5:GLY:HA3	4:A:5655:HOH:O	2.13	0.48
1:B:158:PRO:HD2	1:B:165:LEU:HD22	1.95	0.48
1:B:106:SER:H	1:B:109:ASN:ND2	2.12	0.48
1:A:97:LYS:HE2	1:A:130:ASP:HB2	1.96	0.47
1:B:143:MSE:HA	1:B:143:MSE:HE2	1.96	0.47
1:B:2:SER:HB3	1:B:300:GLN:HE22	1.79	0.47
1:A:86:ARG:NH1	1:A:86:ARG:HG2	2.29	0.46
1:B:143:MSE:HE2	1:B:146:LEU:HD12	1.98	0.46
1:A:211:ARG:NH1	4:A:5643:HOH:O	2.47	0.46
1:A:28:ARG:HG2	1:A:33:TRP:CZ2	2.51	0.45
1:A:25:MSE:CE	1:A:303:LEU:HD13	2.47	0.45
1:A:25:MSE:HE1	1:A:300:GLN:OE1	2.16	0.45
1:A:303:LEU:C	1:A:303:LEU:HD23	2.42	0.44
1:B:51:TRP:CD1	1:B:307:ARG:HA	2.52	0.44
1:B:157:GLN:HG2	1:B:180:GLU:CG	2.45	0.44
1:B:211:ARG:HB3	1:B:215:ARG:HH12	1.82	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:ILE:HG13	1:A:200:VAL:HG13	1.98	0.44
1:A:122:GLY:HA3	4:A:5684:HOH:O	2.17	0.43
1:B:137:VAL:O	1:B:141:VAL:HG23	2.18	0.43
1:A:50:TRP:CE3	1:A:205:GLN:HB3	2.54	0.43
1:A:126:ARG:NE	1:A:153:GLU:OE2	2.52	0.43
1:A:259:ARG:NH2	4:A:5586:HOH:O	2.50	0.43
1:B:50:TRP:CZ2	1:B:206:PRO:HG3	2.53	0.43
1:B:287:GLN:CD	1:B:287:GLN:H	2.27	0.43
1:A:86:ARG:HG2	1:A:86:ARG:HH11	1.84	0.42
1:B:111:VAL:HB	4:B:5696:HOH:O	2.18	0.42
1:A:126:ARG:NH2	1:A:128:ARG:NH2	2.62	0.42
1:A:220:CYS:HB2	4:A:5616:HOH:O	2.20	0.42
1:B:137:VAL:HG12	1:B:171:ARG:NH2	2.35	0.41
1:A:121:LEU:HB3	1:A:125:GLY:HA3	2.01	0.41
1:B:182:ILE:HG13	1:B:200:VAL:HG13	2.02	0.41
1:B:4:THR:OG1	1:B:5:GLY:N	2.53	0.41
1:B:205:GLN:HB2	1:B:206:PRO:HD3	2.02	0.41
1:A:145:ARG:NH2	4:A:5596:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/327 (94%)	299 (97%)	8 (3%)	2 (1%)	21	8
1	B	309/327 (94%)	301 (97%)	8 (3%)	0	100	100
All	All	618/654 (94%)	600 (97%)	16 (3%)	2 (0%)	36	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	151	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/247 (96%)	234 (99%)	2 (1%)	73	64
1	B	236/247 (96%)	232 (98%)	4 (2%)	53	36
All	All	472/494 (96%)	466 (99%)	6 (1%)	61	46

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	GLU
1	A	189	LEU
1	B	3	LEU
1	B	97	LYS
1	B	180	GLU
1	B	195	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	GLN
1	A	109	ASN
1	A	157	GLN
1	B	109	ASN
1	B	157	GLN
1	B	274	HIS
1	B	300	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	OSB	A	5550	2	16,16,16	1.44	1 (6%)	21,21,21	0.89	0
3	OSB	B	5551	2	16,16,16	1.51	1 (6%)	21,21,21	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OSB	A	5550	2	-	2/13/13/13	0/1/1/1
3	OSB	B	5551	2	-	4/13/13/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	5551	OSB	C1-C2	4.39	1.47	1.40
3	A	5550	OSB	C1-C2	3.88	1.47	1.40

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	5550	OSB	C6-C1-C8-O3
3	A	5550	OSB	C6-C1-C8-C9
3	B	5551	OSB	C6-C1-C8-C9
3	B	5551	OSB	C1-C8-C9-C10
3	B	5551	OSB	C6-C1-C8-O3
3	B	5551	OSB	O3-C8-C9-C10

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/327 (94%)	0.67	30 (9%) 13 15	8, 17, 34, 54	0
1	B	309/327 (94%)	0.54	24 (7%) 19 22	8, 17, 31, 54	0
All	All	618/654 (94%)	0.60	54 (8%) 16 19	8, 17, 32, 54	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	THR	7.3
1	B	4	THR	6.9
1	B	3	LEU	6.1
1	B	2	SER	6.0
1	A	3	LEU	5.8
1	A	5	GLY	5.2
1	A	2	SER	4.5
1	A	259	ARG	4.0
1	B	124	ARG	4.0
1	A	290	ALA	4.0
1	A	187	ASP	3.6
1	A	86	ARG	3.5
1	B	312	GLN	3.5
1	B	195	GLU	3.5
1	A	124	ARG	3.4
1	B	126	ARG	3.4
1	B	123	PRO	3.3
1	B	173	SER	3.3
1	A	292	VAL	3.2
1	A	82	GLU	3.1
1	B	281	ARG	3.0
1	B	5	GLY	2.9
1	A	126	ARG	2.9
1	A	83	GLU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	284	VAL	2.9
1	A	185	ALA	2.9
1	A	186	GLU	2.8
1	A	189	LEU	2.7
1	A	103	ARG	2.7
1	A	288	ARG	2.6
1	B	290	ALA	2.5
1	A	281	ARG	2.4
1	B	103	ARG	2.4
1	A	221	GLY	2.4
1	A	312	GLN	2.4
1	B	300	GLN	2.4
1	A	119	ASP	2.4
1	B	151	GLU	2.4
1	A	268	ASP	2.3
1	A	287	GLN	2.3
1	B	142	ARG	2.3
1	B	85	ALA	2.2
1	A	289	LEU	2.2
1	A	87	ILE	2.2
1	B	89	ALA	2.2
1	B	220	CYS	2.2
1	B	310	TRP	2.1
1	B	288	ARG	2.1
1	A	89	ALA	2.1
1	B	172	VAL	2.1
1	B	307	ARG	2.1
1	B	120	ALA	2.0
1	A	149	ARG	2.0
1	A	136	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	OSB	B	5551	16/16	0.93	0.08	10,14,15,16	0
3	OSB	A	5550	16/16	0.94	0.07	12,15,16,17	0
2	MG	A	401	1/1	0.99	0.06	13,13,13,13	0
2	MG	B	400	1/1	0.99	0.04	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.