



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:21 AM UTC

PDB ID : 2QVY / pdb_00002qvy
Title : 4-Chlorobenzoyl-CoA Ligase/Synthetase, I303G mutation, bound to 3,4-Dichlorobenzoate
Authors : Wu, R.; Reger, A.S.; Cao, J.; Gulick, A.M.; Dunaway-Mariano, D.
Deposited on : 2007-08-09
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	504	5% 70% 26% ..

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3752 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-Chlorobenzoate CoA Ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	499	Total	C	N	O	S	0	0	0
			3716	2349	666	688	13			

- Molecule 2 is 3,4-dichlorobenzoate (CCD ID: 34Z) (formula: $C_7H_4Cl_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	X	1	Total	C	Cl	O	0	0
			11	7	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	X	25	Total	O	0	0
			25	25		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-Chlorobenzoate CoA Ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	127.81Å 127.81Å 71.42Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.76 30.00 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-2.76) 98.7 (30.00-2.76)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.191 , 0.257 0.189 , 0.255	Depositor DCC
R_{free} test set	872 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	64.6	Xtriage
Anisotropy	0.105	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3752	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 34Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.97	2/3784 (0.1%)	1.14	14/5158 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	47	HIS	CE1-NE2	13.43	1.46	1.32
1	X	47	HIS	CG-ND1	5.27	1.44	1.38

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	424	ALA	CA-C-N	-7.67	112.09	119.76
1	X	424	ALA	C-N-CA	-7.67	112.09	119.76
1	X	288	VAL	CB-CA-C	-6.60	103.23	112.14
1	X	232	ARG	CA-C-N	6.39	127.83	119.84
1	X	232	ARG	C-N-CA	6.39	127.83	119.84
1	X	268	SER	N-CA-C	6.14	121.29	113.55
1	X	268	SER	CA-C-N	6.12	132.72	121.70
1	X	268	SER	C-N-CA	6.12	132.72	121.70
1	X	439	ARG	N-CA-C	-5.95	98.13	110.80
1	X	447	CYS	CA-C-N	-5.53	113.38	122.67
1	X	447	CYS	C-N-CA	-5.53	113.38	122.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	431	VAL	CB-CA-C	-5.28	102.36	110.50
1	X	184	PHE	N-CA-C	-5.19	106.20	112.54
1	X	182	VAL	N-CA-C	-5.16	106.77	111.67

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	438	GLN	Peptide
1	X	439	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	3716	0	3706	88	0
2	X	11	0	3	1	0
3	X	25	0	0	1	0
All	All	3752	0	3709	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:438:GLN:HA	1:X:440:TRP:HA	1.55	0.88
1:X:333:ARG:HD3	1:X:344:ILE:CD1	2.04	0.86
1:X:119:GLN:HG3	1:X:119:GLN:O	1.79	0.80
1:X:128:PHE:HB2	1:X:131:ASP:OD2	1.79	0.80
1:X:99:ARG:NH1	1:X:167:LEU:HD12	1.99	0.77
1:X:488:ASN:HB3	1:X:494:LEU:HD21	1.71	0.70
1:X:268:SER:H	1:X:269:LEU:HB2	1.59	0.68
1:X:38:ARG:HH11	1:X:38:ARG:HG3	1.58	0.68
1:X:165:THR:HG23	1:X:166:GLY:H	1.58	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:167:LEU:HD22	1:X:168:PRO:HD2	1.75	0.67
1:X:431:VAL:HG23	1:X:485:LEU:HD13	1.78	0.66
1:X:38:ARG:HG3	1:X:38:ARG:NH1	2.12	0.63
1:X:239:LEU:O	1:X:243:GLU:HB2	1.99	0.62
1:X:333:ARG:HD3	1:X:344:ILE:HD13	1.80	0.62
1:X:198:VAL:HG12	1:X:245:VAL:HA	1.80	0.62
1:X:95:GLU:O	1:X:99:ARG:HG3	2.03	0.59
1:X:39:VAL:HG13	1:X:73:ALA:HA	1.85	0.59
1:X:188:GLN:HE21	1:X:311:ASN:HD22	1.50	0.58
1:X:377:GLN:NE2	1:X:382:ARG:HH11	2.02	0.57
1:X:370:GLN:H	1:X:370:GLN:CD	2.15	0.54
1:X:350:GLU:OE2	1:X:398:LEU:HD13	2.07	0.54
1:X:181:ARG:HG2	1:X:310:MET:CG	2.38	0.54
1:X:431:VAL:CG2	1:X:485:LEU:HD13	2.38	0.54
1:X:82:ALA:HA	1:X:158:ILE:HB	1.90	0.53
1:X:274:LEU:HD23	1:X:297:PRO:HD2	1.92	0.52
1:X:431:VAL:HG21	1:X:493:VAL:HG11	1.91	0.52
1:X:377:GLN:NE2	1:X:382:ARG:HD2	2.26	0.51
1:X:203:MET:HG2	1:X:249:PHE:O	2.11	0.51
1:X:247:SER:HB2	1:X:276:HIS:HB2	1.93	0.51
1:X:354:ILE:O	1:X:354:ILE:HG13	2.09	0.51
1:X:60:VAL:HB	1:X:107:ILE:HG22	1.92	0.51
1:X:308:GLU:OE1	1:X:308:GLU:N	2.37	0.50
1:X:288:VAL:O	1:X:289:LEU:C	2.54	0.49
1:X:451:ARG:O	1:X:452:LEU:C	2.54	0.49
1:X:52:ARG:HB3	1:X:53:PRO:HD2	1.95	0.49
1:X:265:ALA:HB1	1:X:269:LEU:HD12	1.95	0.49
1:X:315:MET:HE3	1:X:316:ARG:O	2.13	0.48
1:X:9:ARG:O	1:X:13:THR:HG23	2.13	0.48
1:X:150:ARG:HA	1:X:154:GLN:HE22	1.78	0.48
1:X:440:TRP:HE3	1:X:441:GLY:H	1.60	0.48
1:X:472:ASP:HA	1:X:475:ARG:HH21	1.79	0.48
1:X:204:PRO:O	1:X:210:GLY:HA3	2.14	0.47
1:X:280:ALA:HA	1:X:303:GLY:O	2.14	0.47
1:X:161:THR:O	1:X:162:SER:CB	2.62	0.47
1:X:404:MET:HE1	1:X:411:ASN:ND2	2.30	0.47
1:X:77:LEU:O	1:X:148:PRO:HG3	2.15	0.47
1:X:404:MET:HE3	1:X:413:HIS:CE1	2.50	0.47
1:X:495:ARG:O	1:X:496:ARG:C	2.58	0.47
1:X:448:VAL:CG2	1:X:456:LEU:HD13	2.44	0.47
1:X:243:GLU:O	1:X:244:GLN:HB2	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:244:GLN:HA	1:X:244:GLN:OE1	2.15	0.46
1:X:162:SER:HA	1:X:168:PRO:HA	1.96	0.46
1:X:271:LEU:HB2	1:X:297:PRO:HG3	1.96	0.46
1:X:302:ASN:O	1:X:314:TYR:HA	2.16	0.46
1:X:413:HIS:HA	1:X:414:PRO:HD2	1.62	0.46
1:X:302:ASN:HB2	1:X:318:PRO:HG3	1.98	0.45
1:X:61:ALA:HB3	1:X:67:VAL:HG21	1.98	0.45
1:X:161:THR:O	1:X:162:SER:HB3	2.16	0.45
1:X:449:VAL:HA	1:X:450:PRO:HD3	1.80	0.45
1:X:308:GLU:HG2	1:X:381:TYR:CE1	2.52	0.45
1:X:38:ARG:O	1:X:42:VAL:HG23	2.16	0.45
1:X:181:ARG:HG2	1:X:310:MET:HG2	1.98	0.45
1:X:240:VAL:HA	1:X:245:VAL:CG2	2.47	0.45
1:X:345:VAL:C	1:X:346:ALA:O	2.60	0.45
1:X:404:MET:HG3	1:X:413:HIS:CE1	2.52	0.45
1:X:18:HIS:O	1:X:31:THR:HA	2.17	0.44
1:X:404:MET:HE2	1:X:411:ASN:HB3	2.00	0.44
1:X:188:GLN:HE21	1:X:311:ASN:ND2	2.13	0.44
1:X:198:VAL:CG1	1:X:245:VAL:HA	2.45	0.43
1:X:269:LEU:HA	1:X:269:LEU:HD23	1.63	0.42
1:X:305:GLY:HA3	2:X:505:34Z:C3	2.49	0.42
1:X:174:PRO:HD3	1:X:363:VAL:HG13	2.00	0.42
1:X:299:GLU:OE2	1:X:316:ARG:HD2	2.19	0.42
1:X:377:GLN:HE21	1:X:382:ARG:HH11	1.66	0.42
1:X:99:ARG:HH12	1:X:167:LEU:HD12	1.81	0.42
1:X:289:LEU:O	1:X:292:VAL:HB	2.19	0.42
1:X:154:GLN:HE21	1:X:175:GLN:HG3	1.84	0.42
1:X:149:GLN:HE21	1:X:149:GLN:HB2	1.69	0.41
1:X:345:VAL:HG23	1:X:346:ALA:O	2.20	0.41
1:X:117:ILE:C	1:X:119:GLN:N	2.79	0.41
1:X:211:PHE:O	1:X:216:VAL:HG23	2.20	0.41
1:X:12:ALA:HB2	1:X:36:ARG:HG3	2.02	0.41
1:X:133:VAL:HA	1:X:138:PRO:HA	2.01	0.40
1:X:172:ILE:HB	1:X:363:VAL:HG22	2.03	0.40
1:X:170:ALA:O	1:X:365:TYR:HA	2.21	0.40
1:X:265:ALA:HA	3:X:507:HOH:O	2.21	0.40
1:X:331:GLU:HB3	1:X:356:ALA:HB2	2.04	0.40
1:X:435:LEU:HD23	1:X:435:LEU:HA	1.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	493/504 (98%)	449 (91%)	34 (7%)	10 (2%)	6	11

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	346	ALA
1	X	439	ARG
1	X	269	LEU
1	X	51	LEU
1	X	437	ASP
1	X	440	TRP
1	X	338	GLY
1	X	358	SER
1	X	472	ASP
1	X	50	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	376/399 (94%)	345 (92%)	31 (8%)	10	20

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	21	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	23	VAL
1	X	29	ARG
1	X	90	SER
1	X	92	GLU
1	X	99	ARG
1	X	102	MET
1	X	103	THR
1	X	119	GLN
1	X	140	SER
1	X	161	THR
1	X	169	LYS
1	X	188	GLN
1	X	229	GLU
1	X	232	ARG
1	X	242	GLN
1	X	243	GLU
1	X	248	LEU
1	X	290	GLU
1	X	294	GLN
1	X	317	GLN
1	X	347	ASN
1	X	355	VAL
1	X	358	SER
1	X	370	GLN
1	X	377	GLN
1	X	401	VAL
1	X	448	VAL
1	X	455	THR
1	X	472	ASP
1	X	484	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	18	HIS
1	X	149	GLN
1	X	154	GLN
1	X	175	GLN
1	X	241	GLN
1	X	242	GLN
1	X	294	GLN
1	X	295	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	311	ASN
1	X	317	GLN
1	X	347	ASN
1	X	377	GLN
1	X	411	ASN
1	X	413	HIS
1	X	484	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	34Z	X	505	-	11,11,11	1.86	1 (9%)	15,15,15	2.88	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	34Z	X	505	-	-	0/4/4/4	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	X	505	34Z	C3-CL3	5.60	1.86	1.73

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	X	505	34Z	C2-C3-C4	-4.85	113.82	120.28
2	X	505	34Z	C5-C6-C1	-4.79	115.68	120.80
2	X	505	34Z	C2-C3-CL3	4.47	125.76	118.45
2	X	505	34Z	C6-C1-C2	4.45	124.40	119.25
2	X	505	34Z	C5-C4-C3	3.73	126.61	120.00
2	X	505	34Z	C3-C4-CL4	-3.26	113.21	120.84
2	X	505	34Z	O1-C-C1	2.13	120.30	114.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	X	505	34Z	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	499/504 (99%)	0.57	25 (5%) 34 34	37, 56, 80, 96	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	25	ALA	4.8
1	X	409	GLY	3.4
1	X	109	VAL	3.3
1	X	269	LEU	2.8
1	X	439	ARG	2.8
1	X	28	LEU	2.7
1	X	244	GLN	2.6
1	X	143	PRO	2.4
1	X	436	ALA	2.4
1	X	51	LEU	2.3
1	X	387	ALA	2.3
1	X	489	ALA	2.3
1	X	165	THR	2.3
1	X	266	GLY	2.3
1	X	349	GLU	2.3
1	X	234	VAL	2.2
1	X	376	LEU	2.2
1	X	334	ILE	2.2
1	X	330	SER	2.2
1	X	128	PHE	2.2
1	X	139	TYR	2.1
1	X	260	ALA	2.1
1	X	312	SER	2.0
1	X	237	LEU	2.0
1	X	240	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	34Z	X	505	11/11	0.93	0.16	52,54,57,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.