



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:00 AM UTC

PDB ID : 3QVK / pdb_00003qvk
Title : Allantoin racemase from *Klebsiella pneumoniae*
Authors : French, J.B.; Neau, D.B.; Ealick, S.E.
Deposited on : 2011-02-25
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	7% 97% .
1	B	245	14% 94% 6%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative hydantoin racemase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	1	0
			1779	1121	319	330	9			
1	B	245	Total	C	N	O	S	0	2	0
			1782	1123	318	332	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	SER	CYS	conflict	UNP A6T9E8
A	184	SER	CYS	conflict	UNP A6T9E8
A	193	GLN	GLU	conflict	UNP A6T9E8
B	79	SER	CYS	conflict	UNP A6T9E8
B	184	SER	CYS	conflict	UNP A6T9E8
B	193	GLN	GLU	conflict	UNP A6T9E8

- Molecule 2 is 1-(2,5-DIOXO-2,5-DIHYDRO-1H-IMIDAZOL-4-YL)UREA (CCD ID: 2AL) (formula: C₄H₄N₄O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 11	C 4	N 4	O 3	0	0
2	B	1	Total 22	C 8	N 8	O 6	0	1

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	156	Total 156	O 156	0	0
3	B	136	Total 136	O 136	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative hydantoin racemase



- Molecule 1: Putative hydantoin racemase



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	124.90Å 124.90Å 126.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	108.16 – 2.00 108.16 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.4 (108.16-2.00) 98.4 (108.16-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.221 , 0.246 0.222 , 0.245	Depositor DCC
R_{free} test set	1975 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 26.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3886	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2AL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1806	0.80	0/2453
1	B	0.43	0/1809	0.78	0/2458
All	All	0.43	0/3615	0.79	0/4911

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1779	0	1811	3	0
1	B	1782	0	1807	9	0
2	A	11	0	4	0	0
2	B	22	0	8	0	0
3	A	156	0	0	0	0
3	B	136	0	0	0	0
All	All	3886	0	3630	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:138[B]:GLN:HE21	1:B:138[B]:GLN:H	1.41	0.67
1:B:138[B]:GLN:HE21	1:B:138[B]:GLN:N	2.01	0.58
1:B:10:ASN:O	1:B:40:PRO:HD3	2.10	0.52
1:A:10:ASN:O	1:A:40:PRO:HD3	2.10	0.52
1:B:53:GLU:OE2	1:B:81:GLY:HA3	2.11	0.51
1:B:161:GLN:HG3	1:B:190:LEU:HD11	1.95	0.49
1:A:144:HIS:HB3	1:A:167:ARG:HD2	1.95	0.48
1:A:53:GLU:OE2	1:A:81:GLY:HA3	2.17	0.44
1:B:106:MET:HE2	1:B:106:MET:HA	2.00	0.44
1:B:11:PRO:HG2	1:B:79:SER:HB2	1.99	0.43
1:B:138[B]:GLN:H	1:B:138[B]:GLN:NE2	2.12	0.43
1:B:17:MET:HE1	1:B:150:VAL:HG12	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/245 (100%)	241 (99%)	3 (1%)	0	100	100
1	B	245/245 (100%)	241 (98%)	4 (2%)	0	100	100
All	All	489/490 (100%)	482 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/183 (98%)	177 (99%)	2 (1%)	65	73
1	B	179/183 (98%)	176 (98%)	3 (2%)	53	60
All	All	358/366 (98%)	353 (99%)	5 (1%)	59	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	167	ARG
1	A	190	LEU
1	B	15	LEU
1	B	167	ARG
1	B	189	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	HIS
1	A	226	HIS
1	B	245	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2AL	A	451	-	9,11,11	1.99	3 (33%)	7,15,15	2.62	2 (28%)
2	2AL	B	451[B]	-	9,11,11	2.12	4 (44%)	7,15,15	2.63	2 (28%)
2	2AL	B	451[A]	-	9,11,11	2.11	4 (44%)	7,15,15	2.60	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2AL	A	451	-	-	4/4/16/16	0/1/1/1
2	2AL	B	451[B]	-	-	4/4/16/16	0/1/1/1
2	2AL	B	451[A]	-	-	2/4/16/16	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	451[B]	2AL	C2-N3	-4.90	1.33	1.40
2	B	451[A]	2AL	C2-N3	-4.83	1.33	1.40
2	A	451	2AL	C2-N3	-4.52	1.34	1.40
2	B	451[A]	2AL	C4-N3	-2.79	1.33	1.38
2	B	451[B]	2AL	C4-N3	-2.76	1.33	1.38
2	A	451	2AL	C4-N3	-2.53	1.34	1.38
2	A	451	2AL	C8-N9	-2.05	1.34	1.40
2	B	451[B]	2AL	C5-N7	-2.04	1.33	1.38
2	B	451[A]	2AL	C8-N9	-2.04	1.34	1.40
2	B	451[B]	2AL	C8-N9	-2.04	1.34	1.40
2	B	451[A]	2AL	C5-N7	-2.03	1.33	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	451[B]	2AL	C5-N7-C8	-5.79	105.78	111.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	451[A]	2AL	C5-N7-C8	-5.72	105.84	111.49
2	A	451	2AL	C5-N7-C8	-5.68	105.89	111.49
2	B	451[B]	2AL	C4-C5-N7	3.62	108.28	104.47
2	B	451[A]	2AL	C4-C5-N7	3.57	108.23	104.47
2	A	451	2AL	C4-C5-N7	3.52	108.18	104.47

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	451	2AL	C5-C4-N3-C2
2	A	451	2AL	N9-C4-N3-C2
2	B	451[A]	2AL	C5-C4-N3-C2
2	B	451[A]	2AL	N9-C4-N3-C2
2	B	451[B]	2AL	C5-C4-N3-C2
2	B	451[B]	2AL	N9-C4-N3-C2
2	A	451	2AL	N1-C2-N3-C4
2	A	451	2AL	O2-C2-N3-C4
2	B	451[B]	2AL	N1-C2-N3-C4
2	B	451[B]	2AL	O2-C2-N3-C4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/245 (100%)	0.78	18 (7%) 21 19	19, 30, 38, 40	1 (0%)
1	B	245/245 (100%)	0.94	35 (14%) 6 5	14, 30, 44, 50	2 (0%)
All	All	490/490 (100%)	0.86	53 (10%) 11 10	14, 30, 40, 50	3 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	SER	5.0
1	B	155	ASP	4.8
1	B	150	VAL	4.2
1	B	3	SER	4.1
1	A	247	PHE	4.0
1	A	246	PRO	4.0
1	B	189	THR	3.9
1	B	190	LEU	3.6
1	B	149	PRO	3.6
1	A	192[A]	GLN	3.5
1	A	189	THR	3.5
1	B	151	LEU	3.2
1	A	240	GLN	3.1
1	A	242	GLN	3.1
1	B	173	LYS	3.1
1	A	186	GLY	3.0
1	B	69	GLU	3.0
1	B	187	MET	3.0
1	B	148	LEU	2.9
1	B	154	GLU	2.8
1	A	187	MET	2.8
1	B	191	ALA	2.7
1	A	147	ASP	2.7
1	B	167	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	158	GLY	2.6
1	B	196	ARG	2.6
1	B	70	GLN	2.6
1	A	15	LEU	2.5
1	B	186	GLY	2.5
1	B	226	HIS	2.5
1	B	15	LEU	2.5
1	B	153	LEU	2.5
1	A	199	ARG	2.4
1	B	163	LYS	2.3
1	B	157	SER	2.3
1	B	38	VAL	2.2
1	A	190	LEU	2.2
1	B	147	ASP	2.2
1	B	188	ALA	2.2
1	B	165	ARG	2.2
1	A	159	LEU	2.2
1	A	191	ALA	2.2
1	B	171	ALA	2.2
1	A	129	HIS	2.2
1	A	19	GLU	2.2
1	B	152	ALA	2.2
1	B	237	LEU	2.1
1	B	169	ILE	2.1
1	B	14	SER	2.1
1	B	19	GLU	2.1
1	B	156	GLY	2.1
1	B	239	GLY	2.0
1	A	196	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	2AL	B	451[A]	11/11	0.58	0.21	40,41,41,41	11
2	2AL	B	451[B]	11/11	0.58	0.21	43,43,43,44	11
2	2AL	A	451	11/11	0.83	0.14	39,40,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.