



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 07:30 AM UTC

PDB ID : 7QVJ / pdb_00007qvj
Title : ESTROGEN RECEPTOR ALPHA IN COMPLEX WITH COMPOUND 29
Authors : Breed, J.
Deposited on : 2022-01-21
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1054 (1.68-1.68)
Clashscore	190562	1078 (1.68-1.68)
Ramachandran outliers	187476	1068 (1.68-1.68)
Sidechain outliers	187428	1067 (1.68-1.68)
RSRZ outliers	180081	1055 (1.68-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	9% 85% 10% 6%
1	B	252	15% 87% 7% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	238	Total	C	N	O	S	0	3	0
			1823	1163	322	324	14			
1	B	239	Total	C	N	O	S	0	1	0
			1795	1152	309	320	14			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	303	GLY	-	expression tag	UNP P03372
A	304	SER	-	expression tag	UNP P03372
A	305	HIS	-	expression tag	UNP P03372
A	306	MET	-	expression tag	UNP P03372
A	381	SER	CYS	engineered mutation	UNP P03372
A	417	SER	CYS	engineered mutation	UNP P03372
A	530	SER	CYS	engineered mutation	UNP P03372
A	536	SER	LEU	engineered mutation	UNP P03372
B	303	GLY	-	expression tag	UNP P03372
B	304	SER	-	expression tag	UNP P03372
B	305	HIS	-	expression tag	UNP P03372
B	306	MET	-	expression tag	UNP P03372
B	381	SER	CYS	engineered mutation	UNP P03372
B	417	SER	CYS	engineered mutation	UNP P03372
B	530	SER	CYS	engineered mutation	UNP P03372
B	536	SER	LEU	engineered mutation	UNP P03372

- Molecule 2 is 2,2-bis(fluoranyl)-3-[(1 {R},3 {R})-1-[6-fluoranyl-3-[2-(3-fluoranylpropylamino)ethoxy]-2-methyl-phenyl]-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-2-yl]propan-1-ol (CCD ID: H09) (formula: C₂₇H₃₃F₄N₃O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 36	C 27	F 4	N 3	O 2	0	0
2	B	1	Total 36	C 27	F 4	N 3	O 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	37	Total O 37 37	0	0
3	B	28	Total O 28 28	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Estrogen receptor



• Molecule 1: Estrogen receptor



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.90Å 51.79Å 85.51Å 90.00° 93.86° 90.00°	Depositor
Resolution (Å)	43.91 – 1.68 43.91 – 1.68	Depositor EDS
% Data completeness (in resolution range)	95.3 (43.91-1.68) 95.3 (43.91-1.68)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 1.68Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.205 , 0.216 0.213 , 0.225	Depositor DCC
R_{free} test set	2466 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3755	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: H09

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/1856	0.96	1/2519 (0.0%)
1	B	0.63	0/1825	0.94	0/2477
All	All	0.63	0/3681	0.95	1/4996 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	336	PRO	N-CA-CB	6.30	109.93	103.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	467	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1823	0	1774	19	1
1	B	1795	0	1737	10	1
2	A	36	0	0	2	0
2	B	36	0	0	0	0
3	A	37	0	0	0	0
3	B	28	0	0	1	0
All	All	3755	0	3511	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359[B]:ASN:OD1	1:A:547:HIS:CD2	2.34	0.79
1:A:533:VAL:O	2:A:601:H09:C24	2.40	0.69
1:B:368:VAL:HG22	3:B:702:HOH:O	1.92	0.69
1:A:454:LEU:O	1:A:458:VAL:CG1	2.47	0.62
1:B:358:ILE:HD13	1:B:543:MET:HB3	1.85	0.58
1:A:358:ILE:HD13	1:A:379:LEU:HD21	1.85	0.58
1:A:359[B]:ASN:OD1	1:A:547:HIS:NE2	2.36	0.57
1:A:522:MET:HE1	1:A:534:VAL:CG1	2.37	0.54
1:A:316:VAL:HG21	1:A:489:LEU:HD21	1.89	0.54
1:A:454:LEU:O	1:A:458:VAL:HG13	2.08	0.53
1:B:396:MET:HE3	1:B:436:ARG:HA	1.91	0.53
1:A:522:MET:HE1	1:A:534:VAL:HG11	1.92	0.52
1:B:352:ARG:O	1:B:355:VAL:HG12	2.10	0.52
1:A:362:LYS:HD3	1:A:547:HIS:ND1	2.26	0.51
1:A:310:LEU:O	1:A:481:LYS:HE3	2.11	0.51
1:A:359[B]:ASN:OD1	1:A:547:HIS:HD2	1.93	0.50
1:A:383:TRP:CZ2	1:A:535:PRO:HD2	2.47	0.49
1:A:533:VAL:O	2:A:601:H09:C23	2.62	0.48
1:A:476[B]:HIS:HD2	1:B:434:ARG:HH22	1.62	0.47
1:A:476[B]:HIS:ND1	1:A:476[B]:HIS:C	2.73	0.47
1:B:330:GLU:C	1:B:331:TYR:O	2.59	0.46
1:B:379:LEU:HD12	1:B:544:LEU:HD11	1.98	0.46
1:B:315:MET:HE3	1:B:481:LYS:HG2	1.99	0.45
1:B:330:GLU:O	1:B:331:TYR:O	2.35	0.45
1:A:403:LEU:HD13	1:A:409:LEU:HD13	2.01	0.43
1:A:396:MET:O	1:A:436:ARG:HD3	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:362:LYS:HD3	1:A:547:HIS:CE1	2.54	0.42
1:B:330:GLU:O	1:B:331:TYR:C	2.63	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359[B]:ASN:ND2	1:B:328:TYR:OH[3_545]	1.36	0.84

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/252 (92%)	231 (99%)	2 (1%)	0	100	100
1	B	234/252 (93%)	227 (97%)	5 (2%)	2 (1%)	14	3
All	All	467/504 (93%)	458 (98%)	7 (2%)	2 (0%)	30	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	331	TYR
1	B	469	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/226 (83%)	186 (99%)	2 (1%)	65	47
1	B	179/226 (79%)	177 (99%)	2 (1%)	65	47
All	All	367/452 (81%)	363 (99%)	4 (1%)	65	47

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	417	SER
1	A	422	VAL
1	B	421	MET
1	B	425	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	488	HIS
1	B	513	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	H09	A	601	-	38,39,39	0.27	0	51,56,56	0.54	0
2	H09	B	601	-	38,39,39	0.31	0	51,56,56	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	H09	A	601	-	-	6/18/36/36	0/4/4/4
2	H09	B	601	-	-	3/18/36/36	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	H09	F-C19-C20-O
2	A	601	H09	F1-C19-C20-O
2	A	601	H09	N2-C24-C25-C26
2	B	601	H09	F-C19-C20-O
2	B	601	H09	F1-C19-C20-O
2	A	601	H09	O1-C22-C23-N2
2	A	601	H09	C22-C23-N2-C24
2	B	601	H09	C23-C22-O1-C2
2	A	601	H09	C25-C24-N2-C23


There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	H09	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	238/252 (94%)	0.80	22 (9%) 14 17	15, 44, 77, 102	4 (1%)
1	B	239/252 (94%)	1.01	39 (16%) 4 5	18, 45, 88, 118	2 (0%)
All	All	477/504 (94%)	0.91	61 (12%) 7 9	15, 45, 82, 118	6 (1%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	465	THR	6.5
1	B	336	PRO	6.4
1	B	469	LEU	6.3
1	A	461	PHE	6.1
1	A	336	PRO	5.4
1	B	466	LEU	5.3
1	B	533	VAL	5.1
1	B	332	ASP	4.7
1	A	334	THR	4.6
1	B	528	MET	4.5
1	B	550	HIS	4.4
1	A	458	VAL	4.4
1	A	533	VAL	4.4
1	A	552	PRO	4.0
1	A	469	LEU	3.8
1	A	468	SER	3.7
1	B	425	PHE	3.7
1	A	340	ALA	3.7
1	B	468	SER	3.5
1	A	413	ASN	3.4
1	B	413	ASN	3.4
1	A	459	TYR	3.3
1	B	308	LEU	3.3
1	B	464	SER	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	341	SER	3.2
1	B	462	LEU	3.1
1	B	306	MET	3.1
1	B	415	GLY	3.1
1	B	467	LYS	3.0
1	B	513	HIS	3.0
1	B	470	GLU	3.0
1	B	424	ILE	2.9
1	B	331	TYR	2.9
1	A	537	TYR	2.9
1	B	307	ALA	2.8
1	B	526	TYR	2.8
1	B	534	VAL	2.7
1	A	333	PRO	2.6
1	B	337	PHE	2.6
1	A	470	GLU	2.6
1	A	331	TYR	2.6
1	A	528	MET	2.5
1	B	549	LEU	2.5
1	A	339	GLU	2.5
1	A	497	LEU	2.4
1	A	460	THR	2.4
1	B	326	ILE	2.4
1	B	547	HIS	2.4
1	B	463	SER	2.4
1	B	310	LEU	2.4
1	B	435	PHE	2.4
1	A	342	MET	2.4
1	B	472	LYS	2.2
1	B	459	TYR	2.1
1	B	476	HIS	2.1
1	B	305	HIS	2.1
1	A	343	MET	2.1
1	A	332	ASP	2.1
1	B	475	ILE	2.1
1	B	461	PHE	2.0
1	B	546	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	H09	A	601	36/36	0.88	0.11	38,43,65,70	0
2	H09	B	601	36/36	0.88	0.12	36,42,70,76	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around H09 A 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around H09 B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.