



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:48 PM UTC

PDB ID : 2QW6 / pdb_00002qw6
Title : Crystal structure of the C-terminal domain of an AAA ATPase from *Enterococcus faecium* DO
Authors : Bonanno, J.B.; Rutter, M.; Bain, K.T.; Ozyurt, S.; Smith, D.; Wasserman, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2007-08-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	112	4% 66% 9% • 24%
1	B	112	% 64% 11% • 24%
1	C	112	4% 59% 15% • 26%
1	D	112	% 62% 12% • 23%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2593 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AAA ATPase, central region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	85	Total	C	N	O	S	0	0	0
			633	401	110	119	3			
1	B	85	Total	C	N	O	S	0	0	0
			633	401	110	119	3			
1	C	83	Total	C	N	O	S	0	0	0
			620	393	108	116	3			
1	D	86	Total	C	N	O	S	0	0	0
			642	407	112	120	3			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	227	MET	-	expression tag	UNP Q3XY27
A	228	SER	-	expression tag	UNP Q3XY27
A	229	LEU	-	expression tag	UNP Q3XY27
A	331	GLU	-	expression tag	UNP Q3XY27
A	332	GLY	-	expression tag	UNP Q3XY27
A	333	HIS	-	expression tag	UNP Q3XY27
A	334	HIS	-	expression tag	UNP Q3XY27
A	335	HIS	-	expression tag	UNP Q3XY27
A	336	HIS	-	expression tag	UNP Q3XY27
A	337	HIS	-	expression tag	UNP Q3XY27
A	338	HIS	-	expression tag	UNP Q3XY27
B	227	MET	-	expression tag	UNP Q3XY27
B	228	SER	-	expression tag	UNP Q3XY27
B	229	LEU	-	expression tag	UNP Q3XY27
B	331	GLU	-	expression tag	UNP Q3XY27
B	332	GLY	-	expression tag	UNP Q3XY27
B	333	HIS	-	expression tag	UNP Q3XY27
B	334	HIS	-	expression tag	UNP Q3XY27
B	335	HIS	-	expression tag	UNP Q3XY27
B	336	HIS	-	expression tag	UNP Q3XY27
B	337	HIS	-	expression tag	UNP Q3XY27

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	338	HIS	-	expression tag	UNP Q3XY27
C	227	MET	-	expression tag	UNP Q3XY27
C	228	SER	-	expression tag	UNP Q3XY27
C	229	LEU	-	expression tag	UNP Q3XY27
C	331	GLU	-	expression tag	UNP Q3XY27
C	332	GLY	-	expression tag	UNP Q3XY27
C	333	HIS	-	expression tag	UNP Q3XY27
C	334	HIS	-	expression tag	UNP Q3XY27
C	335	HIS	-	expression tag	UNP Q3XY27
C	336	HIS	-	expression tag	UNP Q3XY27
C	337	HIS	-	expression tag	UNP Q3XY27
C	338	HIS	-	expression tag	UNP Q3XY27
D	227	MET	-	expression tag	UNP Q3XY27
D	228	SER	-	expression tag	UNP Q3XY27
D	229	LEU	-	expression tag	UNP Q3XY27
D	331	GLU	-	expression tag	UNP Q3XY27
D	332	GLY	-	expression tag	UNP Q3XY27
D	333	HIS	-	expression tag	UNP Q3XY27
D	334	HIS	-	expression tag	UNP Q3XY27
D	335	HIS	-	expression tag	UNP Q3XY27
D	336	HIS	-	expression tag	UNP Q3XY27
D	337	HIS	-	expression tag	UNP Q3XY27
D	338	HIS	-	expression tag	UNP Q3XY27


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	22	Total O 22 22	0	0
2	B	17	Total O 17 17	0	0
2	C	13	Total O 13 13	0	0
2	D	13	Total O 13 13	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: AAA ATPase, central region




- Molecule 1: AAA ATPase, central region



- Molecule 1: AAA ATPase, central region



- Molecule 1: AAA ATPase, central region



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	62.87Å 87.02Å 87.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 20.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.30) 99.8 (20.00-2.30)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.94 (at 2.31Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.224 , 0.275 (Not available) , 0.233	Depositor DCC
R_{free} test set	1128 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.891	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.092 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2593	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6842e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/640	1.02	0/870
1	B	0.87	0/640	1.01	1/870 (0.1%)
1	C	0.91	0/626	1.06	0/850
1	D	0.93	0/649	1.05	1/881 (0.1%)
All	All	0.90	0/2555	1.04	2/3471 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	270	GLY	N-CA-C	5.23	122.84	115.30
1	D	256	VAL	N-CA-C	5.10	115.71	110.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	633	0	651	8	0
1	B	633	0	651	8	0
1	C	620	0	639	12	0
1	D	642	0	664	12	0
2	A	22	0	0	3	0
2	B	17	0	0	0	0
2	C	13	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	13	0	0	0	0
All	All	2593	0	2605	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:ALA:O	1:A:268:GLU:HG3	1.73	0.89
1:D:301:ALA:CB	1:D:320:VAL:HG21	2.09	0.83
1:C:287:ILE:HD13	1:C:324:LEU:HD22	1.65	0.77
1:D:311:PRO:O	1:D:314:ARG:HG2	1.89	0.72
1:A:287:ILE:HD12	1:A:324:LEU:HD22	1.75	0.68
1:D:317:LEU:O	1:D:321:VAL:HG12	1.95	0.66
1:C:301:ALA:CB	1:C:320:VAL:CG2	2.78	0.62
1:C:248:GLN:HG3	1:C:282:ILE:HD11	1.83	0.60
1:C:301:ALA:CB	1:C:320:VAL:HG21	2.34	0.57
1:D:301:ALA:HB3	1:D:320:VAL:HG21	1.87	0.57
1:C:264:ALA:O	1:C:268:GLU:HG3	2.05	0.57
1:A:278:ARG:HD2	2:A:23:HOH:O	2.08	0.53
1:A:248:GLN:HG3	1:A:282:ILE:HD11	1.91	0.52
1:D:301:ALA:CB	1:D:320:VAL:CG2	2.83	0.51
1:B:301:ALA:CB	1:B:320:VAL:HG21	2.41	0.51
1:D:243:VAL:HG22	1:D:266:LEU:HD21	1.95	0.49
1:B:296:ALA:HB2	1:C:297:ARG:HG3	1.96	0.48
1:A:241:TYR:N	2:A:75:HOH:O	2.47	0.47
1:B:301:ALA:HB3	1:B:320:VAL:HG21	1.95	0.47
1:A:248:GLN:NE2	2:A:89:HOH:O	2.50	0.45
1:C:301:ALA:HB1	1:C:320:VAL:CG2	2.48	0.44
1:D:278:ARG:HA	1:D:278:ARG:HD3	1.78	0.43
1:C:267:VAL:HG13	1:C:310:LEU:HD22	1.99	0.43
1:C:281:VAL:HG13	1:D:315:ILE:HD13	1.98	0.43
1:C:302:VAL:O	1:C:306:GLU:HG3	2.16	0.43
1:A:312:GLU:OE1	1:D:277:ARG:NH2	2.51	0.43
1:B:267:VAL:HG13	1:B:310:LEU:HD22	2.00	0.42
1:B:278:ARG:HA	1:B:278:ARG:HD3	1.83	0.42
1:B:248:GLN:O	1:B:252:ARG:HG3	2.21	0.41
1:D:301:ALA:HB2	1:D:320:VAL:CG2	2.50	0.41
1:B:284:TYR:O	1:C:319:ASP:OD1	2.38	0.41
1:D:310:LEU:HD23	1:D:313:ALA:HB3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281:VAL:HG13	1:B:315:ILE:HD13	2.01	0.41
1:C:301:ALA:HB3	1:C:320:VAL:HG21	2.02	0.41
1:D:263:LEU:HD22	1:D:321:VAL:HG11	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	81/112 (72%)	81 (100%)	0	0	100	100
1	B	81/112 (72%)	81 (100%)	0	0	100	100
1	C	79/112 (70%)	78 (99%)	1 (1%)	0	100	100
1	D	82/112 (73%)	80 (98%)	2 (2%)	0	100	100
All	All	323/448 (72%)	320 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	66/87 (76%)	64 (97%)	2 (3%)	36	53
1	B	66/87 (76%)	63 (96%)	3 (4%)	24	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	64/87 (74%)	62 (97%)	2 (3%)	35	52
1	D	67/87 (77%)	64 (96%)	3 (4%)	24	37
All	All	263/348 (76%)	253 (96%)	10 (4%)	29	44

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	278	ARG
1	A	319	ASP
1	B	319	ASP
1	B	320	VAL
1	B	322	VAL
1	C	278	ARG
1	C	326	LEU
1	D	243	VAL
1	D	321	VAL
1	D	326	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	85/112 (75%)	0.17	4 (4%) 36 38	28, 37, 53, 61	0
1	B	85/112 (75%)	0.22	1 (1%) 76 77	31, 39, 52, 58	0
1	C	83/112 (74%)	0.38	5 (6%) 27 29	31, 40, 53, 63	0
1	D	86/112 (76%)	0.14	1 (1%) 76 77	26, 37, 48, 55	0
All	All	339/448 (75%)	0.22	11 (3%) 50 52	26, 39, 53, 63	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	287	ILE	4.0
1	A	287	ILE	3.4
1	A	268	GLU	3.3
1	A	328	PRO	3.1
1	B	287	ILE	3.0
1	C	286	ASP	2.5
1	D	319	ASP	2.5
1	C	291	ASN	2.5
1	C	326	LEU	2.1
1	A	326	LEU	2.1
1	C	268	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.