



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:21 AM UTC

PDB ID : 2QWA / pdb_00002qwa
Title : THE X-RAY STRUCTURE OF A DRUG RESISTANT VARIANT R292K
OF TERN N9 INFLUENZA VIRUS NEURAMINIDASE
Authors : Varghese, J.N.
Deposited on : 1998-04-07
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	388	 86% 14% .
2	B	7	 100%
3	C	2	 100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NEURAMINIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	22	0	0
			3065	1914	536	592	23			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	292	LYS	ARG	engineered mutation	UNP P03472

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	399	Total	O	0	0
			399	399		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: NEURAMINIDASE

Chain A:  86% 14%




- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	181.40 Å 181.40 Å 181.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.70	Depositor
% Data completeness (in resolution range)	87.0 (6.00-1.70)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
Refinement program	X-PLOR 3.0	Depositor
R, R_{free}	0.171 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3591	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MAN, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	1/3148 (0.0%)	1.05	16/4287 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	294	ASN	CG-OD1	5.26	1.33	1.23

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	GLN	N-CA-C	11.90	123.81	111.07
1	A	299	ASN	N-CA-C	-7.65	99.20	110.52
1	A	319	SER	N-CA-C	7.00	117.65	109.60
1	A	297	GLY	N-CA-C	6.91	122.49	111.64
1	A	229	GLU	N-CA-C	6.37	119.02	110.35
1	A	434	ASP	N-CA-C	6.30	120.58	113.02
1	A	125	ASP	N-CA-C	-6.08	101.09	110.58
1	A	245	SER	N-CA-C	5.85	117.84	110.24
1	A	180	SER	N-CA-C	5.70	116.92	108.60
1	A	277	GLU	N-CA-C	5.45	118.76	111.24
1	A	151	ASP	N-CA-C	5.39	119.02	112.23
1	A	323	THR	N-CA-C	5.35	120.49	112.94
1	A	204	VAL	CB-CA-C	5.25	118.22	110.77
1	A	113	ASP	N-CA-C	5.24	117.73	109.39
1	A	165	SER	N-CA-C	-5.10	102.74	110.39
1	A	305	ILE	N-CA-C	5.00	115.17	108.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3065	0	2893	32	0
2	B	83	0	70	0	0
3	C	28	0	24	0	0
4	A	14	0	13	0	0
5	A	2	0	0	0	0
6	A	399	0	0	5	4
All	All	3591	0	3000	32	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:ARG:HD3	1:A:209:ARG:NH2	1.82	0.93
1:A:87:LEU:H	1:A:233:HIS:HD2	1.24	0.84
1:A:168:THR:H	1:A:170:ASN:HD21	1.26	0.81
1:A:170:ASN:HB3	6:A:1317(P):HOH:O	1.88	0.74
1:A:445:SER:HB2	6:A:1383(S):HOH:O	1.88	0.73
1:A:97:TRP:H	1:A:395:GLN:HE22	1.38	0.71
1:A:98:HIS:HE1	1:A:419:ARG:HH11	1.38	0.68
1:A:98:HIS:CE1	1:A:419:ARG:HH11	2.15	0.65
1:A:172:ARG:HD3	1:A:209:ARG:HH22	1.68	0.57
1:A:87:LEU:H	1:A:233:HIS:CD2	2.14	0.57
1:A:82:ARG:N	6:A:1179(G):HOH:O	2.38	0.57
1:A:168:THR:H	1:A:170:ASN:ND2	2.00	0.55
1:A:172:ARG:CD	1:A:209:ARG:NH2	2.64	0.54
1:A:221:ASN:HB3	1:A:244:GLY:HA2	1.90	0.53
1:A:144:HIS:HD2	6:A:1036(W):HOH:O	1.94	0.50
1:A:98:HIS:HD2	1:A:99:ILE:O	1.97	0.47
1:A:228:SER:HB3	1:A:350:LYS:HE2	1.97	0.47
1:A:295:TRP:CG	1:A:296:GLN:HG3	2.50	0.46
1:A:238:PRO:HA	1:A:255:TYR:O	2.16	0.46
1:A:174:GLU:OE1	1:A:209:ARG:NH2	2.49	0.45
1:A:170:ASN:HD22	1:A:170:ASN:H	1.65	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:320:PRO:HG2	1:A:389:LYS:HE2	1.99	0.44
1:A:410:PHE:HZ	6:A:1383(S):HOH:O	2.00	0.44
1:A:121:TYR:CG	1:A:228:SER:HA	2.54	0.42
1:A:172:ARG:HD3	1:A:209:ARG:CZ	2.45	0.42
1:A:367:SER:HB2	1:A:400:ASN:HD21	1.84	0.42
1:A:188:THR:HG22	1:A:207:TYR:CZ	2.55	0.42
1:A:84:PHE:CE1	1:A:187:LYS:HD2	2.55	0.42
1:A:168:THR:OG1	1:A:170:ASN:ND2	2.53	0.42
1:A:324:ASP:O	1:A:327:ARG:HD3	2.19	0.41
1:A:247:THR:HG22	1:A:346:ASN:HB3	2.02	0.41
1:A:292:LYS:HE2	1:A:294:ASN:HB2	2.04	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1362(R):HOH:O	6:A:1365(R):HOH:O[31_555]	0.23	1.97
6:A:1375(R):HOH:O	6:A:1378(R):HOH:O[48_555]	0.44	1.76
6:A:1321(Q):HOH:O	6:A:1322(Q):HOH:O[48_555]	0.55	1.65
6:A:1278(N):HOH:O	6:A:1278(N):HOH:O[48_555]	0.71	1.49

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/388 (100%)	367 (95%)	19 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/341 (100%)	338 (99%)	3 (1%)	70	62

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	ASN
1	A	204	VAL
1	A	220	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	HIS
1	A	144	HIS
1	A	170	ASN
1	A	221	ASN
1	A	233	HIS
1	A	296	GLN
1	A	381	ASN
1	A	395	GLN
1	A	400	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	1.23	2 (14%)	17,19,21	1.23	1 (5%)
2	NAG	B	2	2	14,14,15	1.19	1 (7%)	17,19,21	1.49	3 (17%)
2	BMA	B	3	2	11,11,12	1.16	1 (9%)	15,15,17	0.80	0
2	MAN	B	4	2	11,11,12	1.60	2 (18%)	15,15,17	1.52	4 (26%)
2	MAN	B	5	2	11,11,12	1.01	0	15,15,17	1.18	2 (13%)
2	MAN	B	6	2	11,11,12	1.91	4 (36%)	15,15,17	1.13	1 (6%)
2	MAN	B	7	2	11,11,12	1.04	0	15,15,17	0.80	1 (6%)
3	NAG	C	1	3,1	14,14,15	1.18	1 (7%)	17,19,21	1.58	5 (29%)
3	NAG	C	2	3	14,14,15	1.69	3 (21%)	17,19,21	4.34	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	0/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	MAN	B	7	2	-	0/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	1/6/23/26	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	NAG	C4-C5	4.25	1.62	1.53
2	B	6	MAN	C4-C5	4.13	1.61	1.53
2	B	4	MAN	C2-C3	3.23	1.57	1.52
3	C	2	NAG	C1-C2	-3.22	1.48	1.52
2	B	1	NAG	C1-C2	3.11	1.56	1.52
2	B	6	MAN	O5-C5	3.08	1.49	1.43
3	C	1	NAG	O5-C5	2.86	1.49	1.43
2	B	1	NAG	C4-C5	2.73	1.58	1.53
2	B	6	MAN	C2-C3	2.43	1.56	1.52
2	B	6	MAN	C4-C3	2.38	1.58	1.52
2	B	4	MAN	O5-C5	2.37	1.48	1.43
2	B	2	NAG	O5-C1	-2.11	1.40	1.43
2	B	3	BMA	C4-C3	2.04	1.57	1.52
3	C	2	NAG	C4-C3	2.02	1.57	1.52

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C2-N2-C7	14.10	141.79	122.90
3	C	2	NAG	C4-C3-C2	-6.62	101.31	111.02
3	C	2	NAG	C8-C7-N2	-5.51	106.97	116.12
3	C	2	NAG	O7-C7-C8	3.78	128.78	122.05
2	B	1	NAG	C1-C2-N2	-3.14	105.48	110.43
2	B	2	NAG	C4-C3-C2	-3.01	106.61	111.02
3	C	1	NAG	O6-C6-C5	-2.92	101.39	111.33
3	C	2	NAG	C1-O5-C5	2.80	115.94	112.19
2	B	4	MAN	O2-C2-C1	-2.76	102.90	109.22
2	B	4	MAN	C1-O5-C5	2.72	115.83	112.19
3	C	2	NAG	O3-C3-C4	2.69	116.72	110.38
2	B	6	MAN	C1-O5-C5	2.67	115.76	112.19
2	B	4	MAN	O6-C6-C5	-2.65	102.31	111.33
2	B	5	MAN	O4-C4-C3	-2.60	104.24	110.38
3	C	1	NAG	C4-C3-C2	-2.56	107.27	111.02
2	B	5	MAN	C2-C3-C4	-2.51	106.45	110.86
2	B	4	MAN	C3-C4-C5	-2.48	105.74	110.23
3	C	1	NAG	C3-C4-C5	2.32	114.44	110.23
2	B	7	MAN	O4-C4-C3	-2.28	105.01	110.38
3	C	1	NAG	C6-C5-C4	-2.26	107.46	113.02
3	C	1	NAG	C1-O5-C5	2.23	115.18	112.19
2	B	2	NAG	O7-C7-C8	-2.19	118.15	122.05
3	C	2	NAG	O5-C1-C2	-2.14	107.98	111.29
2	B	2	NAG	C8-C7-N2	2.02	119.46	116.12

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	2	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	478(A)	1	14,14,15	1.51	1 (7%)	17,19,21	1.87	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	478(A)	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	478(A)	NAG	C1-C2	-4.76	1.45	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	478(A)	NAG	C3-C4-C5	-3.53	103.83	110.23
4	A	478(A)	NAG	C2-N2-C7	-3.32	118.45	122.90
4	A	478(A)	NAG	C8-C7-N2	2.93	120.98	116.12
4	A	478(A)	NAG	O7-C7-C8	-2.85	116.98	122.05
4	A	478(A)	NAG	O4-C4-C5	2.43	115.31	109.32
4	A	478(A)	NAG	O5-C5-C4	-2.02	105.91	110.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.