



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 03:05 AM UTC

PDB ID : 8QWQ / pdb_00008qwq
BMRB ID : 34872
Title : Magic-angle spinning NMR Structure of Opa60 in Lipid Bilayers
Authors : Forster, M.C.; Andreas, L.B.
Deposited on : 2023-10-19

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLID-STATE NMR

The overall completeness of chemical shifts assignment is 32%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	252	 29% • 12% 56%

2 Ensemble composition and analysis

This entry contains 10 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *target function*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:10-A:17, A:55-A:61, A:66-A:74, A:115-A:125, A:132-A:143, A:192-A:201, A:207-A:218, A:227-A:238 (81)	0.67	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 6, 7, 8
2	5, 10
Single-model clusters	3; 9

3 Entry composition


There is only 1 type of molecule in this entry. The entry contains 1755 atoms, of which 871 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Opacity protein opA60 (Fragment).

Mol	Chain	Residues	Atoms					Trace
1	A	111	Total	C	H	N	O	0
			1755	568	871	164	152	

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q04884
A	240	LYS	-	expression tag	UNP Q04884
A	241	LEU	-	expression tag	UNP Q04884
A	242	ALA	-	expression tag	UNP Q04884
A	243	ALA	-	expression tag	UNP Q04884
A	244	ALA	-	expression tag	UNP Q04884
A	245	LEU	-	expression tag	UNP Q04884
A	246	GLU	-	expression tag	UNP Q04884
A	247	HIS	-	expression tag	UNP Q04884
A	248	HIS	-	expression tag	UNP Q04884
A	249	HIS	-	expression tag	UNP Q04884
A	250	HIS	-	expression tag	UNP Q04884
A	251	HIS	-	expression tag	UNP Q04884
A	252	HIS	-	expression tag	UNP Q04884



4.2.5 Score per residue for model 5

- Molecule 1: Opacity protein opA60 (Fragment)


Chain A: 24% 8% 12% 56%




4.2.6 Score per residue for model 6

- Molecule 1: Opacity protein opA60 (Fragment)


Chain A: 23% 9% 12% 56%





4.2.10 Score per residue for model 10

- Molecule 1: Opacity protein opA60 (Fragment)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 2000 calculated structures, 10 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	405
Number of shifts mapped to atoms	405
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	32%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	635	624	623	4±2
All	All	6350	6240	6230	42

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:211:ALA:HB2	1:A:233:LEU:HD12	0.91	1.38	8	1
1:A:12:VAL:HG22	1:A:59:VAL:HG23	0.88	1.46	4	3
1:A:12:VAL:HG12	1:A:59:VAL:HG13	0.65	1.68	6	1
1:A:196:ILE:HG22	1:A:214:ARG:HB3	0.60	1.73	1	2
1:A:140:VAL:HG12	1:A:193:LEU:CB	0.60	2.26	9	1
1:A:14:ALA:HB3	1:A:235:VAL:HG22	0.59	1.73	3	1
1:A:12:VAL:HG22	1:A:59:VAL:CG2	0.59	2.26	4	2
1:A:56:ARG:HG3	1:A:73:ALA:HB2	0.58	1.74	2	3
1:A:136:ILE:HG22	1:A:197:ALA:HB1	0.58	1.75	9	1
1:A:196:ILE:HG22	1:A:214:ARG:CB	0.56	2.30	1	1
1:A:17:ALA:HB2	1:A:232:SER:HB3	0.55	1.79	4	1
1:A:136:ILE:HG22	1:A:197:ALA:CB	0.55	2.31	9	1
1:A:14:ALA:HB3	1:A:235:VAL:CG2	0.55	2.31	3	1
1:A:70:ALA:HB1	1:A:119:GLY:O	0.51	2.05	2	3

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:12:VAL:HG12	1:A:59:VAL:HG23	0.51	1.81	9	1
1:A:12:VAL:HG12	1:A:59:VAL:CG2	0.50	2.35	9	1
1:A:56:ARG:HB3	1:A:73:ALA:HB2	0.49	1.84	10	1
1:A:118:LEU:HD13	1:A:119:GLY:N	0.48	2.23	3	1
1:A:123:ILE:HG13	1:A:196:ILE:HD11	0.47	1.86	8	1
1:A:211:ALA:CB	1:A:233:LEU:HD12	0.47	2.26	8	1
1:A:123:ILE:HG23	1:A:135:TYR:CD2	0.47	2.45	4	1
1:A:140:VAL:HG12	1:A:193:LEU:HB2	0.46	1.87	9	1
1:A:12:VAL:HG23	1:A:59:VAL:HG12	0.46	1.87	7	1
1:A:197:ALA:HB3	1:A:213:TYR:CE1	0.46	2.45	5	2
1:A:135:TYR:C	1:A:136:ILE:HD12	0.46	2.35	7	1
1:A:56:ARG:CG	1:A:73:ALA:HB2	0.46	2.41	1	1
1:A:12:VAL:HG13	1:A:59:VAL:HG22	0.44	1.87	10	1
1:A:59:VAL:O	1:A:70:ALA:HB3	0.44	2.13	9	1
1:A:134:PRO:HA	1:A:199:VAL:HG12	0.42	1.91	9	2
1:A:12:VAL:HG23	1:A:59:VAL:CG1	0.42	2.45	7	2
1:A:12:VAL:HG12	1:A:59:VAL:CB	0.40	2.47	9	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	76/252 (30%)	75±1 (99±1%)	1±1 (1±1%)	0±0 (0±1%)	31	76
All	All	760/2520 (30%)	751 (99%)	6 (1%)	3 (0%)	31	76

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	66	TRP	2
1	A	132	PHE	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/213 (28%)	47±4 (78±6%)	13±4 (22±6%)	2	29
All	All	600/2130 (28%)	466 (78%)	134 (22%)	2	29

All 36 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	228	THR	9
1	A	233	LEU	9
1	A	133	LYS	8
1	A	209	LEU	8
1	A	74	ARG	6
1	A	58	SER	6
1	A	208	THR	6
1	A	207	LEU	5
1	A	218	TRP	5
1	A	117	SER	5
1	A	227	LYS	5
1	A	15	ASP	4
1	A	120	LEU	4
1	A	232	SER	4
1	A	193	LEU	4
1	A	230	GLU	4
1	A	118	LEU	4
1	A	16	LEU	4
1	A	121	SER	3
1	A	139	ARG	3
1	A	236	ARG	3
1	A	238	ARG	3
1	A	214	ARG	3
1	A	135	TYR	2
1	A	196	ILE	2
1	A	210	ASP	2
1	A	11	TYR	2
1	A	68	ILE	2
1	A	13	GLN	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	132	PHE	1
1	A	237	TYR	1
1	A	67	ARG	1
1	A	116	SER	1
1	A	142	TYR	1
1	A	201	PHE	1
1	A	56	ARG	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 32% for the well-defined parts and 26% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chemical_shifts_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	405
Number of shifts mapped to atoms	405
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	88	0.56 ± 0.15	Should be checked
$^{13}\text{C}_\beta$	74	-0.71 ± 0.14	Should be checked
$^{13}\text{C}'$	76	1.17 ± 0.18	Should be applied
^{15}N	84	0.31 ± 0.65	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 32%, i.e. 356 atoms were assigned a chemical shift out of a possible 1106. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	291/409 (71%)	72/169 (43%)	146/162 (90%)	73/78 (94%)
Sidechain	65/558 (12%)	0/369 (0%)	65/164 (40%)	0/25 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	0/139 (0%)	0/66 (0%)	0/69 (0%)	0/4 (0%)
Overall	356/1106 (32%)	72/604 (12%)	211/395 (53%)	73/107 (68%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 26%, i.e. 405 atoms were assigned a chemical shift out of a possible 1546. 0 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	331/561 (59%)	83/232 (36%)	164/222 (74%)	84/107 (79%)
Sidechain	74/809 (9%)	0/527 (0%)	74/237 (31%)	0/45 (0%)
Aromatic	0/176 (0%)	0/85 (0%)	0/86 (0%)	0/5 (0%)
Overall	405/1546 (26%)	83/844 (10%)	238/545 (44%)	84/157 (54%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis [i](#)

8.1 Conformationally restricting restraints [i](#)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	273
Intra-residue ($ i-j =0$)	0
Sequential ($ i-j =1$)	0
Medium range ($ i-j >1$ and $ i-j <5$)	0
Long range ($ i-j \geq 5$)	137
Inter-chain	0
Hydrogen bond restraints	136
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	1.1
Number of long range restraints per residue ¹	1.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations [i](#)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model [i](#)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.3	0.14
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Medium range (i-j >1 & i-j <5)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range (i-j ≥5)	137	50.2	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	1	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	136	49.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	136	49.8	2	1.5	0.7	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	273	100.0	2	0.7	0.7	0	0.0	0.0
Backbone-Backbone	1	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	272	99.6	2	0.7	0.7	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis


9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	1	0	1	0.11	0.11	0.0	0.11
5	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	2	0	2	0.12	0.14	0.02	0.12
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble ⓘ

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 137(IR:0, SQ:0, MR:0, LR:137, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	1	10.0
0	0	0	0	0	0	2	20.0
0	0	0	0	0	0	3	30.0

Continued on next page...


Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	4	40.0
0	0	0	0	0	0	5	50.0
0	0	0	0	0	0	6	60.0
0	0	0	0	0	0	7	70.0
0	0	0	0	0	0	8	80.0
0	0	0	0	0	0	9	90.0
0	0	0	0	0	0	10	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble ⓘ




9.4 Most violated distance restraints in the ensemble ⓘ

9.4.1 Histogram : Distribution of mean distance violations ⓘ

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,116)	1:201:A:PHE:O	1:209:A:LEU:H	2	0.12	0.02	0.12

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram

9.5.2 Table : All distance violations

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,116)	1:201:A:PHE:O	1:209:A:LEU:H	7	0.14
(1,116)	1:201:A:PHE:O	1:209:A:LEU:H	4	0.11
(1,115)	1:201:A:PHE:O	1:209:A:LEU:N	7	0.11

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found