



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:43 PM UTC

PDB ID : 1QXN / pdb_00001qxn
Title : Solution Structure of the 30 kDa Polysulfide-sulfur Transferase Homodimer from Wolinella Succinogenes
Authors : Lin, Y.J.; Dancea, F.; Loehr, F.; Klimmek, O.; Pfeiffer-Marek, S.; Nilges, M.; Wienk, H.; Kroeger, A.; Rueterjans, H.
Deposited on : 2003-09-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	137	34% 45% 7% 14%
1	B	137	32% 47% 6% 15%

2 Ensemble composition and analysis

This entry contains 10 models. Model 1 is the overall representative, medoid model (most similar to other models). The authors have identified model 8 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:65, A:70-A:89, A:95-A:129, B:3-B:65, B:70-B:89, B:95-B:128 (235)	0.76	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 3, 6, 7, 8
2	9, 10
Single-model clusters	2; 4; 5

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4338 atoms, of which 2176 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called sulfide dehydrogenase.

Mol	Chain	Residues	Atoms						Trace
1	A	137	Total	C	H	N	O	S	0
			2164	680	1088	189	200	7	
1	B	137	Total	C	H	N	O	S	0
			2164	680	1088	189	200	7	

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	130	ARG	-	expression tag	UNP Q56748
A	131	SER	-	expression tag	UNP Q56748
A	132	HIS	-	expression tag	UNP Q56748
A	133	HIS	-	expression tag	UNP Q56748
A	134	HIS	-	expression tag	UNP Q56748
A	135	HIS	-	expression tag	UNP Q56748
A	136	HIS	-	expression tag	UNP Q56748
A	137	HIS	-	expression tag	UNP Q56748
B	130	ARG	-	expression tag	UNP Q56748
B	131	SER	-	expression tag	UNP Q56748
B	132	HIS	-	expression tag	UNP Q56748
B	133	HIS	-	expression tag	UNP Q56748
B	134	HIS	-	expression tag	UNP Q56748
B	135	HIS	-	expression tag	UNP Q56748
B	136	HIS	-	expression tag	UNP Q56748
B	137	HIS	-	expression tag	UNP Q56748

- Molecule 2 is PENTASULFIDE-SULFUR (CCD ID: PS5) (formula: S₅).




Mol	Chain	Residues	Atoms	
2	A	1	Total	S
			5	5
2	B	1	Total	S
			5	5

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: sulfide dehydrogenase




- Molecule 1: sulfide dehydrogenase




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: sulfide dehydrogenase



● Molecule 1: sulfide dehydrogenase



5 Refinement protocol and experimental data overview

The models were refined using the following method: *Simulated annealing with torsion angle dynamics*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
ARIA	structure solution	1.1
ARIA	refinement	1.1

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PS5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.65±0.78	0±0/937 (0.0± 0.0%)	0.73±0.06	0±0/1263 (0.0± 0.0%)
1	B	0.65±0.78	0±0/929 (0.0± 0.0%)	0.76±0.15	0±1/1252 (0.0± 0.0%)
All	All	1.01	2/18660 (0.0%)	0.75	3/25150 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.3
1	B	0.0±0.0	0.1±0.3
All	All	0	2

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	47	ASP	CB-CG	90.48	3.78	1.52	3	1
1	A	47	ASP	CB-CG	90.29	3.77	1.52	3	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	47	ASP	CA-CB-CG	29.78	142.38	112.60	3	1
1	A	47	ASP	CB-CG-OD1	-18.50	75.84	118.40	3	1
1	B	47	ASP	CB-CG-OD1	-16.42	80.63	118.40	3	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	47	ASP	Sidechain	1
1	B	47	ASP	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	918	934	933	28±5
1	B	910	930	929	30±4
2	A	5	0	0	0±0
2	B	5	0	0	1±1
All	All	18380	18640	18620	542

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

5 of 246 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:47:ASP:CG	1:B:46:ARG:HH11	1.27	1.36	3	1
1:A:47:ASP:OD1	1:B:46:ARG:NH1	1.18	1.77	3	1
1:A:47:ASP:CG	1:B:47:ASP:HB3	1.14	1.66	3	1
1:A:47:ASP:CG	1:B:46:ARG:NH1	1.09	2.09	3	1
1:A:59:VAL:HG23	1:A:62:TYR:HB2	0.94	1.35	1	3

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/137 (86%)	110±2 (93±1%)	6±2 (5±2%)	2±1 (1±1%)	11	58
1	B	117/137 (85%)	109±2 (93±1%)	6±1 (5±1%)	2±1 (2±1%)	9	53
All	All	2350/2740 (86%)	2188 (93%)	125 (5%)	37 (2%)	10	55

5 of 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	108	LYS	8
1	A	89	CYS	7
1	B	89	CYS	6
1	A	108	LYS	4
1	A	106	GLY	3

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/114 (87%)	57±3 (58±3%)	42±3 (42±3%)	0	4
1	B	98/114 (86%)	56±3 (57±4%)	42±3 (43±4%)	0	4
All	All	1970/2280 (86%)	1131 (57%)	839 (43%)	0	4

5 of 150 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	24	MET	10
1	A	33	LEU	10
1	A	45	VAL	10
1	A	51	LEU	10
1	A	60	LYS	10

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	PS5	A	138	1	2,4,4	0.44±0.02	0±0 (0±0%)
2	PS5	B	139	1	2,4,4	0.45±0.02	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	PS5	A	138	1	1,3,3	0.07±0.05	0±0 (0±0%)
2	PS5	B	139	1	1,3,3	0.10±0.06	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PS5	B	139	1	-	0±0,2,2,2	-
2	PS5	A	138	1	-	0±0,2,2,2	-

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique torsion outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Models (Total)
2	A	138	PS5	S1-S2-S3-S4	1
2	A	138	PS5	S2-S3-S4-S5	1
2	B	139	PS5	S1-S2-S3-S4	1
2	B	139	PS5	S2-S3-S4-S5	1

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided