



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:22 AM UTC

PDB ID : 4QXQ / pdb_00004qxq
Title : Crystal structure of hSTING(S162A/Q266I) in complex with DMXAA
Authors : Gao, P.; Patel, D.J.
Deposited on : 2014-07-21
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	188	73% 22% . .
1	B	188	72% 19% 6% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3031 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Stimulator of interferon genes protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1460	919	259	276	6			
1	B	182	Total	C	N	O	S	0	0	0
			1460	919	259	276	6			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	154	SER	-	expression tag	UNP Q86WV6
A	162	ALA	SER	engineered mutation	UNP Q86WV6
A	232	ARG	HIS	engineered mutation	UNP Q86WV6
A	266	ILE	GLN	engineered mutation	UNP Q86WV6
B	154	SER	-	expression tag	UNP Q86WV6
B	162	ALA	SER	engineered mutation	UNP Q86WV6
B	232	ARG	HIS	engineered mutation	UNP Q86WV6
B	266	ILE	GLN	engineered mutation	UNP Q86WV6

- Molecule 2 is (5,6-dimethyl-9-oxo-9H-xanthen-4-yl)acetic acid (CCD ID: 1YE) (formula: C₁₇H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	17	4		
2	B	1	Total	C	O	0	0
			21	17	4		

- Molecule 3 is SULFATE ION (CCD ID: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	26	Total 26	O 26	0	0
4	B	38	Total 38	O 38	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Stimulator of interferon genes protein



- Molecule 1: Stimulator of interferon genes protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, α , β , γ	147.95Å 147.95Å 36.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.42 50.00 – 2.42	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.42) 99.9 (50.00-2.42)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.42Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.183 , 0.229 0.195 , 0.229	Depositor DCC
R_{free} test set	900 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	52.0	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.066 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3031	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 1YE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	1/1488 (0.1%)	0.95	0/2018
1	B	0.94	2/1488 (0.1%)	1.01	2/2018 (0.1%)
All	All	0.87	3/2976 (0.1%)	0.98	2/4036 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	238	ARG	C-O	-5.13	1.17	1.23
1	A	167	TYR	C-O	-5.10	1.17	1.24
1	B	238	ARG	CA-C	-5.01	1.46	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	238	ARG	N-CA-C	-8.10	97.44	110.32
1	B	237	ASP	N-CA-C	5.16	118.84	112.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1460	0	1439	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1460	0	1439	21	0
2	A	21	0	13	2	0
2	B	21	0	13	1	0
3	A	5	0	0	0	0
4	A	26	0	0	1	0
4	B	38	0	0	0	0
All	All	3031	0	2904	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:230:GLY:HA3	1:B:235:ILE:HD12	1.71	0.71
1:A:334:ARG:O	1:A:335:GLN:NE2	2.22	0.68
1:A:320:ASP:OD1	1:A:322:SER:N	2.31	0.63
1:B:165:ILE:HA	1:B:169:ARG:HD2	1.83	0.60
1:B:194:VAL:HB	1:B:256:THR:HG22	1.84	0.59
1:A:213:SER:HB3	4:A:513:HOH:O	2.02	0.59
1:A:319:ASP:OD1	1:A:320:ASP:N	2.40	0.55
1:A:179:ILE:HD13	1:A:225:LEU:HD21	1.87	0.55
2:A:401:1YE:H12	2:B:401:1YE:H12	1.89	0.54
1:B:329:VAL:HG12	1:B:333:LEU:HD12	1.91	0.53
1:B:334:ARG:O	1:B:335:GLN:NE2	2.26	0.52
1:B:169:ARG:HB2	1:B:304:GLU:OE2	2.10	0.52
1:B:186:TYR:CE2	1:B:189:LEU:HD12	2.46	0.51
2:A:401:1YE:OAC	1:B:238:ARG:NH1	2.43	0.51
1:A:179:ILE:HD11	1:A:258:VAL:HG21	1.92	0.50
1:A:209:PRO:HG3	1:A:215:ALA:HB2	1.94	0.50
1:A:183:ASN:O	1:A:187:ASN:HB2	2.11	0.50
1:A:197:ARG:NH1	1:A:310:ARG:HD3	2.26	0.49
1:A:170:LEU:HD13	1:A:235:ILE:HD13	1.95	0.48
1:A:324:SER:HB3	1:A:327:GLN:HB2	1.96	0.48
1:B:190:LEU:HD11	1:B:194:VAL:HG12	1.96	0.48
1:B:265:LEU:HD23	1:B:268:LEU:HD12	1.95	0.48
1:A:328:GLU:CD	1:A:331:ARG:HH21	2.21	0.48
1:B:180:ARG:HG3	1:B:180:ARG:HH21	1.78	0.47
1:B:236:LYS:HE2	1:B:237:ASP:OD2	2.15	0.46
1:B:227:GLN:HG2	1:B:240:TYR:O	2.16	0.46
1:A:334:ARG:C	1:A:335:GLN:HE21	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:186:TYR:CD2	1:B:189:LEU:HD12	2.52	0.45
1:A:222:LEU:HD11	1:A:244:ILE:HG22	2.00	0.44
1:A:299:LEU:HD22	1:A:305:SER:CB	2.48	0.44
1:B:279:PHE:HZ	1:B:287:GLN:HG3	1.83	0.44
1:A:210:ASP:HA	1:B:233:ALA:HB2	2.00	0.44
1:A:333:LEU:HD23	1:A:333:LEU:HA	1.75	0.43
1:B:289:LYS:O	1:B:293:ARG:HG3	2.18	0.42
1:B:202:LEU:HD11	1:B:311:LEU:HD22	2.00	0.42
1:A:199:TYR:HD2	1:A:312:ILE:HD11	1.86	0.41
1:B:230:GLY:CA	1:B:235:ILE:HD12	2.45	0.41
1:B:159:LEU:HD23	1:B:159:LEU:HA	1.92	0.40
1:B:196:GLN:HE21	1:B:196:GLN:N	2.19	0.40
1:A:179:ILE:HG23	1:A:194:VAL:CG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/188 (96%)	176 (98%)	4 (2%)	0	100	100
1	B	180/188 (96%)	174 (97%)	6 (3%)	0	100	100
All	All	360/376 (96%)	350 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/163 (96%)	140 (89%)	17 (11%)	6	9
1	B	157/163 (96%)	133 (85%)	24 (15%)	3	3
All	All	314/326 (96%)	273 (87%)	41 (13%)	4	5

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	155	VAL
1	A	159	LEU
1	A	165	ILE
1	A	176	GLN
1	A	180	ARG
1	A	184	GLN
1	A	194	VAL
1	A	196	GLN
1	A	198	LEU
1	A	213	SER
1	A	220	ARG
1	A	238	ARG
1	A	296	GLU
1	A	306	GLN
1	A	316	GLU
1	A	326	SER
1	A	335	GLN
1	B	169	ARG
1	B	171	ILE
1	B	176	GLN
1	B	180	ARG
1	B	184	GLN
1	B	189	LEU
1	B	190	LEU
1	B	194	VAL
1	B	196	GLN
1	B	197	ARG
1	B	210	ASP
1	B	213	SER
1	B	225	LEU
1	B	231	ASP
1	B	232	ARG
1	B	241	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	258	VAL
1	B	276	GLN
1	B	281	ARG
1	B	304	GLU
1	B	322	SER
1	B	326	SER
1	B	333	LEU
1	B	335	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	ASN
1	B	196	GLN
1	B	228	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1YE	A	401	-	23,23,23	2.46	8 (34%)	33,34,34	1.21	3 (9%)
2	1YE	B	401	-	23,23,23	2.75	8 (34%)	33,34,34	1.04	3 (9%)
3	SO4	A	402	-	4,4,4	0.27	0	6,6,6	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1YE	A	401	-	-	2/4/4/4	0/3/3/3
2	1YE	B	401	-	-	0/4/4/4	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	1YE	CAK-CAP	-6.77	1.42	1.51
2	A	401	1YE	CAK-CAP	-6.04	1.43	1.51
2	B	401	1YE	CAS-CAQ	-5.31	1.37	1.48
2	B	401	1YE	CAR-CAQ	-5.15	1.38	1.48
2	B	401	1YE	CAB-CAO	-4.84	1.42	1.51
2	A	401	1YE	CAR-CAQ	-4.76	1.38	1.48
2	A	401	1YE	CAB-CAO	-4.59	1.42	1.51
2	A	401	1YE	CAS-CAQ	-4.42	1.39	1.48
2	B	401	1YE	CAA-CAN	-3.68	1.44	1.51
2	A	401	1YE	OAL-CAT	-3.33	1.34	1.39
2	B	401	1YE	OAL-CAT	-3.20	1.34	1.39
2	B	401	1YE	CAK-CAM	3.09	1.57	1.51
2	A	401	1YE	CAA-CAN	-2.99	1.45	1.51
2	B	401	1YE	OAL-CAU	-2.36	1.35	1.39
2	A	401	1YE	OAL-CAU	-2.26	1.35	1.39
2	A	401	1YE	CAK-CAM	2.08	1.55	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	1YE	OAC-CAM-CAK	-2.97	113.99	122.94
2	B	401	1YE	CAB-CAO-CAN	2.38	123.78	119.67
2	B	401	1YE	CAP-CAK-CAM	2.22	117.90	113.10
2	B	401	1YE	CAB-CAO-CAT	-2.17	117.85	121.24
2	A	401	1YE	CAP-CAK-CAM	2.09	117.61	113.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	1YE	CAR-CAU-CAP	2.07	122.76	119.46

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	1YE	CAP-CAK-CAM-OAE
2	A	401	1YE	CAP-CAK-CAM-OAC


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	1YE	2	0
2	B	401	1YE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/188 (96%)	-0.14	2 (1%) 78 75	24, 51, 89, 173	0
1	B	182/188 (96%)	-0.10	2 (1%) 78 75	25, 50, 103, 142	0
All	All	364/376 (96%)	-0.12	4 (1%) 78 75	24, 51, 96, 173	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	ALA	2.9
1	B	228	GLN	2.5
1	B	233	ALA	2.4
1	A	215	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	402	5/5	0.64	0.26	208,209,210,210	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	1YE	A	401	21/21	0.97	0.05	23,28,39,41	0
2	1YE	B	401	21/21	0.98	0.05	18,26,33,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 1YE A 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 1YE B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.