



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:16 PM UTC

PDB ID : 1QYX / pdb_00001qyx
Title : Crystal structure of human estrogenic 17beta-hydroxysteroid dehydrogenase complex with androstenedione and NADP
Authors : Shi, R.; Lin, S.X.
Deposited on : 2003-09-12
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	327	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	600	-	X	-	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 2300 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Estradiol 17 beta-dehydrogenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	2	0
			2131	1353	378	389	11			

- Molecule 2 is 4-ANDROSTENE-3-17-DIONE (CCD ID: ASD) (formula: $C_{19}H_{26}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	19	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	115	Total 115	O 115	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Estradiol 17 beta-dehydrogenase 1



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.70Å 44.03Å 60.79Å 90.00° 99.52° 90.00°	Depositor
Resolution (Å)	10.00 – 1.89	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.89)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.186 , 0.227	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2300	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, ASD, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/2180	1.57	14/2956 (0.5%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ARG	CD-NE-CZ	8.53	136.34	124.40
1	A	252	ARG	CG-CD-NE	-7.89	94.64	112.00
1	A	77	GLU	CB-CG-CD	6.70	123.99	112.60
1	A	252	ARG	NH1-CZ-NH2	6.67	127.97	119.30
1	A	252	ARG	NE-CZ-NH2	-6.57	113.29	119.20
1	A	258	ARG	N-CA-C	5.93	117.75	111.28
1	A	26	PRO	O-C-N	-5.83	115.11	122.17
1	A	68	ASP	CA-CB-CG	5.56	118.16	112.60
1	A	189	HIS	CB-CA-C	-5.33	102.57	110.67
1	A	237	ALA	O-C-N	-5.25	116.56	122.12
1	A	234	GLU	CB-CG-CD	5.22	121.47	112.60
1	A	109	SER	CA-CB-OG	-5.08	100.93	111.10
1	A	76	ARG	NE-CZ-NH1	5.05	126.55	121.50
1	A	109	SER	CB-CA-C	-5.00	102.48	110.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2131	0	2169	17	0
2	A	21	0	26	2	0
3	A	27	0	11	0	0
4	A	6	0	4	0	0
5	A	115	0	0	0	0
All	All	2300	0	2210	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:LEU:HD11	1:A:244:LEU:HG	1.86	0.56
1:A:263:LEU:O	1:A:267:LEU:HG	2.11	0.50
1:A:13:GLY:O	1:A:17:HIS:HD2	1.95	0.49
1:A:179:HIS:HB3	1:A:249:PRO:HG2	1.96	0.48
1:A:86:VAL:CG1	1:A:244:LEU:HD21	2.44	0.47
1:A:95:LEU:HB2	1:A:110:VAL:HG21	1.97	0.47
1:A:16:LEU:HD23	1:A:16:LEU:C	2.41	0.46
1:A:25:ASP:OD2	1:A:27:SER:OG	2.26	0.45
1:A:174:LEU:HB3	1:A:175:PRO:HD3	1.99	0.44
1:A:244:LEU:HD23	1:A:244:LEU:HA	1.74	0.44
1:A:42:GLN:NE2	1:A:46:TRP:HE1	2.15	0.43
1:A:55:PRO:HA	1:A:56:PRO:HD3	1.92	0.43
1:A:143:VAL:HG11	2:A:500:ASD:H72	2.00	0.43
1:A:25:ASP:HA	1:A:26:PRO:HD3	1.91	0.42
1:A:64:LEU:HD23	1:A:64:LEU:C	2.44	0.42
1:A:86:VAL:HG11	1:A:244:LEU:HD21	2.02	0.42
1:A:11:SER:HB3	1:A:36:LEU:HD23	2.03	0.41
2:A:500:ASD:H193	2:A:500:ASD:H112	1.90	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/327 (84%)	273 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/257 (90%)	227 (98%)	4 (2%)	53	52

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	SER
1	A	77	GLU
1	A	111	LEU
1	A	131	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	HIS
1	A	42	GLN
1	A	232	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAP	A	361	-	29,29,52	1.54	6 (20%)	44,45,80	2.33	11 (25%)
4	GOL	A	600	-	5,5,5	4.77	5 (100%)	5,5,5	4.39	3 (60%)
2	ASD	A	500	-	24,24,24	1.52	5 (20%)	39,39,39	3.24	19 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	361	-	-	3/15/31/67	0/3/3/5
4	GOL	A	600	-	-	2/4/4/4	-
2	ASD	A	500	-	-	-	0/4/4/4

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	600	GOL	C3-C2	-7.53	1.23	1.51
4	A	600	GOL	O1-C1	4.60	1.61	1.42
4	A	600	GOL	O3-C3	4.03	1.59	1.42
3	A	361	NAP	C8A-N7A	3.74	1.38	1.31
4	A	600	GOL	O2-C2	-3.23	1.34	1.43
2	A	500	ASD	C16-C17	3.13	1.56	1.51
4	A	600	GOL	C1-C2	-3.03	1.40	1.51
3	A	361	NAP	C4A-N3A	3.02	1.40	1.34
2	A	500	ASD	C12-C13	2.98	1.59	1.54
2	A	500	ASD	C7-C8	2.86	1.58	1.53
3	A	361	NAP	P2B-O2B	2.73	1.64	1.59
3	A	361	NAP	C6A-N1A	2.66	1.43	1.35
2	A	500	ASD	C4-C5	2.50	1.38	1.34
3	A	361	NAP	C4A-N9A	-2.24	1.33	1.37
2	A	500	ASD	C13-C14	-2.22	1.50	1.54
3	A	361	NAP	O5B-C5B	-2.07	1.36	1.44

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	ASD	C18-C13-C14	7.49	123.74	112.99
2	A	500	ASD	O2-C17-C13	7.44	135.73	125.98
3	A	361	NAP	C5A-C4A-N3A	-7.11	116.92	126.72
4	A	600	GOL	O2-C2-C3	6.87	137.63	109.18
3	A	361	NAP	O4B-C1B-N9A	6.83	121.21	108.09
2	A	500	ASD	O2-C17-C16	-6.42	117.93	125.75
4	A	600	GOL	O3-C3-C2	6.02	137.49	110.38
2	A	500	ASD	C11-C9-C10	-5.71	106.05	113.08
2	A	500	ASD	C18-C13-C12	-5.71	104.58	111.10
2	A	500	ASD	C1-C10-C9	-5.08	102.01	108.74
2	A	500	ASD	C11-C9-C8	4.90	118.61	111.78
2	A	500	ASD	C16-C15-C14	-4.62	95.99	103.00
3	A	361	NAP	N3A-C4A-N9A	4.17	134.27	127.17
3	A	361	NAP	O4B-C1B-C2B	-4.16	99.43	106.59
2	A	500	ASD	C13-C14-C8	4.00	117.34	113.13
3	A	361	NAP	O2A-PA-O5B	3.98	117.05	106.67
3	A	361	NAP	C2A-N3A-C4A	3.75	120.98	111.83
2	A	500	ASD	C12-C13-C17	3.74	122.01	116.71
2	A	500	ASD	C15-C14-C8	-3.57	113.41	119.10
4	A	600	GOL	O1-C1-C2	3.45	125.93	110.38
2	A	500	ASD	C2-C3-C4	3.42	121.93	116.73
3	A	361	NAP	O3-PA-O1A	-3.22	98.28	110.83
2	A	500	ASD	C7-C8-C9	-3.09	106.82	110.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	361	NAP	C6A-C5A-C4A	3.04	121.33	117.18
3	A	361	NAP	N3A-C2A-N1A	-2.94	124.13	128.58
2	A	500	ASD	O1-C3-C2	-2.84	115.94	121.59
3	A	361	NAP	C5A-C4A-N9A	2.75	108.81	105.81
2	A	500	ASD	C6-C7-C8	-2.48	107.29	111.64
2	A	500	ASD	C11-C12-C13	2.26	117.59	112.78
2	A	500	ASD	C19-C10-C1	2.25	112.86	109.43
2	A	500	ASD	C16-C17-C13	-2.23	106.34	108.56
2	A	500	ASD	C5-C4-C3	-2.17	120.32	123.67
3	A	361	NAP	C1B-N9A-C8A	2.06	131.67	127.09

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	361	NAP	C5B-O5B-PA-O1A
3	A	361	NAP	C5B-O5B-PA-O2A
3	A	361	NAP	C5B-O5B-PA-O3
4	A	600	GOL	O1-C1-C2-C3
4	A	600	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	ASD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.