



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 03:17 PM UTC

PDB ID : 7QY8 / pdb_00007qy8
Title : The structure of T. forsythia NanH
Authors : Rafferty, J.; Stafford, G.; Satur, M.
Deposited on : 2022-01-27
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	519	4% 90% 9%
1	B	519	4% 89% 10%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 16489 atoms, of which 8053 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BNR/Asp-box repeat protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	519	Total	C	H	N	O	S	128	0	0
			8008	2499	3994	723	772	20			
1	B	519	Total	C	H	N	O	S	131	1	0
			7994	2497	3983	720	774	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	237	ALA	ASP	conflict	UNP G8UIQ1
B	237	ALA	ASP	conflict	UNP G8UIQ1

- Molecule 2 is 3-sialyllactose (CCD ID: I2K) (formula: $C_{23}H_{39}NO_{19}$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O		11	0
			81	23	38	1	19			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	H	N	O	11	0
			81	23	38	1	19		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	171	Total	O	0	0
			171	171		
3	B	154	Total	O	0	0
			154	154		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BNR/Asp-box repeat protein



- Molecule 1: BNR/Asp-box repeat protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	80.06Å 80.06Å 349.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.90 – 2.06 72.90 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.90-2.06) 99.4 (72.90-2.06)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.245 , 0.279 0.242 , 0.276	Depositor DCC
R_{free} test set	3636 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	37.8	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 24.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16489	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: I2K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	0/4092	1.26	0/5559
1	B	1.01	0/4089	1.25	0/5557
All	All	1.01	0/8181	1.25	0/11116

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	475	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4014	3994	3975	34	0
1	B	4011	3983	3959	35	0
2	A	43	38	0	0	0
2	B	43	38	0	0	0
3	A	171	0	0	1	0
3	B	154	0	0	1	0
All	All	8436	8053	7934	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:MET:HG3	1:A:196:MET:HE1	1.50	0.94
1:B:263:MET:HE1	1:B:324:MET:CB	2.04	0.88
1:A:263:MET:HE1	1:A:324:MET:CB	2.05	0.86
1:B:263:MET:HE1	1:B:324:MET:HB2	1.65	0.79
1:A:263:MET:HE1	1:A:324:MET:HB2	1.65	0.77
1:B:84:VAL:HB	1:B:135:VAL:HG11	1.68	0.74
1:B:263:MET:HE1	1:B:324:MET:HB3	1.68	0.73
1:A:263:MET:HE1	1:A:324:MET:HB3	1.69	0.72
1:B:166:LEU:H	1:B:254:GLN:HE22	1.41	0.66
1:A:166:LEU:H	1:A:254:GLN:HE22	1.44	0.66
1:A:432:VAL:H	1:A:448:ASN:HD22	1.44	0.65
1:B:432:VAL:H	1:B:448:ASN:HD22	1.45	0.65
1:A:160:MET:HG3	1:A:196:MET:CE	2.28	0.62
1:A:39:GLN:NE2	1:A:56:ARG:HH11	1.98	0.62
1:A:488:HIS:HD2	1:A:489:HIS:ND1	1.98	0.61
1:B:488:HIS:HD2	1:B:489:HIS:ND1	1.98	0.60
1:A:39:GLN:HG2	1:A:56:ARG:HB2	1.87	0.57
1:A:173:THR:C	3:A:701:HOH:O	2.48	0.56
1:B:213:ILE:HG21	1:B:280:ASP:HA	1.87	0.56
1:A:213:ILE:HG21	1:A:280:ASP:HA	1.87	0.55
1:B:494:ALA:N	3:B:701:HOH:O	2.30	0.54
1:A:148:VAL:H	1:A:152:ASN:HD21	1.55	0.53
1:B:148:VAL:H	1:B:152:ASN:HD21	1.56	0.53
1:B:284:LEU:HD12	1:B:284:LEU:C	2.33	0.53
1:A:506:HIS:HA	1:A:550:LEU:O	2.09	0.52
1:A:284:LEU:HD12	1:A:284:LEU:C	2.34	0.52
1:B:506:HIS:HA	1:B:550:LEU:O	2.10	0.52
1:B:71:LEU:C	1:B:71:LEU:HD12	2.35	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:363:MET:HE1	1:B:438:LEU:CD2	2.42	0.50
1:B:263:MET:HE2	1:B:265:PHE:CE2	2.47	0.49
1:A:54:LEU:HD22	1:A:158:VAL:CG1	2.43	0.49
1:B:363:MET:HE1	1:B:438:LEU:HD21	1.94	0.49
1:A:422:MET:HB2	1:A:431:ALA:HB3	1.94	0.49
1:B:263:MET:HE3	1:B:338:SER:CB	2.43	0.48
1:A:263:MET:HE3	1:A:338:SER:CB	2.44	0.48
1:A:263:MET:HE2	1:A:265:PHE:CE2	2.49	0.47
1:A:72:THR:OG1	1:A:176:GLU:HB3	2.16	0.46
1:B:422:MET:HB2	1:B:431:ALA:HB3	1.96	0.46
1:B:160:MET:O	1:B:162:ARG:NH1	2.50	0.45
1:A:341:ILE:HG22	1:A:341:ILE:O	2.17	0.45
1:A:203:ALA:HB2	1:A:539:HIS:CE1	2.53	0.44
1:B:286:ASP:OD1	1:B:391:ARG:NH1	2.50	0.44
1:B:248:ARG:NH1	1:B:331:ASP:O	2.51	0.44
1:A:98:THR:HG22	1:B:310:ASN:HD21	1.82	0.43
1:A:116:ARG:NH2	1:B:207:GLY:HA2	2.33	0.43
1:B:345:VAL:HB	1:B:397:ILE:HD12	2.01	0.43
1:B:203:ALA:HB2	1:B:539:HIS:CE1	2.54	0.43
1:B:363:MET:HE2	1:B:418:LEU:HD11	2.00	0.43
1:A:263:MET:CE	1:A:324:MET:HB2	2.43	0.42
1:B:263:MET:HE2	1:B:265:PHE:CZ	2.54	0.42
1:A:283:ILE:O	1:A:359:ARG:HA	2.20	0.42
1:A:263:MET:HE3	1:A:338:SER:HB3	2.01	0.42
1:A:234:ASN:OD1	1:A:236:VAL:HG22	2.19	0.42
1:A:263:MET:CE	1:A:324:MET:HB3	2.44	0.42
1:A:84:VAL:HG12	1:A:135:VAL:HG21	2.03	0.41
1:A:345:VAL:HB	1:A:397:ILE:HD12	2.02	0.41
1:B:263:MET:HE3	1:B:338:SER:HB2	2.02	0.41
1:B:263:MET:HE3	1:B:338:SER:HB3	2.01	0.41
1:B:234:ASN:OD1	1:B:236:VAL:HG22	2.20	0.41
1:A:278:VAL:HA	1:A:297:ALA:O	2.21	0.41
1:B:283:ILE:O	1:B:359:ARG:HA	2.20	0.41
1:B:325:MET:CE	1:B:341:ILE:HD13	2.51	0.41
1:A:352:PHE:O	1:A:374:PHE:HA	2.21	0.40
1:B:488:HIS:CD2	1:B:489:HIS:ND1	2.85	0.40
1:B:278:VAL:HA	1:B:297:ALA:O	2.21	0.40
1:A:54:LEU:HD22	1:A:158:VAL:HG13	2.02	0.40
1:B:74:ARG:HA	1:B:137:ASN:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/519 (100%)	494 (96%)	22 (4%)	1 (0%)	43	38
1	B	518/519 (100%)	493 (95%)	24 (5%)	1 (0%)	43	38
All	All	1035/1038 (100%)	987 (95%)	46 (4%)	2 (0%)	43	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	456	ILE
1	B	456	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/445 (100%)	436 (98%)	7 (2%)	55	55
1	B	442/445 (99%)	436 (99%)	6 (1%)	59	60
All	All	885/890 (99%)	872 (98%)	13 (2%)	57	58

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	98	THR
1	A	143	THR
1	A	148	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	255	THR
1	A	551	ILE
1	A	552	ARG
1	B	68	LEU
1	B	71	LEU
1	B	134	THR
1	B	143	THR
1	B	485	GLU
1	B	551	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	52	ASN
1	A	137	ASN
1	A	152	ASN
1	A	180	ASN
1	A	202	HIS
1	A	241	HIS
1	A	254	GLN
1	A	399	GLN
1	A	425	ASN
1	A	448	ASN
1	A	488	HIS
1	B	52	ASN
1	B	152	ASN
1	B	180	ASN
1	B	202	HIS
1	B	241	HIS
1	B	254	GLN
1	B	310	ASN
1	B	399	GLN
1	B	448	ASN
1	B	488	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	I2K	A	601	-	42,44,44	0.72	2 (4%)	58,64,64	1.18	5 (8%)
2	I2K	B	601	-	42,44,44	0.64	1 (2%)	58,64,64	0.95	4 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	I2K	A	601	-	-	18/45/85/85	0/2/2/2
2	I2K	B	601	-	-	19/45/85/85	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	I2K	O18-C21	-2.60	1.21	1.30
2	A	601	I2K	O18-C21	-2.49	1.21	1.30
2	A	601	I2K	O6-C9	2.40	1.45	1.42

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	I2K	C6-C5-C4	4.36	121.25	113.05
2	A	601	I2K	C22-C23-C3	3.56	115.22	109.72
2	A	601	I2K	O5-C4-C5	2.92	111.21	106.65
2	B	601	I2K	C6-C5-C4	2.82	118.36	113.05
2	A	601	I2K	C8-C22-C23	2.64	116.08	110.73
2	B	601	I2K	C4-C3-N1	2.63	115.11	110.91
2	B	601	I2K	C22-C23-C3	2.46	113.53	109.72
2	A	601	I2K	C8-O5-C4	-2.07	109.50	114.36
2	B	601	I2K	O6-C8-C21	-2.05	104.65	109.86

There are no chirality outliers.

All (37) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	I2K	O1-C2-N1-C3
2	A	601	I2K	C3-C4-C5-C6
2	A	601	I2K	C3-C4-C5-O2
2	A	601	I2K	O5-C4-C5-C6
2	A	601	I2K	O5-C4-C5-O2
2	A	601	I2K	O18-C21-C8-O6
2	A	601	I2K	C15-C14-C17-C18
2	A	601	I2K	C14-C17-C18-C19
2	A	601	I2K	O13-C17-C18-O14
2	A	601	I2K	O13-C17-C18-C19
2	B	601	I2K	O1-C2-N1-C3
2	B	601	I2K	C4-C3-N1-C2
2	B	601	I2K	C3-C4-C5-C6
2	B	601	I2K	C3-C4-C5-O2
2	B	601	I2K	O5-C4-C5-C6
2	B	601	I2K	O5-C4-C5-O2
2	B	601	I2K	O3-C6-C7-O4
2	B	601	I2K	O18-C21-C8-O6
2	B	601	I2K	C14-C15-C16-O12
2	B	601	I2K	O11-C15-C16-O12
2	B	601	I2K	C14-C17-C18-C19
2	B	601	I2K	O13-C17-C18-O14
2	B	601	I2K	O13-C17-C18-C19
2	A	601	I2K	C1-C2-N1-C3
2	B	601	I2K	C1-C2-N1-C3
2	B	601	I2K	C5-C6-C7-O4
2	B	601	I2K	O9-C13-O10-C14
2	A	601	I2K	O9-C11-C12-O8
2	A	601	I2K	C5-C6-C7-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	601	I2K	O3-C6-C7-O4
2	B	601	I2K	O17-C21-C8-O6
2	A	601	I2K	C14-C17-C18-O14
2	B	601	I2K	C14-C17-C18-O14
2	A	601	I2K	O17-C21-C8-O6
2	B	601	I2K	C23-C3-N1-C2
2	A	601	I2K	C10-C11-C12-O8
2	A	601	I2K	C4-C3-N1-C2


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand I2K A 601





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	519/519 (100%)	0.50	19 (3%)	45	46	31, 47, 70, 101	0
1	B	519/519 (100%)	0.76	21 (4%)	42	43	30, 53, 69, 80	1 (0%)
All	All	1038/1038 (100%)	0.63	40 (3%)	43	44	30, 50, 70, 101	1 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	LYS	4.1
1	B	34	ASP	3.4
1	A	135	VAL	3.1
1	B	207	GLY	3.0
1	B	292	VAL	2.8
1	B	547	ILE	2.8
1	B	262	ALA	2.8
1	A	36	VAL	2.8
1	B	261	ILE	2.7
1	A	132	VAL	2.6
1	A	413	VAL	2.6
1	A	71	LEU	2.5
1	B	266	GLY	2.5
1	A	183	PRO	2.4
1	B	360	GLY	2.4
1	A	134	THR	2.4
1	B	199	GLY	2.4
1	A	552	ARG	2.4
1	B	137	ASN	2.3
1	B	274	GLY	2.3
1	A	35	SER	2.3
1	A	148	VAL	2.3
1	B	293	TRP	2.3
1	A	137	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	335	TRP	2.3
1	B	87	VAL	2.2
1	B	413	VAL	2.2
1	B	471	ILE	2.2
1	A	184	ALA	2.2
1	A	496	LEU	2.2
1	A	125	TYR	2.2
1	A	178	VAL	2.2
1	A	92	ALA	2.2
1	B	552	ARG	2.2
1	B	285	VAL	2.1
1	B	209	ALA	2.1
1	B	358	GLY	2.1
1	B	47	VAL	2.1
1	A	37	TYR	2.0
1	A	187	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	I2K	B	601	43/43	0.69	0.15	53,84,111,115	11
2	I2K	A	601	43/43	0.75	0.15	44,80,106,112	11

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around I2K B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around I2K A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.