



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 05:28 AM UTC

PDB ID : 8QZ3 / pdb_00008qz3
Title : Crystal structure of human two pore domain potassium ion channel TREK-2 (K2P10.1) in complex with an activatory nanobody (Nb67)
Authors : Baronina, A.; Pike, A.C.W.; Rodstrom, K.E.J.; Ang, J.; Bushell, S.R.; Chalk, R.; Mukhopadhyay, S.M.M.; Pardon, E.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Burgess-Brown, N.A.; Tucker, S.J.; Steyaert, J.; Carpenter, E.P.; Structural Genomics Consortium (SGC)
Deposited on : 2023-10-26
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	23% 91% 5%
1	B	274	21% 93% .
2	C	137	4% 81% 12% 7%
2	D	137	11% 80% 12% 7%
2	E	137	15% 82% 10% 8%

Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7082 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium channel subfamily K member 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			1960	1308	303	345	4			
1	B	266	Total	C	N	O	S	0	0	0
			2033	1354	317	358	4			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	74	MET	-	initiating methionine	UNP P57789
A	149	GLN	ASN	engineered mutation	UNP P57789
A	152	GLN	ASN	engineered mutation	UNP P57789
A	153	GLN	ASN	engineered mutation	UNP P57789
A	341	ALA	-	expression tag	UNP P57789
A	342	GLU	-	expression tag	UNP P57789
A	343	ASN	-	expression tag	UNP P57789
A	344	LEU	-	expression tag	UNP P57789
A	345	TYR	-	expression tag	UNP P57789
A	346	PHE	-	expression tag	UNP P57789
A	347	GLN	-	expression tag	UNP P57789
B	74	MET	-	initiating methionine	UNP P57789
B	149	GLN	ASN	engineered mutation	UNP P57789
B	152	GLN	ASN	engineered mutation	UNP P57789
B	153	GLN	ASN	engineered mutation	UNP P57789
B	341	ALA	-	expression tag	UNP P57789
B	342	GLU	-	expression tag	UNP P57789
B	343	ASN	-	expression tag	UNP P57789
B	344	LEU	-	expression tag	UNP P57789
B	345	TYR	-	expression tag	UNP P57789
B	346	PHE	-	expression tag	UNP P57789
B	347	GLN	-	expression tag	UNP P57789


- Molecule 2 is a protein called Nanobody 67.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	127	Total	C	N	O	S	0	2	0
			977	610	172	191	4			
2	D	127	Total	C	N	O	S	0	0	0
			959	599	168	188	4			
2	E	126	Total	C	N	O	S	0	0	0
			949	594	165	186	4			

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	K	0	0
			3	3		
3	B	2	Total	K	0	0
			2	2		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	53	Total	O	0	0
			53	53		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	43	Total 43	O 43	0	0
5	C	48	Total 48	O 48	0	0
5	D	27	Total 27	O 27	0	0
5	E	20	Total 20	O 20	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Potassium channel subfamily K member 10




- Molecule 1: Potassium channel subfamily K member 10




- Molecule 2: Nanobody 67



- Molecule 2: Nanobody 67



- Molecule 2: Nanobody 67



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	51.26Å 102.10Å 215.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.81 – 2.40 58.81 – 2.40	Depositor EDS
% Data completeness (in resolution range)	75.6 (58.81-2.40) 75.6 (58.81-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.40Å)	Xtriage
Refinement program	BUSTER 2.10.4	Depositor
R, R_{free}	0.278 , 0.291 (Not available) , 0.272	Depositor DCC
R_{free} test set	1747 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	7082	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/2009	0.97	0/2743
1	B	0.55	0/2083	0.94	0/2837
2	C	0.50	0/1003	0.80	1/1357 (0.1%)
2	D	0.49	0/979	0.81	2/1326 (0.2%)
2	E	0.51	0/969	0.80	1/1313 (0.1%)
All	All	0.53	0/7043	0.90	4/9576 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	8	GLY	N-CA-C	-7.07	103.77	112.68
2	D	110	PHE	CA-CB-CG	6.27	120.07	113.80
2	C	8	GLY	N-CA-C	6.08	124.70	115.63
2	D	8	GLY	N-CA-C	5.69	123.95	115.64

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1960	0	1954	9	0
1	B	2033	0	2051	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	977	0	932	8	0
2	D	959	0	907	10	0
2	E	949	0	894	7	0
3	A	3	0	0	0	0
3	B	2	0	0	0	0
4	D	8	0	14	1	0
5	A	53	0	0	0	0
5	B	43	0	0	0	0
5	C	48	0	0	0	0
5	D	27	0	0	0	0
5	E	20	0	0	0	0
All	All	7082	0	6752	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:329:VAL:O	1:A:332:LYS:HG2	1.96	0.66
2:C:83:MET:HE1	2:C:123:VAL:HG11	1.78	0.65
2:D:83:MET:HE1	2:D:123:VAL:HG11	1.82	0.62
1:A:245:ILE:HG12	1:A:319:VAL:HG22	1.82	0.61
2:E:83:MET:HE1	2:E:123:VAL:HG11	1.82	0.61
2:D:51:LEU:HD13	2:D:58:THR:HG22	1.85	0.58
2:D:33:VAL:HB	2:D:99:ARG:HB3	1.87	0.57
1:A:190:ILE:HG23	1:B:276:VAL:HG21	1.87	0.56
2:D:83:MET:HE2	2:D:86:LEU:HD21	1.89	0.55
2:C:33:VAL:HB	2:C:99:ARG:HB3	1.89	0.55
2:E:33:VAL:HB	2:E:99:ARG:HB3	1.90	0.53
2:D:39:GLU:HB2	2:D:45:ARG:HG2	1.91	0.53
2:C:83:MET:HE2	2:C:86:LEU:HD21	1.89	0.53
2:E:83:MET:HE2	2:E:86:LEU:HD21	1.91	0.52
2:E:51:LEU:HD13	2:E:58:THR:HG22	1.90	0.52
1:A:276:VAL:HG21	1:B:190:ILE:HG23	1.92	0.51
1:A:144:VAL:HG13	1:B:141:ASN:HB3	1.93	0.51
2:D:45:ARG:NH1	2:D:111:TRP:CZ2	2.83	0.46
2:C:91:THR:HG23	2:C:124:THR:HA	1.97	0.46
2:D:91:THR:HG23	2:D:124:THR:HA	1.96	0.46
1:A:137:LEU:HD23	1:B:137:LEU:HD23	1.98	0.45
1:A:149:GLN:HG2	1:A:292:ASN:HA	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:91:THR:HG23	2:E:124:THR:HA	1.97	0.45
2:D:45:ARG:NH1	2:D:111:TRP:CH2	2.83	0.44
1:B:328:ARG:HB3	2:D:107:THR:HB	1.99	0.43
1:A:137:LEU:HD11	1:B:146:PRO:HG2	2.00	0.43
1:B:335:LYS:HD2	4:D:401:MPD:H51	2.00	0.43
2:C:51:LEU:HD13	2:C:58:THR:HG22	1.99	0.43
1:A:121:HIS:CG	1:B:124:VAL:HG22	2.53	0.43
2:D:40:ALA:HB3	2:D:43:LYS:HB2	2.00	0.43
2:C:45:ARG:NH1	2:C:117:TRP:CE3	2.87	0.43
2:E:50:THR:HG22	2:E:59:TYR:HB2	2.01	0.42
1:B:112:LEU:HD21	2:C:106:GLY:HA3	2.02	0.42
2:E:40:ALA:HB3	2:E:43:LYS:HB2	2.03	0.41
2:C:50:THR:HG22	2:C:59:TYR:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/226 (76%)	171 (99%)	2 (1%)	63	81
1	B	141/226 (62%)	139 (99%)	2 (1%)	59	79
2	C	98/105 (93%)	97 (99%)	1 (1%)	68	84
2	D	95/105 (90%)	95 (100%)	0	100	100
2	E	93/105 (89%)	93 (100%)	0	100	100
All	All	600/767 (78%)	595 (99%)	5 (1%)	73	86

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	149	GLN
1	A	316	PHE
1	B	316	PHE
1	B	338	VAL
2	C	102	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	106	GLN
1	B	127	GLN
2	E	122	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPD	D	401	-	7,7,7	0.11	0	9,10,10	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	D	401	-	-	2/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	401	MPD	C2-C3-C4-O4
4	D	401	MPD	C2-C3-C4-C5

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	401	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	260/274 (94%)	1.24	62 (23%)	2 1	27, 49, 74, 82	0
1	B	266/274 (97%)	1.41	58 (21%)	2 2	29, 45, 78, 85	0
2	C	127/137 (92%)	0.45	6 (4%)	36 32	12, 28, 36, 45	2 (1%)
2	D	127/137 (92%)	1.00	15 (11%)	9 6	29, 39, 52, 70	0
2	E	126/137 (91%)	1.25	21 (16%)	4 3	34, 45, 59, 68	0
All	All	906/959 (94%)	1.15	162 (17%)	3 3	12, 42, 70, 85	2 (0%)

All (162) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	151	SER	7.7
1	B	149	GLN	6.6
1	B	74	MET	6.3
1	B	339	GLY	6.0
1	B	231	VAL	5.4
1	B	152	GLN	5.2
1	B	232	SER	5.1
1	B	233	GLN	4.9
1	B	153	GLN	4.9
1	B	338	VAL	4.8
2	E	9	GLY	4.7
2	D	9	GLY	4.7
1	A	234	THR	4.7
1	B	154	SER	4.5
2	E	8	GLY	4.5
1	B	150	SER	4.4
1	B	296	ASN	4.3
1	B	221	ARG	4.3
1	A	147	ILE	4.1
1	A	328	ARG	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	234	THR	4.0
1	A	149	GLN	3.9
2	E	126	SER	3.8
1	B	237	ARG	3.8
1	B	241	THR	3.8
2	C	1	GLN	3.6
1	A	333	LYS	3.6
2	E	11	LEU	3.6
1	A	244	PHE	3.6
1	B	156	HIS	3.5
1	A	330	LEU	3.5
1	B	236	ILE	3.5
1	A	231	VAL	3.4
1	B	148	GLY	3.4
1	A	226	PHE	3.3
2	D	55	GLY	3.3
1	B	295	ILE	3.3
2	D	108	ALA	3.3
2	C	120	GLY	3.3
2	E	1	GLN	3.2
2	C	27	ARG	3.2
1	A	241	THR	3.2
2	C	46	ASP	3.2
1	A	74	MET	3.2
1	A	293	ALA	3.1
1	A	242	ILE	3.1
2	E	51	LEU	3.1
1	B	177	ASN	3.1
1	A	262	TYR	3.1
1	B	334	THR	3.0
2	E	45	ARG	3.0
1	A	292	ASN	3.0
1	A	206	GLY	3.0
1	A	152	GLN	3.0
1	A	151	SER	3.0
1	A	294	GLY	3.0
1	B	294	GLY	3.0
1	A	295	ILE	3.0
1	B	226	PHE	3.0
1	B	141	ASN	3.0
1	A	148	GLY	2.9
1	B	216	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	75	LYS	2.9
1	B	230	GLN	2.9
1	B	225	VAL	2.8
2	D	40	ALA	2.8
1	A	332	LYS	2.8
1	B	224	LYS	2.8
1	A	235	LYS	2.8
1	A	326	TRP	2.7
1	B	332	LYS	2.7
1	B	242	ILE	2.7
1	B	331	SER	2.7
1	A	233	GLN	2.7
1	A	150	SER	2.7
2	D	7	SER	2.7
2	D	32	TYR	2.7
1	A	213	THR	2.7
1	B	293	ALA	2.7
1	A	75	LYS	2.6
1	B	229	LYS	2.6
1	B	147	ILE	2.6
1	A	108	ASN	2.6
1	A	327	LEU	2.6
2	C	2	VAL	2.6
2	E	49	ALA	2.6
1	B	164	PHE	2.5
1	A	331	SER	2.5
1	B	144	VAL	2.5
2	E	5	VAL	2.5
2	E	17	SER	2.5
1	B	228	LYS	2.5
1	A	80	ILE	2.5
1	A	153	GLN	2.5
1	B	290	GLY	2.4
1	B	239	ILE	2.4
1	A	154	SER	2.4
2	D	54	GLY	2.4
1	A	263	ILE	2.4
1	A	301	TYR	2.4
1	A	156	HIS	2.4
1	A	221	ARG	2.4
1	B	235	LYS	2.4
1	A	208	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	244	PHE	2.4
1	B	321	SER	2.4
1	B	159	LEU	2.4
1	A	247	ALA	2.4
1	A	89	VAL	2.3
1	B	179	ALA	2.3
2	E	42	GLY	2.3
1	A	141	ASN	2.3
1	A	92	GLY	2.3
1	A	324	GLY	2.3
1	A	243	LEU	2.3
1	B	336	GLU	2.3
1	B	337	GLU	2.3
2	C	42	GLY	2.3
1	A	236	ILE	2.3
1	A	260	PHE	2.3
2	E	44	GLU	2.3
1	A	246	LEU	2.2
1	A	240	SER	2.2
2	D	117	TRP	2.2
1	A	298	ARG	2.2
1	B	222	VAL	2.2
1	A	302	LYS	2.2
1	A	239	ILE	2.2
2	E	7	SER	2.2
2	E	47	PHE	2.2
1	B	142	ALA	2.2
2	D	11	LEU	2.2
1	A	299	GLU	2.2
2	E	46	ASP	2.2
2	D	12	VAL	2.1
2	D	1	GLN	2.1
2	D	42	GLY	2.1
1	B	272	ILE	2.1
1	A	215	PHE	2.1
1	A	144	VAL	2.1
2	E	12	VAL	2.1
2	E	18	LEU	2.1
1	B	301	TYR	2.1
2	D	127	SER	2.1
1	A	238	VAL	2.1
1	B	300	TRP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	16	GLY	2.1
2	D	39	GLU	2.1
2	E	39	GLU	2.1
1	A	228	LYS	2.1
2	E	71	SER	2.1
1	A	229	LYS	2.0
1	A	146	PRO	2.0
1	B	155	SER	2.0
1	B	238	VAL	2.0
2	E	14	ALA	2.0
1	B	292	ASN	2.0
1	B	335	LYS	2.0
1	A	245	ILE	2.0
1	A	255	ILE	2.0
2	D	58	THR	2.0
1	A	232	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	K	B	601	1/1	0.71	0.20	48,48,48,48	0
3	K	A	403	1/1	0.74	0.09	37,37,37,37	0
4	MPD	D	401	8/8	0.82	0.20	47,47,48,48	0
3	K	A	402	1/1	0.86	0.11	50,50,50,50	0
3	K	B	602	1/1	0.87	0.06	37,37,37,37	0
3	K	A	401	1/1	0.93	0.14	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.