



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 06:24 PM UTC

PDB ID : 2R0K / pdb_00002r0k
Title : Protease domain of HGFA with inhibitor Fab58
Authors : Eigenbrot, C.; Shia, S.
Deposited on : 2007-08-20
Resolution : 3.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1025 (3.56-3.48)
Clashscore	190562	1079 (3.56-3.48)
Ramachandran outliers	187476	1052 (3.56-3.48)
Sidechain outliers	187428	1053 (3.56-3.48)
RSRZ outliers	180081	1024 (3.56-3.48)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	10% 61% 22% • 16%
2	L	214	13% 66% 29% •
3	H	225	8% 62% 27% • • 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hepatocyte growth factor activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	239	Total	C	N	O	S	0	0	0
			1824	1156	316	339	13			

- Molecule 2 is a protein called antibody light chain of Fab58.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1637	1024	271	336	6			


- Molecule 3 is a protein called antibody heavy chain of Fab58, Fab portion only.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	213	Total	C	N	O	S	7	0	0
			1581	1001	266	308	6			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hepatocyte growth factor activator




- Molecule 2: antibody light chain of Fab58



- Molecule 3: antibody heavy chain of Fab58, Fab portion only





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	188.66Å 75.61Å 69.14Å 90.00° 92.67° 90.00°	Depositor
Resolution (Å)	50.00 – 3.51 50.00 – 3.51	Depositor EDS
% Data completeness (in resolution range)	87.7 (50.00-3.51) 87.8 (50.00-3.51)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.55 (at 3.48Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.251 , 0.312 0.258 , 0.312	Depositor DCC
R_{free} test set	526 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	47.6	Xtriage
Anisotropy	0.225	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.030 for -h,-k,l	Xtriage
F_o, F_c correlation	0.81	EDS
Total number of atoms	5042	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1878	0.84	3/2558 (0.1%)
2	L	0.52	1/1673 (0.1%)	0.85	2/2273 (0.1%)
3	H	0.74	2/1619 (0.1%)	0.92	7/2208 (0.3%)
All	All	0.59	3/5170 (0.1%)	0.87	12/7039 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	1	GLU	N-CA	-19.23	1.09	1.46
3	H	192	GLN	CB-CG	7.85	1.76	1.52
2	L	58	VAL	CA-CB	5.37	1.58	1.54

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	36	GLY	O-C-N	6.46	131.10	122.70
1	A	219	GLY	O-C-N	6.30	130.04	122.67
3	H	99	SER	N-CA-C	6.08	117.38	108.14
3	H	192	GLN	CA-CB-CG	-5.69	102.72	114.10
3	H	157	GLY	N-CA-C	-5.60	107.29	114.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1824	0	1739	37	0
2	L	1637	0	1586	46	0
3	H	1581	0	1550	65	0
All	All	5042	0	4875	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:192:GLN:CG	3:H:192:GLN:CB	1.76	1.64
3:H:153:SER:HA	3:H:154:TRP:HB2	1.41	1.03
1:A:217:ASP:HB2	3:H:28:THR:HG21	1.47	0.96
2:L:37:GLN:HB2	2:L:47:LEU:HD21	1.57	0.87
1:A:217:ASP:HB2	3:H:28:THR:CG2	2.07	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/283 (84%)	213 (90%)	22 (9%)	2 (1%)	16	50
2	L	212/214 (99%)	192 (91%)	16 (8%)	4 (2%)	6	33
3	H	209/225 (93%)	182 (87%)	21 (10%)	6 (3%)	3	25
All	All	658/722 (91%)	587 (89%)	59 (9%)	12 (2%)	6	34

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	187	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	H	98	PHE
3	H	153	SER
3	H	154	TRP
1	A	171	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/237 (84%)	187 (94%)	12 (6%)	17	44
2	L	188/188 (100%)	167 (89%)	21 (11%)	6	26
3	H	174/185 (94%)	154 (88%)	20 (12%)	5	25
All	All	561/610 (92%)	508 (91%)	53 (9%)	8	31

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	183	LYS
3	H	32	SER
3	H	186	SER
2	L	185	ASP
2	L	214	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	198	HIS
3	H	155	ASN
2	L	210	ASN
3	H	200	HIS
3	H	39	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	239/283 (84%)	1.90	27 (11%)	10 6	0, 7, 45, 63	15 (6%)
2	L	214/214 (100%)	1.47	27 (12%)	8 6	3, 15, 43, 55	7 (3%)
3	H	213/225 (94%)	0.71	17 (7%)	18 12	1, 11, 36, 62	3 (1%)
All	All	666/722 (92%)	1.38	71 (10%)	11 7	0, 11, 42, 63	25 (3%)

The worst 5 of 71 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	187	CYS	44.0
1	A	186	ASP	38.8
2	L	189	HIS	30.0
1	A	219	GLY	28.9
1	A	221	ARG	26.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.