



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 27, 2026 – 08:42 PM UTC

PDB ID : 4R0A / pdb_00004r0a
Title : Crystal structure of human TLR8 in complex with uridine mononucleoside
Authors : Tanji, H.; Ohto, U.; Shimizu, T.
Deposited on : 2014-07-30
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6567 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	748	Total	C	N	O	S	0	0	0
			5994	3833	1020	1122	19			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	ARG	-	expression tag	UNP Q9NR97
A	24	SER	-	expression tag	UNP Q9NR97
A	25	PRO	-	expression tag	UNP Q9NR97
A	26	TRP	-	expression tag	UNP Q9NR97
A	828	GLU	-	expression tag	UNP Q9NR97
A	829	PHE	-	expression tag	UNP Q9NR97
A	830	LEU	-	expression tag	UNP Q9NR97
A	831	VAL	-	expression tag	UNP Q9NR97
A	832	PRO	-	expression tag	UNP Q9NR97
A	833	ARG	-	expression tag	UNP Q9NR97

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(2-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	5	Total	C	N	O	0	0	0
			61	34	2	25			


- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is URIDINE (CCD ID: URI) (formula: $C_9H_{12}N_2O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			17	9	2	6		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	330	Total	O	0	0
			330	330		

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	105.44Å 133.32Å 120.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.58 – 1.90 48.58 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.3 (48.58-1.90) 98.3 (48.58-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.200 , 0.260 0.207 , 0.265	Depositor DCC
R_{free} test set	3309 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	27.2	Xtriage
Anisotropy	0.134	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6567	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.76	0	17,19,21	1.55	3 (17%)
2	NAG	B	2	2	14,14,15	0.85	1 (7%)	17,19,21	1.80	3 (17%)
2	BMA	B	3	2	11,11,12	1.04	1 (9%)	15,15,17	1.87	4 (26%)
2	MAN	B	4	2	11,11,12	1.08	0	15,15,17	1.03	1 (6%)
2	MAN	B	5	2	11,11,12	0.68	0	13,15,17	4.36	4 (30%)
3	NAG	C	1	1,3	14,14,15	1.02	1 (7%)	17,19,21	1.36	2 (11%)
3	NAG	C	2	3	14,14,15	0.88	1 (7%)	17,19,21	1.36	2 (11%)
4	NAG	D	1	4,1	14,14,15	0.61	0	17,19,21	1.21	1 (5%)
4	NAG	D	2	4	14,14,15	0.90	0	17,19,21	1.52	3 (17%)
4	BMA	D	3	4	11,11,12	0.69	0	15,15,17	1.23	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	2/2/18/22	0/1/1/1
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
4	NAG	D	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	BMA	O5-C1	-2.46	1.39	1.43
3	C	1	NAG	O5-C5	-2.35	1.38	1.43
2	B	2	NAG	O5-C1	-2.16	1.40	1.43
3	C	2	NAG	C1-C2	2.08	1.55	1.52

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	5	MAN	O1-C1-O5	-14.29	76.05	108.34
2	B	2	NAG	O5-C1-C2	-5.71	102.45	111.29
2	B	5	MAN	O1-C1-C2	-4.60	90.37	111.39
2	B	3	BMA	C1-C2-C3	4.47	116.16	109.64
4	D	1	NAG	C1-O5-C5	3.81	117.30	112.19
2	B	1	NAG	O6-C6-C5	-3.67	98.85	111.33
2	B	1	NAG	O5-C1-C2	-3.05	106.58	111.29
2	B	3	BMA	O2-C2-C3	-3.02	103.89	110.15
3	C	1	NAG	C1-C2-N2	-3.00	105.70	110.43
3	C	1	NAG	O5-C1-C2	-2.88	106.84	111.29
3	C	2	NAG	O6-C6-C5	-2.81	101.78	111.33
4	D	3	BMA	C2-C3-C4	-2.74	106.05	110.86
2	B	3	BMA	O4-C4-C5	2.72	116.03	109.32
2	B	5	MAN	O5-C5-C4	2.54	114.27	109.70
3	C	2	NAG	C3-C4-C5	-2.53	105.64	110.23
4	D	2	NAG	O7-C7-N2	2.49	126.38	121.98
2	B	3	BMA	O4-C4-C3	-2.46	104.58	110.38
2	B	1	NAG	O5-C5-C6	2.43	112.40	107.66
2	B	5	MAN	C2-C3-C4	-2.19	107.51	110.67
2	B	4	MAN	C1-O5-C5	2.16	115.08	112.19
4	D	2	NAG	C4-C3-C2	-2.15	107.86	111.02
2	B	2	NAG	C1-O5-C5	-2.14	109.31	112.19
4	D	2	NAG	O5-C1-C2	-2.06	108.10	111.29
2	B	2	NAG	O5-C5-C6	2.02	111.59	107.66

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	5	MAN	O5-C5-C6-O6
2	B	5	MAN	C4-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







4.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	NAG	A	904	1	14,14,15	1.00	1 (7%)	17,19,21	1.43	2 (11%)
6	NAG	A	913	1	14,14,15	0.72	0	17,19,21	2.03	7 (41%)
6	NAG	A	902	1	14,14,15	0.57	0	17,19,21	0.99	1 (5%)
6	NAG	A	905	1	14,14,15	0.69	0	17,19,21	2.57	4 (23%)
6	NAG	A	903	1	14,14,15	0.66	0	17,19,21	1.11	1 (5%)
6	NAG	A	917	1	14,14,15	1.39	2 (14%)	17,19,21	2.05	4 (23%)
5	URI	A	901	-	18,18,18	0.27	0	26,26,26	0.48	0
6	NAG	A	918	1	14,14,15	0.99	1 (7%)	17,19,21	1.81	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	904	1	-	0/6/23/26	0/1/1/1
6	NAG	A	913	1	-	2/6/23/26	0/1/1/1
6	NAG	A	902	1	-	0/6/23/26	0/1/1/1
6	NAG	A	905	1	-	0/6/23/26	0/1/1/1
6	NAG	A	903	1	-	2/6/23/26	0/1/1/1
6	NAG	A	917	1	-	0/6/23/26	0/1/1/1
5	URI	A	901	-	-	4/6/22/22	0/2/2/2
6	NAG	A	918	1	-	2/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	917	NAG	C2-N2	-2.96	1.41	1.46
6	A	917	NAG	O5-C5	-2.71	1.38	1.43
6	A	904	NAG	O5-C1	-2.53	1.39	1.43
6	A	918	NAG	O5-C1	-2.41	1.39	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	905	NAG	C1-O5-C5	8.90	124.11	112.19
6	A	917	NAG	C1-C2-N2	-6.44	100.28	110.43
6	A	918	NAG	O5-C1-C2	-5.29	103.11	111.29
6	A	913	NAG	C4-C3-C2	-3.81	105.44	111.02
6	A	913	NAG	O5-C1-C2	-3.58	105.75	111.29
6	A	918	NAG	C1-C2-N2	-3.47	104.97	110.43
6	A	913	NAG	O7-C7-C8	-3.27	116.24	122.05
6	A	904	NAG	O5-C1-C2	-3.21	106.33	111.29
6	A	905	NAG	C4-C3-C2	-3.08	106.50	111.02
6	A	902	NAG	C1-O5-C5	2.93	116.11	112.19
6	A	913	NAG	O7-C7-N2	2.87	127.05	121.98
6	A	913	NAG	C1-C2-N2	2.79	114.83	110.43
6	A	905	NAG	C2-N2-C7	-2.77	119.18	122.90
6	A	904	NAG	O5-C5-C4	-2.75	104.14	110.83
6	A	917	NAG	O6-C6-C5	-2.39	103.20	111.33
6	A	918	NAG	C8-C7-N2	-2.34	112.24	116.12
6	A	913	NAG	O5-C5-C4	-2.20	105.47	110.83
6	A	913	NAG	C1-O5-C5	2.20	115.13	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	917	NAG	C8-C7-N2	-2.18	112.50	116.12
6	A	917	NAG	C4-C3-C2	-2.17	107.84	111.02
6	A	903	NAG	O5-C5-C4	-2.13	105.63	110.83
6	A	905	NAG	C3-C4-C5	-2.02	106.56	110.23

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	903	NAG	O5-C5-C6-O6
6	A	903	NAG	C4-C5-C6-O6
6	A	918	NAG	O5-C5-C6-O6
6	A	918	NAG	C4-C5-C6-O6
6	A	913	NAG	C1-C2-N2-C7
5	A	901	URI	O4'-C1'-N1-C6
6	A	913	NAG	C3-C2-N2-C7
5	A	901	URI	C2'-C1'-N1-C6
5	A	901	URI	O4'-C1'-N1-C2
5	A	901	URI	C2'-C1'-N1-C2

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data ⓘ

5.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	748/811 (92%)	0.49	68 (9%) 15 15	20, 31, 76, 106	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	780	ILE	6.1
1	A	791	LEU	5.2
1	A	702	PHE	5.0
1	A	758	THR	4.8
1	A	815	VAL	4.6
1	A	764	LEU	4.4
1	A	760	THR	4.4
1	A	86	LEU	4.1
1	A	793	VAL	4.1
1	A	778	CYS	4.0
1	A	786	TRP	3.9
1	A	761	THR	3.9
1	A	781	GLY	3.9
1	A	814	ILE	3.9
1	A	85	GLY	3.8
1	A	817	LEU	3.8
1	A	765	SER	3.7
1	A	100	VAL	3.6
1	A	802	ILE	3.6
1	A	762	THR	3.6
1	A	783	PHE	3.6
1	A	756	LEU	3.6
1	A	64	TYR	3.5
1	A	749	LYS	3.5
1	A	732	LEU	3.5
1	A	806	PRO	3.5
1	A	757	GLU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	779	ASP	3.3
1	A	236	TYR	3.2
1	A	45	VAL	3.0
1	A	774	PHE	3.0
1	A	470	PHE	2.9
1	A	748	LEU	2.9
1	A	123	ASN	2.8
1	A	799	VAL	2.8
1	A	61	VAL	2.8
1	A	807	GLY	2.8
1	A	752	ASN	2.6
1	A	726	HIS	2.6
1	A	751	ILE	2.6
1	A	701	LEU	2.6
1	A	811	GLY	2.6
1	A	795	ILE	2.6
1	A	735	VAL	2.6
1	A	808	ASP	2.5
1	A	678	PHE	2.5
1	A	46	ILE	2.4
1	A	84	GLN	2.4
1	A	813	SER	2.3
1	A	776	CYS	2.3
1	A	57	VAL	2.3
1	A	727	LEU	2.3
1	A	775	GLU	2.2
1	A	803	CYS	2.2
1	A	122	LEU	2.2
1	A	582	ASN	2.2
1	A	812	LYS	2.2
1	A	730	GLY	2.2
1	A	34	TYR	2.2
1	A	724	ILE	2.1
1	A	773	PRO	2.1
1	A	810	ARG	2.1
1	A	164	GLU	2.1
1	A	113	GLY	2.1
1	A	784	ARG	2.1
1	A	42	ASN	2.1
1	A	41	GLN	2.0
1	A	281	PHE	2.0

5.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	B	5	11/12	0.78	0.12	56,61,74,74	0
2	MAN	B	4	11/12	0.81	0.16	35,47,52,53	0
4	BMA	D	3	11/12	0.86	0.10	34,39,50,52	0
3	NAG	C	2	14/15	0.87	0.09	32,38,44,49	0
4	NAG	D	2	14/15	0.93	0.07	23,26,32,39	0
2	BMA	B	3	11/12	0.93	0.08	29,37,49,50	0
2	NAG	B	1	14/15	0.94	0.06	19,21,24,28	0
3	NAG	C	1	14/15	0.95	0.06	20,22,25,26	0
2	NAG	B	2	14/15	0.95	0.07	22,24,32,33	0
4	NAG	D	1	14/15	0.96	0.06	20,22,23,24	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	A	903	14/15	0.81	0.13	57,63,69,81	0
6	NAG	A	904	14/15	0.82	0.13	44,49,58,58	0
6	NAG	A	902	14/15	0.83	0.11	39,64,72,73	0
6	NAG	A	918	14/15	0.83	0.13	44,53,63,63	0
6	NAG	A	905	14/15	0.84	0.12	45,55,63,67	0
6	NAG	A	913	14/15	0.87	0.12	35,45,57,59	0
6	NAG	A	917	14/15	0.96	0.06	21,23,28,31	0
5	URI	A	901	17/17	0.96	0.06	21,25,35,37	0

5.5 Other polymers [i](#)

There are no such residues in this entry.