



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 01:44 PM EDT

PDB ID : 4R0V / pdb_00004r0v
Title : [FeFe]-hydrogenase Oxygen Inactivation is Initiated by the Modification and Degradation of the H cluster 2Fe Subcluster
Authors : Swanson, S.D.; Ratzloff, M.W.; Mulder, D.W.; Artz, J.H.; Ghose, S.; Hoffman, A.; White, S.; Zadvornyy, O.A.; Broderick, J.B.; Bothner, B.; King, P.W.; Peters, J.W.
Deposited on : 2014-08-01
Resolution : 2.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	514	9% 68% 11% • 18%
1	B	514	7% 67% 12% • 20%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ARS	A	604	-	-	X	-
4	ARS	B	604	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fe-hydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	S	0	2	0
			3179	1998	550	606	25			
1	B	410	Total	C	N	O	S	0	2	0
			3121	1963	539	594	25			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	498	LEU	-	expression tag	UNP Q9FYU1
A	499	GLU	-	expression tag	UNP Q9FYU1
A	500	ILE	-	expression tag	UNP Q9FYU1
A	501	GLU	-	expression tag	UNP Q9FYU1
A	502	GLY	-	expression tag	UNP Q9FYU1
A	503	ARG	-	expression tag	UNP Q9FYU1
A	504	GLN	-	expression tag	UNP Q9FYU1
A	505	LEU	-	expression tag	UNP Q9FYU1
A	506	GLY	-	expression tag	UNP Q9FYU1
A	507	TRP	-	expression tag	UNP Q9FYU1
A	508	SER	-	expression tag	UNP Q9FYU1
A	509	HIS	-	expression tag	UNP Q9FYU1
A	510	PRO	-	expression tag	UNP Q9FYU1
A	511	GLN	-	expression tag	UNP Q9FYU1
A	512	PHE	-	expression tag	UNP Q9FYU1
A	513	GLU	-	expression tag	UNP Q9FYU1
A	514	LYS	-	expression tag	UNP Q9FYU1
B	498	LEU	-	expression tag	UNP Q9FYU1
B	499	GLU	-	expression tag	UNP Q9FYU1
B	500	ILE	-	expression tag	UNP Q9FYU1
B	501	GLU	-	expression tag	UNP Q9FYU1
B	502	GLY	-	expression tag	UNP Q9FYU1
B	503	ARG	-	expression tag	UNP Q9FYU1
B	504	GLN	-	expression tag	UNP Q9FYU1
B	505	LEU	-	expression tag	UNP Q9FYU1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	506	GLY	-	expression tag	UNP Q9FYU1
B	507	TRP	-	expression tag	UNP Q9FYU1
B	508	SER	-	expression tag	UNP Q9FYU1
B	509	HIS	-	expression tag	UNP Q9FYU1
B	510	PRO	-	expression tag	UNP Q9FYU1
B	511	GLN	-	expression tag	UNP Q9FYU1
B	512	PHE	-	expression tag	UNP Q9FYU1
B	513	GLU	-	expression tag	UNP Q9FYU1
B	514	LYS	-	expression tag	UNP Q9FYU1

- Molecule 2 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		
3	B	2	Total	Cl	0	0
			2	2		

- Molecule 4 is ARSENIC (CCD ID: ARS) (formula: As).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	As 1	0	0
4	B	1	Total 1	As 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	147	Total 147	O 147	0	0
5	B	146	Total 146	O 146	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.64Å 70.96Å 94.69Å 90.00° 91.91° 90.00°	Depositor
Resolution (Å)	35.00 – 2.29 35.00 – 2.29	Depositor EDS
% Data completeness (in resolution range)	88.2 (35.00-2.29) 88.2 (35.00-2.29)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.206 , 0.252 0.217 , 0.255	Depositor DCC
R_{free} test set	2098 reflections (4.50%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtriage
Anisotropy	0.371	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.037 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6615	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 96.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.0524e-10. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ARS, SF4, CL, CSO, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.23	9/3220 (0.3%)	1.21	13/4357 (0.3%)
1	B	1.21	6/3161 (0.2%)	1.25	13/4276 (0.3%)
All	All	1.22	15/6381 (0.2%)	1.23	26/8633 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	4
All	All	0	6

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	150	THR	C-N	9.29	1.46	1.33
1	A	88	CSD	C-N	9.08	1.46	1.33
1	B	89	VAL	C-N	-8.03	1.22	1.33
1	A	151	GLU	C-N	-7.45	1.23	1.33
1	A	127	VAL	N-CA	-6.99	1.37	1.46

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	CSD	O-C-N	-18.30	93.73	123.00
1	B	88	CSD	CA-C-N	15.40	149.42	121.70
1	B	88	CSD	C-N-CA	15.40	149.42	121.70
1	A	88	CSD	O-C-N	-11.38	104.79	123.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	151	GLU	O-C-N	-11.19	110.26	122.12

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	238	CSO	Mainchain
1	A	87	VAL	Mainchain
1	B	169[A]	CSO	Mainchain
1	B	169[B]	CSO	Mainchain
1	B	87	VAL	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3179	0	3162	69	0
1	B	3121	0	3108	68	0
2	A	8	0	0	1	0
2	B	8	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	9	0
4	B	1	0	0	10	0
5	A	147	0	0	5	0
5	B	146	0	0	5	0
All	All	6615	0	6270	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance ($\text{\AA}$)	Clash overlap ($\text{\AA}$)
1:A:361:THR:CG2	1:A:362:PRO:HD2	1.51	1.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:361:THR:HG23	1:A:362:PRO:CD	1.52	1.38
1:B:169[B]:CSO:SG	1:B:415[B]:MET:HE1	1.70	1.29
1:A:475:HIS:NE2	4:B:604:ARS:AS	2.40	1.14
4:A:604:ARS:AS	1:B:475:HIS:NE2	2.40	1.14

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	411/514 (80%)	401 (98%)	9 (2%)	1 (0%)	43	55
1	B	403/514 (78%)	392 (97%)	11 (3%)	0	100	100
All	All	814/1028 (79%)	793 (97%)	20 (2%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	446	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/399 (84%)	329 (99%)	4 (1%)	63	79

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	328/399 (82%)	325 (99%)	3 (1%)	70	84
All	All	661/798 (83%)	654 (99%)	7 (1%)	65	81

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	348	HIS
1	B	143	SER
1	B	315	ARG
1	B	202	VAL
1	A	341	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	86	HIS
1	B	195	GLN
1	B	484	HIS
1	B	440	GLN
1	B	456	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	B	88	1	4,7,8	2.95	1 (25%)	1,8,10	3.15	1 (100%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	A	238	1	3,6,7	1.01	0	1,6,8	1.27	0
1	CSO	B	238	1	3,6,7	1.01	0	1,6,8	0.50	0
1	CSO	A	169[A]	-	3,6,7	1.10	0	1,6,8	1.34	0
1	CSO	B	169[A]	-	3,6,7	1.00	0	1,6,8	0.49	0
1	CSO	B	169[B]	-	3,6,7	0.83	0	1,6,8	1.44	0
1	CSO	A	169[B]	-	3,6,7	1.90	1 (33%)	1,6,8	2.05	1 (100%)
1	CSD	A	88	1	4,7,8	3.17	1 (25%)	1,8,10	3.68	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	B	88	1	-	1/2/6/8	-
1	CSO	A	238	1	-	0/1/5/7	-
1	CSO	B	238	1	-	1/1/5/7	-
1	CSO	A	169[A]	-	-	0/1/5/7	-
1	CSO	B	169[A]	-	-	1/1/5/7	-
1	CSO	B	169[B]	-	-	0/1/5/7	-
1	CSO	A	169[B]	-	-	1/1/5/7	-
1	CSD	A	88	1	-	1/2/6/8	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	88	CSD	OD1-SG	-5.74	1.42	1.47
1	B	88	CSD	OD1-SG	-5.33	1.42	1.47
1	A	169[B]	CSO	CB-CA	2.69	1.60	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	CSD	OD1-SG-CB	3.68	112.37	105.60
1	B	88	CSD	OD1-SG-CB	3.15	111.40	105.60
1	A	169[B]	CSO	CB-CA-C	-2.05	105.22	110.80

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	88	CSD	CA-CB-SG-OD1
1	A	169[B]	CSO	N-CA-CB-SG
1	B	88	CSD	CA-CB-SG-OD1
1	B	238	CSO	N-CA-CB-SG
1	B	169[A]	CSO	N-CA-CB-SG

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	238	CSO	2	0
1	B	238	CSO	2	0
1	B	169[B]	CSO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SF4	B	601	1	0,12,12	-	-	-		
2	SF4	A	601	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	B	601	1	-	-	0/6/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	A	601	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	SF4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	87:VAL	C	88:CSD	N	1.61

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	416/514 (80%)	0.61	47 (11%)	10 11	23, 45, 83, 153	3 (0%)
1	B	407/514 (79%)	0.44	36 (8%)	15 17	23, 44, 78, 123	3 (0%)
All	All	823/1028 (80%)	0.53	83 (10%)	12 14	23, 44, 80, 153	6 (0%)

The worst 5 of 83 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	240	ASP	11.2
1	A	241	ALA	11.2
1	A	391	GLY	9.4
1	B	363	GLY	8.8
1	A	237	PHE	8.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	A	238	7/8	0.49	0.33	45,52,59,60	0
1	CSO	B	238	7/8	0.51	0.34	45,52,59,60	0
1	CSO	B	169[B]	7/8	0.82	0.18	22,33,35,38	4
1	CSO	B	169[A]	7/8	0.82	0.18	32,33,38,47	4
1	CSD	B	88	8/9	0.86	0.15	30,36,44,49	3
1	CSD	A	88	8/9	0.86	0.15	31,35,43,48	3
1	CSO	A	169[B]	7/8	0.92	0.11	25,35,36,47	4
1	CSO	A	169[A]	7/8	0.92	0.11	30,34,38,43	4

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ARS	A	604	1/1	0.41	0.35	69,69,69,69	0
2	SF4	A	601	8/8	0.69	0.18	24,27,40,41	0
4	ARS	B	604	1/1	0.86	0.13	71,71,71,71	0
2	SF4	B	601	8/8	0.87	0.10	24,27,40,41	0
3	CL	A	602	1/1	0.94	0.27	73,73,73,73	0
3	CL	B	603	1/1	0.95	0.19	78,78,78,78	0
3	CL	B	602	1/1	0.98	0.05	38,38,38,38	0
3	CL	A	603	1/1	0.99	0.05	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.