



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:03 AM UTC

PDB ID : 5R0F / pdb_00005r0f
Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment F2X-Entry D06, DMSO-free
Authors : Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.; Weiss, M.S.
Deposited on : 2020-02-12
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	25% 71% 18% 8%
2	B	308	30% 78% 18% • •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	1994	1278	334	371	11	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2568	1645	420	483	20	0	9	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	170	SER	LEU	conflict	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ASP	deletion	UNP P32357

- Molecule 3 is (3-methoxyphenyl)(pyrrolidin-1-yl)methanone (CCD ID: R9G) (formula: $C_{12}H_{15}NO_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	12	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	45	Total	O	0	0
			45	45		
4	B	21	Total	O	0	0
			21	21		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-mRNA-splicing factor 8



• Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.59Å 82.24Å 91.64Å 90.00° 108.25° 90.00°	Depositor
Resolution (Å)	22.34 – 1.97 22.34 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.5 (22.34-1.97) 99.4 (22.34-1.97)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.95 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.261 , 0.304 0.276 , 0.315	Depositor DCC
R_{free} test set	2100 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 70.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4643	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: R9G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	2/2041 (0.1%)	1.11	10/2765 (0.4%)
2	B	0.69	0/2638	0.85	1/3563 (0.0%)
All	All	0.82	2/4679 (0.0%)	0.97	11/6328 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1887	GLY	C-O	-8.97	1.11	1.24
1	A	1882	LEU	C-O	-5.09	1.18	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1940	MET	CG-SD-CE	-11.06	76.56	100.90
1	A	1986	MET	CA-C-N	7.99	135.28	122.50
1	A	1986	MET	C-N-CA	7.99	135.28	122.50
1	A	1987	VAL	N-CA-CB	-7.61	100.06	110.56
1	A	2035	LYS	O-C-N	-6.91	114.53	122.68
2	B	235	GLY	N-CA-C	-5.96	105.72	114.90
1	A	1973	LYS	CB-CA-C	5.92	120.91	110.85
1	A	1993	ASP	CB-CA-C	5.83	122.02	110.42
1	A	2039	LEU	N-CA-CB	-5.77	102.64	110.67
1	A	1990	ASN	CA-CB-CG	-5.50	107.10	112.60
1	A	1986	MET	CB-CG-SD	-5.22	97.04	112.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1994	0	2020	31	0
2	B	2568	0	2441	39	0
3	A	15	0	0	0	0
4	A	45	0	0	1	0
4	B	21	0	0	1	0
All	All	4643	0	4461	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:96:PHE:HB2	2:B:102:MET:HE3	1.56	0.88
1:A:1842:GLU:OE1	4:A:2601:HOH:O	1.92	0.87
2:B:1:MET:N	4:B:401:HOH:O	2.09	0.86
1:A:1918[B]:SER:OG	1:A:1922:ARG:NH2	2.11	0.82
1:A:2062:GLU:O	1:A:2066:LYS:HG2	1.80	0.81
2:B:70:GLN:HB3	2:B:81:MET:HE2	1.61	0.80
1:A:1951:PHE:HB3	1:A:1954:ILE:HD12	1.69	0.74
2:B:96:PHE:CB	2:B:102:MET:HE3	2.18	0.72
1:A:1910:LYS:O	1:A:1940:MET:HE1	1.94	0.68
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.19	0.68
2:B:277:GLU:CD	2:B:277:GLU:H	2.05	0.64
1:A:1878:CYS:SG	1:A:1969:MET:HE3	2.40	0.62
2:B:47:GLN:NE2	2:B:52:SER:O	2.32	0.62
1:A:1962:ARG:O	1:A:1962:ARG:HG2	1.99	0.61
2:B:287:ARG:O	2:B:291:ILE:HD13	1.99	0.61
1:A:1930:PRO:O	1:A:1954:ILE:HG12	2.03	0.59
1:A:2044:THR:OG1	1:A:2047:GLN:HG3	2.07	0.55
1:A:2028:SER:O	1:A:2030:PRO:HD3	2.07	0.54
1:A:1910:LYS:HA	1:A:1940:MET:SD	2.50	0.51
2:B:141:ASP:OD1	2:B:141:ASP:C	2.53	0.51
2:B:41[B]:VAL:HG12	2:B:123:MET:HG2	1.93	0.51
2:B:144:MET:HE2	2:B:176:LEU:HG	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:16:GLY:HA3	2:B:45:HIS:CE1	2.46	0.50
2:B:203[A]:TYR:CZ	2:B:207:THR:HG21	2.45	0.50
2:B:61:ASP:HB2	2:B:123:MET:HG3	1.94	0.50
1:A:1902:GLN:HB2	1:A:1905:LEU:CD2	2.43	0.49
2:B:53:SER:O	2:B:54[A]:MET:HB3	2.12	0.49
2:B:59:TRP:HH2	2:B:229:LEU:CD1	2.26	0.49
1:A:1848:ILE:H	1:A:1931[A]:LYS:NZ	2.11	0.48
2:B:251:CYS:SG	2:B:292:CYS:HB3	2.53	0.48
1:A:1891:LEU:HD22	1:A:1974:LEU:HD22	1.95	0.48
2:B:70:GLN:CB	2:B:81:MET:HE2	2.38	0.48
1:A:2052:GLU:O	1:A:2056[B]:ARG:HG3	2.14	0.48
1:A:1951:PHE:CB	1:A:1954:ILE:HD12	2.42	0.47
2:B:140:VAL:HG12	2:B:141:ASP:N	2.29	0.47
2:B:59:TRP:HH2	2:B:229:LEU:HD12	1.80	0.46
2:B:251:CYS:O	2:B:296:SER:HB2	2.15	0.46
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ1	1.62	0.45
1:A:1835:MET:HE3	1:A:1835:MET:HB3	1.77	0.45
2:B:250:ILE:HG21	2:B:264:LEU:HD22	1.98	0.45
2:B:195:MET:HG2	2:B:199:LEU:HD12	1.99	0.45
1:A:2021:SER:HB3	1:A:2058:LEU:HD12	1.99	0.44
2:B:6:PHE:CE2	2:B:69:ILE:HG21	2.53	0.44
1:A:2064:GLY:O	1:A:2068:ASN:N	2.45	0.44
1:A:1941:LEU:O	1:A:1945:GLU:HB2	2.18	0.44
2:B:18:ASP:OD2	2:B:42:HIS:ND1	2.42	0.44
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.52	0.44
1:A:1910:LYS:HG2	1:A:1940:MET:SD	2.57	0.44
1:A:1859:ARG:HB2	1:A:1875:ILE:HG13	2.00	0.44
1:A:2021:SER:HB3	1:A:2058:LEU:CD1	2.48	0.44
2:B:285:ASN:O	2:B:286:GLU:C	2.61	0.43
1:A:1933:ILE:HD12	1:A:1948:MET:HE2	2.00	0.43
1:A:1851:PHE:O	1:A:1881:THR:HA	2.19	0.43
2:B:231:ALA:O	2:B:235:GLY:N	2.48	0.43
2:B:267:ILE:O	2:B:271:GLN:HG3	2.18	0.43
2:B:277:GLU:CD	2:B:277:GLU:N	2.75	0.43
2:B:214:PHE:O	2:B:215:LYS:HB2	2.19	0.42
2:B:70:GLN:OE1	2:B:81:MET:HE1	2.19	0.42
1:A:1975:SER:O	1:A:1979[A]:MET:SD	2.78	0.42
2:B:10:PRO:O	2:B:26:GLU:HG3	2.19	0.41
2:B:214:PHE:O	2:B:215:LYS:CB	2.68	0.41
2:B:285:ASN:OD1	2:B:288:VAL:HG23	2.20	0.41
2:B:15:ILE:O	2:B:21:SER:HA	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1902:GLN:HB2	1:A:1905:LEU:HD21	2.02	0.41
2:B:118:THR:O	2:B:119:GLU:C	2.63	0.41
2:B:277:GLU:O	2:B:280:SER:OG	2.36	0.40
2:B:292:CYS:HA	2:B:296:SER:HB3	2.02	0.40
1:A:2056[B]:ARG:O	1:A:2060:LEU:HG	2.20	0.40
2:B:225:GLN:O	2:B:229:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/258 (96%)	239 (97%)	8 (3%)	0	100	100
2	B	305/308 (99%)	285 (93%)	19 (6%)	1 (0%)	36	27
All	All	552/566 (98%)	524 (95%)	27 (5%)	1 (0%)	43	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	215	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/233 (97%)	216 (96%)	9 (4%)	28	17
2	B	286/284 (101%)	279 (98%)	7 (2%)	43	35
All	All	511/517 (99%)	495 (97%)	16 (3%)	35	26

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1835	MET
1	A	1847	ASP
1	A	1889	LEU
1	A	1931[A]	LYS
1	A	1931[B]	LYS
1	A	1962	ARG
1	A	1977	VAL
1	A	1986	MET
1	A	2021	SER
2	B	53	SER
2	B	77	LEU
2	B	133	ASP
2	B	174	HIS
2	B	280	SER
2	B	315	GLU
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	R9G	A	2501	-	16,16,16	2.15	6 (37%)	21,21,21	0.98	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	R9G	A	2501	-	-	0/10/17/17	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2501	R9G	C11-N	-3.93	1.39	1.47
3	A	2501	R9G	C10-C9	-3.82	1.23	1.48
3	A	2501	R9G	C8-N	-3.62	1.39	1.47
3	A	2501	R9G	O1-C7	-2.89	1.16	1.22
3	A	2501	R9G	O-C	-2.58	1.35	1.42
3	A	2501	R9G	C2-C1	-2.31	1.34	1.38

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2501	R9G	O-C1-C2	-2.43	108.26	119.82
3	A	2501	R9G	O-C1-C6	2.15	130.26	120.00


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	237/258 (91%)	1.56	65 (27%)	1 1	26, 73, 116, 151	12 (5%)
2	B	300/308 (97%)	1.65	93 (31%)	1 1	33, 84, 140, 178	9 (3%)
All	All	537/566 (94%)	1.61	158 (29%)	1 1	26, 80, 130, 178	21 (3%)

All (158) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1878	CYS	9.5
2	B	54[A]	MET	9.0
2	B	22	PHE	6.0
1	A	1969	MET	5.8
1	A	1979[A]	MET	5.8
1	A	1890	PHE	5.7
1	A	1940	MET	5.2
2	B	89	PHE	5.2
1	A	1879	ILE	4.8
1	A	1840	TYR	4.8
1	A	2015	LEU	4.6
2	B	203[A]	TYR	4.6
2	B	313	TYR	4.5
2	B	316	LEU	4.4
1	A	2017[A]	THR	4.4
1	A	2069	VAL	4.4
2	B	188	ALA	4.3
1	A	2037	TYR	4.0
2	B	170	SER	3.9
2	B	179	THR	3.9
2	B	250	ILE	3.9
1	A	1991	ILE	3.8
2	B	275	LEU	3.7
1	A	1857	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	137	PHE	3.7
2	B	253	SER	3.6
2	B	173	ALA	3.6
2	B	221	PHE	3.6
1	A	1891	LEU	3.6
1	A	1941	LEU	3.5
2	B	103	VAL	3.5
2	B	101	MET	3.4
1	A	1834	ALA	3.4
1	A	2067	TYR	3.4
1	A	1934	ILE	3.3
2	B	291	ILE	3.3
1	A	2032	ILE	3.3
1	A	1843	LEU	3.3
2	B	243	TRP	3.3
1	A	1971	ILE	3.2
2	B	119	GLU	3.2
1	A	1987	VAL	3.2
1	A	1988	LEU	3.2
2	B	1	MET	3.2
2	B	249	LEU	3.2
2	B	129	ILE	3.1
2	B	180	VAL	3.1
2	B	198	PHE	3.1
2	B	282	ILE	3.1
2	B	233	PHE	3.1
2	B	205	LEU	3.0
2	B	152	LEU	3.0
1	A	2043	PHE	3.0
2	B	55	ARG	3.0
2	B	11	ILE	3.0
1	A	1881	THR	3.0
1	A	2033	THR	3.0
2	B	20	TYR	2.9
2	B	53	SER	2.9
2	B	295	SER	2.9
1	A	2061	THR	2.9
2	B	106	PRO	2.9
2	B	172	PRO	2.9
1	A	1989	PHE	2.9
1	A	2051	ILE	2.9
1	A	1876	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	130	VAL	2.8
2	B	96	PHE	2.8
2	B	214	PHE	2.8
2	B	267	ILE	2.8
1	A	1860	VAL	2.8
1	A	1931[A]	LYS	2.8
2	B	71	TYR	2.8
2	B	294	TYR	2.8
1	A	2065	ARG	2.7
1	A	1866	PHE	2.7
2	B	74	LYS	2.7
2	B	175	SER	2.7
2	B	193	HIS	2.7
2	B	120	PHE	2.7
1	A	2035	LYS	2.7
2	B	147	VAL	2.7
1	A	1919	ALA	2.7
2	B	178	TYR	2.7
2	B	279	TYR	2.7
1	A	2060	LEU	2.7
1	A	2066	LYS	2.6
1	A	1889	LEU	2.6
2	B	262	ASP	2.6
2	B	77	LEU	2.6
1	A	1898[A]	VAL	2.6
1	A	2068	ASN	2.6
2	B	254	ALA	2.6
1	A	2036	SER	2.6
1	A	2038	HIS	2.5
2	B	108	ILE	2.5
1	A	1980	LYS	2.5
1	A	1974	LEU	2.5
2	B	261	LEU	2.5
1	A	1855[A]	THR	2.5
2	B	41[A]	VAL	2.5
2	B	73	PRO	2.5
1	A	2058	LEU	2.5
2	B	317	LEU	2.4
2	B	69	ILE	2.4
2	B	235	GLY	2.4
2	B	40	HIS	2.4
2	B	146	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	149	GLU	2.4
2	B	174	HIS	2.4
2	B	220	TYR	2.4
2	B	270	TYR	2.4
2	B	92	ILE	2.4
2	B	181	ILE	2.4
1	A	1849	LYS	2.3
2	B	121	VAL	2.3
1	A	2026	LEU	2.3
2	B	268	LEU	2.3
1	A	1835	MET	2.3
1	A	1848	ILE	2.3
2	B	13	VAL	2.3
1	A	1833	GLY	2.2
1	A	1888	HIS	2.2
1	A	2024[A]	MET	2.2
2	B	201	LYS	2.2
1	A	2004	ALA	2.2
2	B	56	TYR	2.2
2	B	126	ILE	2.2
2	B	186	ARG	2.2
2	B	94	HIS	2.2
2	B	84	ARG	2.2
1	A	1893	ILE	2.1
1	A	1949	LEU	2.1
2	B	283	LEU	2.1
2	B	132	LYS	2.1
2	B	307	LYS	2.1
2	B	211[A]	GLN	2.1
2	B	247	ILE	2.1
2	B	76	GLY	2.1
2	B	91	ASN	2.1
2	B	241	LEU	2.1
2	B	100	GLN	2.1
1	A	2063	TYR	2.1
2	B	111	ASP	2.1
2	B	59	TRP	2.1
2	B	311	ASN	2.1
1	A	1850	LEU	2.1
1	A	1999	ILE	2.0
1	A	2059	ILE	2.0
1	A	1872	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	177[A]	ASN	2.0
1	A	1882	LEU	2.0
1	A	2022	ALA	2.0
2	B	112	ASP	2.0
2	B	217	SER	2.0
2	B	183	PHE	2.0
2	B	150	ASN	2.0
1	A	1977	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	R9G	A	2501	15/15	0.83	0.11	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.