



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:31 PM UTC

PDB ID : 8R02 / pdb_00008r02
Title : Crystal structure of the retromer complex VPS29/VPS35 with the ligand bis-1,3-phenyl guanylylhydrazone, 2a
Authors : Milani, M.; Fagnani, E.
Deposited on : 2023-10-30
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	185	 80% 16% ...
1	B	185	 82% 14% ...
2	C	306	 84% 14% .
2	D	306	 84% 10% . .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vacuolar protein sorting-associated protein 29.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1447	936	242	263	6			
1	B	182	Total	C	N	O	S	0	0	0
			1447	936	242	263	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	initiating methionine	UNP Q9UBQ0
A	-1	GLU	-	expression tag	UNP Q9UBQ0
A	0	HIS	-	expression tag	UNP Q9UBQ0
B	-2	MET	-	initiating methionine	UNP Q9UBQ0
B	-1	GLU	-	expression tag	UNP Q9UBQ0
B	0	HIS	-	expression tag	UNP Q9UBQ0


- Molecule 2 is a protein called Vacuolar protein sorting-associated protein 35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	305	Total	C	N	O	S	0	1	0
			2467	1565	426	464	12			
2	D	293	Total	C	N	O	S	0	0	0
			2371	1507	416	438	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	475	MET	-	initiating methionine	UNP Q96QK1
D	475	MET	-	initiating methionine	UNP Q96QK1

- Molecule 3 is Bis-1,3-phenyl guanyldiazon (CCD ID: XFZ) (formula: C₁₀H₁₄N₈) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	N	0	0
			18	10	8		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	14	Total	O	0	0
			14	14		
4	B	16	Total	O	0	0
			16	16		
4	C	17	Total	O	0	0
			17	17		
4	D	17	Total	O	0	0
			17	17		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vacuolar protein sorting-associated protein 29




- Molecule 1: Vacuolar protein sorting-associated protein 29



- Molecule 2: Vacuolar protein sorting-associated protein 35



- Molecule 2: Vacuolar protein sorting-associated protein 35



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.50Å 140.83Å 141.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.92 – 2.50 19.92 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.92-2.50) 99.7 (19.92-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.202 , 0.264 0.204 , 0.268	Depositor DCC
R_{free} test set	2025 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	75.7	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 77.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.008 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7814	wwPDB-VP
Average B, all atoms (Å ²)	117.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XFZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/1481	1.07	2/2008 (0.1%)
1	B	0.55	0/1481	1.06	3/2008 (0.1%)
2	C	0.54	0/2519	1.08	1/3395 (0.0%)
2	D	0.55	0/2417	1.08	0/3253
All	All	0.55	0/7898	1.07	6/10664 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	D	0	2
All	All	0	3

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	40	LEU	N-CA-C	-8.62	101.61	112.90
1	B	40	LEU	N-CA-C	-8.59	101.40	112.23
1	B	69	TYR	N-CA-CB	6.12	118.77	110.14
2	C	487	GLU	CB-CA-C	-6.03	100.43	110.68
1	B	12	PRO	N-CA-CB	-5.44	96.61	102.60
1	A	69	TYR	N-CA-CB	5.17	117.43	110.14

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	726	ARG	Sidechain
2	D	493	ARG	Sidechain
2	D	681	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1447	0	1459	26	0
1	B	1447	0	1459	16	0
2	C	2467	0	2436	23	0
2	D	2371	0	2367	19	0
3	D	18	0	0	1	0
4	A	14	0	0	0	0
4	B	16	0	0	1	0
4	C	17	0	0	0	0
4	D	17	0	0	1	0
All	All	7814	0	7721	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:ILE:HG21	1:A:159:THR:O	1.58	1.02
1:A:154:ASP:O	1:A:155:ILE:HG23	1.64	0.98
1:A:155:ILE:CG2	1:A:159:THR:O	2.34	0.73
1:A:155:ILE:HG21	1:A:159:THR:C	2.15	0.71
1:A:155:ILE:HG21	1:A:160:VAL:HA	1.75	0.67
2:C:579:LEU:HB3	2:C:580:PRO:HD3	1.80	0.64
2:D:579:LEU:HB3	2:D:580:PRO:HD3	1.80	0.63
1:A:154:ASP:O	1:A:155:ILE:CG2	2.46	0.61
2:C:501:GLU:OE2	2:C:501:GLU:HA	2.02	0.59
1:A:126:ASN:HB3	1:A:182:PRO:HG3	1.87	0.57
1:B:42:THR:HG21	2:D:490:LEU:HD11	1.86	0.57
2:D:663:LYS:N	2:D:664:PRO:HD2	2.21	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:LYS:NZ	4:B:201:HOH:O	2.40	0.55
1:B:156:GLN:HE21	1:B:156:GLN:HA	1.72	0.54
2:C:625:LEU:O	2:C:629:THR:HG23	2.07	0.54
1:B:40:LEU:HD21	1:B:58:ILE:CD1	2.38	0.54
2:C:663:LYS:N	2:C:664:PRO:HD2	2.23	0.54
2:D:726:ARG:NH2	4:D:901:HOH:O	2.30	0.53
2:C:475:MET:HB3	2:C:478:ASP:HB2	1.90	0.53
2:C:507:TYR:OH	2:C:559:LYS:HG2	2.09	0.52
2:C:754:LEU:N	2:C:755:PRO:HD2	2.25	0.52
2:D:650:ARG:HD2	2:D:677:PHE:CE1	2.45	0.52
1:A:157:ALA:O	1:A:159:THR:N	2.43	0.51
1:B:40:LEU:HD21	1:B:58:ILE:HD12	1.92	0.51
2:D:754:LEU:N	2:D:755:PRO:HD2	2.26	0.51
2:D:710:CYS:O	2:D:711:MET:HB2	2.12	0.49
1:B:154:ASP:HB3	1:B:161:VAL:HG13	1.94	0.49
2:C:725:ASN:HA	2:C:728:ILE:HD12	1.95	0.49
2:D:539:LEU:HD23	2:D:560:ILE:HD11	1.94	0.49
1:A:153:MET:HG2	1:A:162:THR:HG22	1.96	0.48
2:C:501:GLU:OE2	2:C:501:GLU:CA	2.61	0.48
1:A:11:ILE:HA	1:A:12:PRO:HA	1.71	0.48
1:A:4:LEU:HB3	1:A:34:ILE:HG12	1.96	0.47
1:A:157:ALA:O	1:A:159:THR:HG22	2.14	0.47
2:D:494:PHE:CZ	2:D:498:LEU:HD11	2.49	0.47
2:C:666:GLN:O	2:C:670:VAL:HG23	2.15	0.47
1:A:160:VAL:N	1:A:179:TYR:O	2.45	0.46
2:C:548:LYS:HA	2:C:548:LYS:HE2	1.97	0.46
1:A:43:LYS:HB3	1:A:43:LYS:HE3	1.53	0.46
2:D:581:LEU:C	2:D:581:LEU:HD23	2.41	0.46
2:C:632:ILE:HG21	2:C:672:THR:HG22	1.98	0.46
1:A:2:LEU:HA	1:A:153:MET:O	2.15	0.45
2:D:678:TRP:CE3	2:D:691:HIS:HA	2.51	0.45
2:C:650:ARG:HD2	2:C:677:PHE:CE1	2.51	0.45
1:A:155:ILE:CG2	1:A:160:VAL:HA	2.47	0.45
2:C:726:ARG:HE	2:C:726:ARG:HB2	1.60	0.44
1:B:43:LYS:O	1:B:46:TYR:N	2.50	0.44
2:C:678:TRP:CE3	2:C:691:HIS:HA	2.53	0.44
1:A:33:HIS:ND1	1:A:55:ASP:OD2	2.51	0.44
2:D:577:ALA:C	2:D:580:PRO:HD2	2.43	0.44
1:A:158:SER:O	1:A:180:LYS:HG3	2.17	0.43
1:B:33:HIS:ND1	1:B:55:ASP:OD2	2.51	0.43
2:D:632:ILE:HG21	2:D:672:THR:HG22	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:HE3	1:A:1:MET:HB2	1.84	0.43
1:B:16:ASN:HB2	2:D:493:ARG:HD2	2.00	0.43
1:B:11:ILE:HA	1:B:12:PRO:HA	1.70	0.43
2:C:495:ILE:HD11	2:C:532:LEU:HD23	2.01	0.43
2:C:597:GLU:O	2:C:598:ASN:CB	2.67	0.43
1:A:6:LEU:O	1:A:36:CYS:HA	2.18	0.42
1:A:40:LEU:HD21	1:A:58:ILE:HD12	2.00	0.42
1:B:40:LEU:H	1:B:40:LEU:HG	1.67	0.42
2:D:525:ILE:HG13	2:D:529:LEU:HD22	2.00	0.42
1:A:157:ALA:C	1:A:159:THR:N	2.78	0.42
1:B:6:LEU:O	1:B:36:CYS:HA	2.19	0.42
2:D:681:ARG:HE	2:D:681:ARG:HB2	1.39	0.42
1:A:12:PRO:HA	1:A:15:CYS:O	2.20	0.41
1:B:9:LEU:HB2	1:B:40:LEU:HA	2.01	0.41
1:B:43:LYS:O	1:B:44:GLU:C	2.60	0.41
2:C:577:ALA:C	2:C:580:PRO:HD2	2.45	0.41
2:D:666:GLN:O	2:D:670:VAL:HG23	2.21	0.41
1:A:40:LEU:HD21	1:A:58:ILE:CD1	2.51	0.41
2:C:748:GLN:OE1	2:C:748:GLN:HA	2.20	0.41
1:B:126:ASN:HB3	1:B:182:PRO:HG2	2.03	0.41
2:D:726:ARG:CZ	3:D:801:XFZ:NAP	2.84	0.40
1:A:121:ALA:HA	1:A:129:TYR:O	2.22	0.40
1:A:155:ILE:HG21	1:A:160:VAL:CA	2.47	0.40
2:C:710:CYS:O	2:C:711:MET:HB2	2.20	0.40
2:D:597:GLU:O	2:D:598:ASN:CB	2.69	0.40
2:C:525:ILE:HB	2:C:529:LEU:HD22	2.02	0.40
2:C:544:LYS:O	2:C:547:SER:HB3	2.22	0.40
2:C:647:GLU:O	2:C:651:THR:HG23	2.21	0.40
1:B:158:SER:O	1:B:180:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/185 (97%)	172 (96%)	8 (4%)	0	100	100
1	B	180/185 (97%)	170 (94%)	10 (6%)	0	100	100
2	C	304/306 (99%)	295 (97%)	9 (3%)	0	100	100
2	D	291/306 (95%)	280 (96%)	11 (4%)	0	100	100
All	All	955/982 (97%)	917 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/163 (98%)	150 (94%)	10 (6%)	16	34
1	B	160/163 (98%)	148 (92%)	12 (8%)	12	26
2	C	266/267 (100%)	252 (95%)	14 (5%)	20	42
2	D	254/267 (95%)	245 (96%)	9 (4%)	32	58
All	All	840/860 (98%)	795 (95%)	45 (5%)	20	41

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	30	LYS
1	A	40	LEU
1	A	43	LYS
1	A	50	LYS
1	A	72	GLN
1	A	118	LYS
1	A	155	ILE
1	A	160	VAL
1	A	164	VAL
1	B	1	MET
1	B	12	PRO
1	B	40	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	43	LYS
1	B	72	GLN
1	B	155	ILE
1	B	156	GLN
1	B	159	THR
1	B	160	VAL
1	B	161	VAL
1	B	170	ASP
1	B	173	LYS
2	C	498	LEU
2	C	501	GLU
2	C	508	LEU
2	C	514	ARG
2	C	542	ARG
2	C	547	SER
2	C	548	LYS
2	C	560	ILE
2	C	611	PHE
2	C	661	LEU
2	C	691	HIS
2	C	711	MET
2	C	726	ARG
2	C	740	ILE
2	D	491	VAL
2	D	525	ILE
2	D	560	ILE
2	D	611	PHE
2	D	651	THR
2	D	681	ARG
2	D	691	HIS
2	D	740	ILE
2	D	761	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	156	GLN
2	C	511	ASN
2	C	523	GLN
2	C	675	HIS
2	C	691	HIS
2	C	741	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	769	HIS
2	D	546	ASN
2	D	566	GLN
2	D	586	GLN
2	D	675	HIS
2	D	741	GLN
2	D	748	GLN
2	D	769	HIS
2	D	771	HIS
2	D	772	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	XFZ	D	801	-	18,18,18	2.44	7 (38%)	22,22,22	3.72	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	XFZ	D	801	-	-	3/12/12/12	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	801	XFZ	NAE-NAD	6.52	1.49	1.40
3	D	801	XFZ	CAF-NAE	3.47	1.35	1.27
3	D	801	XFZ	CAK-CAL	-3.38	1.39	1.47
3	D	801	XFZ	CAL-NAM	3.26	1.35	1.27
3	D	801	XFZ	CAG-CAF	-3.24	1.39	1.47
3	D	801	XFZ	NAM-NAN	2.91	1.44	1.40
3	D	801	XFZ	CAB-NAD	2.26	1.38	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	801	XFZ	CAB-NAD-NAE	13.47	127.98	110.90
3	D	801	XFZ	CAO-NAN-NAM	7.24	120.08	110.90
3	D	801	XFZ	CAF-NAE-NAD	7.22	125.89	113.14

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	801	XFZ	CAB-NAD-NAE-CAF
3	D	801	XFZ	CAJ-CAK-CAL-NAM
3	D	801	XFZ	CAR-CAK-CAL-NAM


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	801	XFZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/185 (98%)	-0.34	0 100 100	64, 104, 157, 188	1 (0%)
1	B	182/185 (98%)	-0.04	0 100 100	74, 123, 183, 221	5 (2%)
2	C	305/306 (99%)	-0.41	0 100 100	67, 113, 173, 201	4 (1%)
2	D	293/306 (95%)	-0.34	0 100 100	61, 108, 156, 181	8 (2%)
All	All	962/982 (97%)	-0.31	0 100 100	61, 112, 168, 221	18 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	XFZ	D	801	18/18	0.64	0.12	90,150,185,185	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.