



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 05:57 PM EDT

PDB ID : 5R1H / pdb_00005r1h
Title : PanDDA analysis group deposition – Auto-refined data of Aar2/RNaseH for ground state model 32, DMSO-free
Authors : Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.; Weiss, M.S.
Deposited on : 2020-02-12
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	
2	B	308	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	2002	1283	335	372	12	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2563	1642	419	482	20	0	9	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	170	SER	LEU	conflict	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ASP	deletion	UNP P32357


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	27	Total O 27 27	0	0
3	B	15	Total O 15 15	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-mRNA-splicing factor 8



• Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.00Å 82.99Å 92.17Å 90.00° 107.57° 90.00°	Depositor
Resolution (Å)	20.55 – 2.06 20.55 – 2.06	Depositor EDS
% Data completeness (in resolution range)	98.9 (20.55-2.06) 98.9 (20.55-2.06)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.268 , 0.320 0.287 , 0.334	Depositor DCC
R_{free} test set	1965 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	59.7	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 67.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4607	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	1/2049 (0.0%)	1.02	1/2775 (0.0%)
2	B	0.69	0/2630	0.86	1/3552 (0.0%)
All	All	0.79	1/4679 (0.0%)	0.93	2/6327 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1933	ILE	CG1-CD1	5.50	1.73	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	308	ILE	CA-CB-CG1	-5.74	100.65	110.40
1	A	1940	MET	CG-SD-CE	-5.16	89.55	100.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2002	0	2029	29	0
2	B	2563	0	2433	37	0
3	A	27	0	0	0	0
3	B	15	0	0	2	0
All	All	4607	0	4462	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.22	0.85
1:A:2062:GLU:O	1:A:2066:LYS:HG2	1.77	0.84
2:B:224:LEU:C	2:B:224:LEU:HD23	2.11	0.75
2:B:70:GLN:HB3	2:B:81:MET:HE1	1.69	0.73
2:B:224:LEU:HD23	2:B:224:LEU:O	1.88	0.73
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	1.72	0.70
2:B:1:MET:N	3:B:402:HOH:O	2.26	0.68
1:A:1979[B]:MET:HA	1:A:1979[B]:MET:HE2	1.79	0.64
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.26	0.61
2:B:300:ASN:O	2:B:303:HIS:NE2	2.35	0.60
1:A:1951:PHE:HB3	1:A:1954:ILE:HD12	1.86	0.56
1:A:2021:SER:O	1:A:2022:ALA:C	2.48	0.55
2:B:55:ARG:HD2	2:B:233:PHE:O	2.06	0.55
1:A:1910:LYS:HG2	1:A:1940:MET:SD	2.47	0.55
2:B:74:LYS:NZ	3:B:401:HOH:O	2.13	0.55
2:B:265:ASP:OD2	2:B:304:ASN:HB2	2.07	0.54
1:A:2047:GLN:O	1:A:2051:ILE:HG12	2.10	0.51
2:B:224:LEU:C	2:B:224:LEU:CD2	2.83	0.51
2:B:287:ARG:O	2:B:291:ILE:HD13	2.10	0.51
2:B:237:TYR:CE2	2:B:241:LEU:HD11	2.47	0.50
1:A:1878:CYS:HA	1:A:1892:LYS:O	2.13	0.49
2:B:77:LEU:O	2:B:77:LEU:HD23	2.13	0.49
1:A:1853:ASP:OD2	1:A:1855[B]:THR:HG23	2.12	0.48
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.48	0.48
2:B:190:ARG:HG3	2:B:203[B]:TYR:CE2	2.48	0.48
2:B:6:PHE:HZ	2:B:44[A]:ILE:HD11	1.79	0.47
2:B:291:ILE:HA	2:B:295:SER:HB2	1.96	0.47
2:B:193:HIS:ND1	2:B:196:GLU:OE1	2.33	0.47
1:A:1940:MET:HA	1:A:1940:MET:HE3	1.96	0.47
2:B:172:PRO:HA	2:B:175:SER:OG	2.15	0.47
2:B:70:GLN:HB3	2:B:81:MET:CE	2.42	0.47
2:B:72:ASP:OD1	2:B:72:ASP:C	2.55	0.46
2:B:201:LYS:HG3	2:B:252:SER:O	2.15	0.46
2:B:85:ASP:C	2:B:85:ASP:OD1	2.58	0.46
2:B:43:VAL:HG13	2:B:43:VAL:O	2.14	0.46
1:A:1942:ASP:HB2	1:A:1943:PRO:HD3	1.98	0.46
1:A:1974:LEU:O	1:A:1977:VAL:HG12	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1857:VAL:HG21	1:A:1913:THR:OG1	2.16	0.45
1:A:1940:MET:HE3	1:A:1940:MET:CA	2.47	0.45
1:A:2062:GLU:HB3	1:A:2066:LYS:HE3	1.99	0.45
2:B:64[B]:MET:HE3	2:B:136:GLN:OE1	2.17	0.45
1:A:1934:ILE:HA	1:A:1957:ARG:O	2.16	0.44
1:A:1914:ALA:HB1	1:A:1944:LEU:HA	1.98	0.44
1:A:1966:SER:C	1:A:1968:ALA:H	2.26	0.44
1:A:1977:VAL:CG1	1:A:1978:VAL:N	2.80	0.44
2:B:54[B]:MET:HE1	2:B:143:SER:HB3	1.99	0.43
1:A:1854:ASP:OD2	1:A:1940:MET:SD	2.77	0.43
2:B:220:TYR:CE1	2:B:246:MET:HE2	2.54	0.43
1:A:1910:LYS:O	1:A:1940:MET:HE1	2.18	0.43
1:A:2007:ARG:O	1:A:2008:LEU:C	2.60	0.43
2:B:315:GLU:C	2:B:317:LEU:N	2.74	0.42
2:B:2:ASN:CG	2:B:62:CYS:HB3	2.45	0.42
2:B:315:GLU:C	2:B:317:LEU:H	2.26	0.42
1:A:1875:ILE:HB	1:A:1979[B]:MET:HE1	2.02	0.42
2:B:244:HIS:HE1	2:B:283:LEU:O	2.02	0.42
2:B:56:TYR:CZ	2:B:77:LEU:HB2	2.55	0.41
1:A:1920:LEU:O	1:A:1924:LEU:HG	2.20	0.41
1:A:2056[B]:ARG:O	1:A:2060:LEU:HG	2.21	0.41
2:B:258:LYS:HD2	2:B:258:LYS:H	1.86	0.41
2:B:214:PHE:O	2:B:215:LYS:HB2	2.21	0.41
1:A:2026:LEU:HG	1:A:2055:MET:CE	2.51	0.41
1:A:2029:ASP:OD1	1:A:2029:ASP:C	2.64	0.41
2:B:3:THR:HG22	2:B:4:VAL:N	2.37	0.40
2:B:69:ILE:HD13	2:B:80:MET:HA	2.03	0.40
2:B:81:MET:HE3	2:B:81:MET:HB2	1.85	0.40
2:B:260:MET:HE2	2:B:260:MET:HB3	1.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/258 (96%)	236 (95%)	11 (4%)	1 (0%)	30	23
2	B	304/308 (99%)	280 (92%)	20 (7%)	4 (1%)	9	3
All	All	552/566 (98%)	516 (94%)	31 (6%)	5 (1%)	16	7

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	54[A]	MET
2	B	54[B]	MET
2	B	210	LEU
2	B	316	LEU
1	A	1964	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/233 (97%)	222 (98%)	4 (2%)	51	51
2	B	285/284 (100%)	276 (97%)	9 (3%)	34	29
All	All	511/517 (99%)	498 (98%)	13 (2%)	43	38

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1835	MET
1	A	1931[A]	LYS
1	A	1931[B]	LYS
1	A	1962	ARG
2	B	2	ASN
2	B	21	SER
2	B	77	LEU
2	B	142	SER
2	B	174	HIS
2	B	224	LEU
2	B	240	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	315	GLU
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1869	ASN
1	A	1907	GLN
1	A	2038	HIS
2	B	2	ASN
2	B	45	HIS
2	B	47	GLN
2	B	66	ASN
2	B	91	ASN
2	B	174	HIS
2	B	259	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	1.37	56 (23%) 2 1	34, 87, 139, 286	12 (5%)
2	B	300/308 (97%)	1.36	74 (24%) 2 1	35, 104, 160, 218	8 (2%)
All	All	537/566 (94%)	1.37	130 (24%) 2 1	34, 96, 154, 286	20 (3%)

All (130) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1878	CYS	13.4
1	A	1940	MET	6.5
2	B	176	LEU	6.0
1	A	1880	PHE	5.6
1	A	2040	TRP	5.5
1	A	1935	VAL	5.4
1	A	1882	LEU	4.9
2	B	146	THR	4.7
2	B	22	PHE	4.7
2	B	78	TYR	4.5
1	A	1996	LEU	4.5
2	B	267	ILE	4.2
1	A	2036	SER	4.2
2	B	257	PRO	4.2
2	B	54[A]	MET	4.2
2	B	283	LEU	4.0
2	B	234	PHE	4.0
2	B	284	LEU	3.7
1	A	1995	TRP	3.7
1	A	1840	TYR	3.6
2	B	313	TYR	3.6
2	B	24	VAL	3.5
1	A	1921	VAL	3.5
1	A	1979[A]	MET	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1986	MET	3.3
2	B	38	ILE	3.3
1	A	1881	THR	3.3
1	A	1875	ILE	3.3
2	B	264	LEU	3.2
2	B	174	HIS	3.2
1	A	2041	PRO	3.2
1	A	2038	HIS	3.2
1	A	1857	VAL	3.1
2	B	170	SER	3.1
2	B	256	VAL	3.0
2	B	302	LEU	3.0
2	B	233	PHE	3.0
2	B	101	MET	3.0
1	A	2026	LEU	3.0
2	B	53	SER	3.0
2	B	140	VAL	3.0
1	A	1841	ALA	3.0
1	A	1956	ILE	3.0
2	B	76	GLY	3.0
1	A	2058	LEU	2.9
1	A	1934	ILE	2.9
1	A	1853	ASP	2.9
1	A	2061	THR	2.9
2	B	120	PHE	2.9
2	B	305	THR	2.9
1	A	1858	TYR	2.9
2	B	104	SER	2.9
1	A	2059	ILE	2.8
2	B	129	ILE	2.8
1	A	2060	LEU	2.8
2	B	317	LEU	2.8
1	A	2017[A]	THR	2.8
2	B	289	TRP	2.8
1	A	1989	PHE	2.8
2	B	6	PHE	2.8
2	B	183	PHE	2.8
1	A	2027	LEU	2.8
2	B	316	LEU	2.7
2	B	52	SER	2.7
2	B	60	PHE	2.7
2	B	100	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	69	ILE	2.7
2	B	108	ILE	2.7
1	A	2015	LEU	2.6
2	B	31	HIS	2.6
2	B	304	ASN	2.6
1	A	1891	LEU	2.6
2	B	117	LEU	2.6
2	B	203[A]	TYR	2.6
2	B	232	MET	2.6
2	B	96	PHE	2.6
1	A	2069	VAL	2.6
2	B	92	ILE	2.6
1	A	1893	ILE	2.5
2	B	147	VAL	2.5
2	B	291	ILE	2.5
1	A	2037	TYR	2.5
1	A	2063	TYR	2.5
1	A	1936	THR	2.5
2	B	139	TYR	2.5
2	B	247	ILE	2.5
2	B	115	TYR	2.4
1	A	1866	PHE	2.4
1	A	2048	TRP	2.4
1	A	1835	MET	2.4
1	A	1877	GLY	2.4
2	B	226	PHE	2.4
2	B	103	VAL	2.4
1	A	1856	ASN	2.3
2	B	77	LEU	2.3
1	A	2021	SER	2.3
2	B	126	ILE	2.3
1	A	1899	TRP	2.3
2	B	243	TRP	2.3
2	B	20	TYR	2.3
2	B	89	PHE	2.3
2	B	191	PRO	2.3
1	A	1874	ALA	2.3
1	A	1833	GLY	2.3
2	B	275	LEU	2.3
1	A	1847	ASP	2.2
2	B	235	GLY	2.2
2	B	261	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1971	ILE	2.2
1	A	1895	HIS	2.2
2	B	193	HIS	2.2
2	B	279	TYR	2.2
1	A	1988	LEU	2.2
2	B	1	MET	2.2
2	B	114	TRP	2.2
1	A	1844	PHE	2.1
1	A	1954	ILE	2.1
2	B	270	TYR	2.1
1	A	1860	VAL	2.1
2	B	145	THR	2.1
2	B	32	GLY	2.1
2	B	137	PHE	2.1
1	A	1859	ARG	2.1
2	B	220	TYR	2.1
2	B	74	LYS	2.1
2	B	67	PHE	2.1
1	A	1992	TYR	2.0
2	B	130	VAL	2.0
2	B	239	SER	2.0
2	B	178	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.