



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:00 AM UTC

PDB ID : 5R1M / pdb_00005r1m
Title : PanDDA analysis group deposition – Auto-refined data of Aar2/RNaseH for ground state model 37, DMSO-free
Authors : Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.; Weiss, M.S.
Deposited on : 2020-02-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	19% 83% 8% 8%
2	B	308	21% 83% 13% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4650 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	1994	1278	334	371	11	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2580	1654	421	485	20	0	9	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	170	SER	LEU	conflict	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ASP	deletion	UNP P32357


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	53	Total	O	0	0
			53	53		
3	B	23	Total	O	0	0
			23	23		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.72Å 82.44Å 92.14Å 90.00° 107.55° 90.00°	Depositor
Resolution (Å)	21.88 – 1.90 21.88 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (21.88-1.90) 99.9 (21.88-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.249 , 0.299 0.260 , 0.308	Depositor DCC
R_{free} test set	2100 reflections (4.24%)	wwPDB-VP
Wilson B-factor (Å ²)	48.9	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4650	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/2041	0.84	0/2765
2	B	0.64	0/2651	0.74	0/3581
All	All	0.68	0/4692	0.79	0/6346

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1994	0	2020	18	0
2	B	2580	0	2450	30	0
3	A	53	0	0	2	0
3	B	23	0	0	3	0
All	All	4650	0	4470	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.28	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2062:GLU:O	1:A:2066:LYS:HG2	1.81	0.80
2:B:1:MET:N	3:B:402:HOH:O	2.14	0.79
2:B:258:LYS:HD2	2:B:258:LYS:H	1.61	0.65
1:A:2062:GLU:OE2	3:A:2101:HOH:O	2.14	0.65
1:A:1910:LYS:O	1:A:1940:MET:HE1	2.02	0.59
2:B:70:GLN:HB3	2:B:81:MET:HE2	1.83	0.59
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	1.85	0.58
2:B:96:PHE:HB2	2:B:102:MET:HE3	1.84	0.58
1:A:2066:LYS:HE2	3:A:2101:HOH:O	2.03	0.57
2:B:74:LYS:NZ	3:B:401:HOH:O	2.13	0.56
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.31	0.55
2:B:251:CYS:SG	2:B:292:CYS:SG	3.06	0.53
2:B:300:ASN:O	2:B:303:HIS:NE2	2.43	0.51
2:B:70:GLN:HB3	2:B:81:MET:CE	2.40	0.51
2:B:6:PHE:HZ	2:B:44[A]:ILE:HD11	1.76	0.51
2:B:54[A]:MET:HE1	3:B:422:HOH:O	2.10	0.51
1:A:2056[B]:ARG:O	1:A:2060:LEU:HG	2.10	0.50
1:A:1835:MET:HE2	1:A:1843:LEU:HD21	1.94	0.49
2:B:309:MET:HE1	2:B:316:LEU:HD23	1.93	0.49
2:B:6:PHE:CD1	2:B:32:GLY:HA2	2.49	0.47
2:B:96:PHE:CB	2:B:102:MET:HE3	2.45	0.47
2:B:287:ARG:O	2:B:291:ILE:HD13	2.15	0.47
2:B:190:ARG:HG3	2:B:203[B]:TYR:CE2	2.50	0.46
2:B:229:LEU:HD23	2:B:232:MET:HE2	1.98	0.46
2:B:2:ASN:CG	2:B:62:CYS:HB3	2.41	0.45
2:B:6:PHE:CZ	2:B:44[A]:ILE:HD11	2.51	0.45
1:A:2062:GLU:HB3	1:A:2066:LYS:HE2	1.99	0.45
1:A:1942:ASP:HB2	1:A:1943:PRO:HD3	2.00	0.44
2:B:118:THR:O	2:B:119:GLU:C	2.61	0.44
2:B:72:ASP:OD1	2:B:72:ASP:C	2.60	0.43
1:A:1941:LEU:O	1:A:1945:GLU:HB2	2.18	0.43
1:A:1951:PHE:HB3	1:A:1954:ILE:HD12	2.01	0.43
2:B:251:CYS:SG	2:B:292:CYS:HB3	2.59	0.43
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.54	0.43
2:B:210:LEU:O	2:B:215:LYS:N	2.50	0.43
1:A:1878:CYS:SG	1:A:1969:MET:HE3	2.59	0.42
1:A:1859:ARG:HH12	1:A:1979[A]:MET:CE	2.31	0.42
2:B:68:TYR:CE2	2:B:81:MET:HE3	2.55	0.42
2:B:260:MET:HE2	2:B:260:MET:HB3	1.96	0.41
2:B:298:GLN:HA	2:B:298:GLN:OE1	2.20	0.41
1:A:2063:TYR:CD1	1:A:2063:TYR:C	2.98	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:190:ARG:HG3	2:B:203[B]:TYR:CZ	2.56	0.41
2:B:188:ALA:HA	2:B:204:TYR:CD1	2.56	0.40
1:A:1859:ARG:NH1	1:A:1979[A]:MET:SD	2.91	0.40
2:B:203[A]:TYR:CZ	2:B:207:THR:HG21	2.57	0.40
1:A:1835:MET:CE	1:A:1843:LEU:HD21	2.52	0.40
2:B:141:ASP:OD1	2:B:144:MET:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/258 (96%)	239 (97%)	8 (3%)	0	100	100
2	B	306/308 (99%)	289 (94%)	15 (5%)	2 (1%)	18	10
All	All	553/566 (98%)	528 (96%)	23 (4%)	2 (0%)	43	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	54[A]	MET
2	B	54[B]	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/233 (97%)	220 (98%)	5 (2%)	45	42
2	B	287/284 (101%)	283 (99%)	4 (1%)	59	59
All	All	512/517 (99%)	503 (98%)	9 (2%)	55	50

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1835	MET
1	A	1931[A]	LYS
1	A	1931[B]	LYS
1	A	1962	ARG
1	A	2058	LEU
2	B	2	ASN
2	B	174	HIS
2	B	275	LEU
2	B	315	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1863	HIS
1	A	1907	GLN
1	A	2038	HIS
2	B	2	ASN
2	B	66	ASN
2	B	290	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	1.24	48 (20%)	3 2	23, 64, 105, 143	12 (5%)
2	B	300/308 (97%)	1.34	64 (21%)	2 2	27, 76, 127, 165	9 (3%)
All	All	537/566 (94%)	1.30	112 (20%)	2 2	23, 71, 120, 165	21 (3%)

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1940	MET	8.0
1	A	1878	CYS	7.4
2	B	74	LYS	5.8
2	B	288	VAL	5.4
2	B	77	LEU	5.4
2	B	22	PHE	5.3
2	B	54[A]	MET	5.0
2	B	275	LEU	4.7
1	A	2060	LEU	4.5
1	A	1979[A]	MET	4.5
1	A	2069	VAL	4.2
1	A	1834	ALA	3.9
2	B	199	LEU	3.8
1	A	1964	PRO	3.8
2	B	52	SER	3.8
2	B	53	SER	3.6
2	B	203[A]	TYR	3.5
2	B	170	SER	3.5
1	A	1948	MET	3.5
2	B	279	TYR	3.4
2	B	96	PHE	3.4
1	A	2051	ILE	3.3
1	A	1835	MET	3.3
2	B	1	MET	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2048	TRP	3.2
1	A	1988	LEU	3.1
2	B	316	LEU	3.1
1	A	1875	ILE	3.1
2	B	108	ILE	3.0
2	B	20	TYR	3.0
2	B	313	TYR	3.0
1	A	2040	TRP	2.9
2	B	14	THR	2.9
1	A	2066	LYS	2.9
2	B	130	VAL	2.9
2	B	183	PHE	2.9
2	B	101	MET	2.9
2	B	139	TYR	2.8
1	A	1866	PHE	2.8
2	B	99	ARG	2.8
2	B	71	TYR	2.8
2	B	291	ILE	2.8
2	B	146	THR	2.7
1	A	1967	ALA	2.7
2	B	100	GLN	2.7
2	B	191	PRO	2.6
2	B	276	PRO	2.6
1	A	1961	LEU	2.6
2	B	69	ILE	2.6
1	A	1833	GLY	2.6
1	A	1995	TRP	2.6
1	A	1853	ASP	2.6
1	A	2034	ILE	2.6
1	A	1954	ILE	2.5
1	A	1956	ILE	2.5
2	B	24	VAL	2.5
1	A	2061	THR	2.5
1	A	1909	ALA	2.5
1	A	1855[A]	THR	2.5
2	B	172	PRO	2.5
2	B	192	GLY	2.4
1	A	1963	LEU	2.4
2	B	129	ILE	2.4
1	A	2035	LYS	2.4
2	B	103	VAL	2.4
1	A	2063	TYR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	175	SER	2.4
2	B	198	PHE	2.4
1	A	1981	ALA	2.4
2	B	87	ALA	2.4
1	A	2043	PHE	2.3
1	A	1838	SER	2.3
2	B	317	LEU	2.3
1	A	1840	TYR	2.3
2	B	145	THR	2.3
2	B	267	ILE	2.3
2	B	283	LEU	2.3
2	B	281	ASP	2.3
1	A	1865	THR	2.2
1	A	2058	LEU	2.2
2	B	73	PRO	2.2
2	B	294	TYR	2.2
1	A	1893	ILE	2.2
2	B	33	ILE	2.2
2	B	179	THR	2.2
2	B	137	PHE	2.2
1	A	1969	MET	2.2
1	A	2065	ARG	2.2
2	B	116	ASN	2.2
1	A	2010	LEU	2.2
2	B	152	LEU	2.2
1	A	1986	MET	2.1
2	B	193	HIS	2.1
1	A	1941	LEU	2.1
2	B	97	LYS	2.1
2	B	258	LYS	2.1
2	B	188	ALA	2.1
1	A	2059	ILE	2.1
1	A	2033	THR	2.1
1	A	1901	GLY	2.1
2	B	106	PRO	2.1
1	A	2037	TYR	2.1
2	B	25	LYS	2.1
2	B	174	HIS	2.1
2	B	5	PRO	2.1
2	B	122[A]	GLN	2.1
2	B	136	GLN	2.1
1	A	1994	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	220	TYR	2.0
1	A	1936	THR	2.0
2	B	238	GLY	2.0
2	B	113	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.