



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 09:54 PM UTC

PDB ID : 7R1E / pdb_00007r1e
Title : Mosquitocidal Cry11Ba determined at pH 10.4 from naturally-occurring nanocrystals by Serial femtosecond crystallography
Authors : Colletier, J.-P.; Sawaya, M.R.; Schibrowsky, N.A.; Cascio, D.; Rodriguez, J.A.
Deposited on : 2022-02-02
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	724	% 77% 10% 14%
1	B	724	% 76% 9% 14%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10031 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pesticidal crystal protein Cry11Ba.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	626	Total	C	N	O	S	0	0	0
			4968	3196	823	937	12			
1	B	621	Total	C	N	O	S	0	0	0
			4932	3168	819	934	11			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	67	Total 67	O 67	0	0
3	B	52	Total 52	O 52	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	167.50Å 157.99Å 57.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.72 – 2.65 35.72 – 2.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (35.72-2.65) 99.9 (35.72-2.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.65Å)	Xtriage
Refinement program	PHENIX 1.20_4459	Depositor
R, R_{free}	0.236 , 0.247 0.237 , 0.247	Depositor DCC
R_{free} test set	4526 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	41.0	Xtriage
Anisotropy	0.308	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10031	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 68.36 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.4242e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.07	0/5089	0.22	0/6918
1	B	0.07	0/5051	0.22	0/6866
All	All	0.07	0/10140	0.22	0/13784

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4968	0	4888	36	0
1	B	4932	0	4857	37	0
2	A	12	0	16	0	0
3	A	67	0	0	0	0
3	B	52	0	0	1	0
All	All	10031	0	9761	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:307:HIS:HA	1:A:462:THR:HG23	1.79	0.64
1:B:82:LEU:HD11	1:B:224:MET:HE2	1.80	0.62
1:A:296:ASP:HB2	1:A:361:VAL:HG23	1.83	0.61
1:A:242:GLU:OE1	1:A:641:GLN:NE2	2.33	0.60
1:B:304:GLU:HB2	1:B:399:GLY:HA2	1.82	0.59
1:A:304:GLU:HB2	1:A:399:GLY:HA2	1.86	0.57
1:A:288:LYS:HB2	1:A:310:LEU:HB3	1.88	0.55
1:B:73:LYS:HD2	1:B:76:ILE:HD11	1.90	0.53
1:A:572:GLY:HA2	1:B:567:ARG:HH22	1.76	0.50
1:B:165:LYS:NZ	3:B:808:HOH:O	2.44	0.50
1:A:71:MET:HE1	1:A:155:GLN:HG3	1.92	0.50
1:B:150:ILE:O	1:B:154:THR:OG1	2.29	0.49
1:A:287:VAL:HG21	1:A:433:VAL:HB	1.96	0.48
1:B:95:GLU:HG2	1:B:99:LYS:HE2	1.96	0.48
1:B:319:ILE:HG23	1:B:455:LYS:HA	1.95	0.48
1:B:321:GLY:HA3	1:B:457:GLU:HB3	1.95	0.48
1:A:324:GLY:HA2	1:A:460:ALA:HB3	1.96	0.47
1:A:280:PRO:HB2	1:A:498:LEU:HB3	1.97	0.46
1:B:71:MET:HE1	1:B:155:GLN:HG3	1.97	0.46
1:A:23:ARG:NH2	1:A:81:LEU:O	2.49	0.46
1:A:319:ILE:HG23	1:A:455:LYS:HA	1.98	0.46
1:A:523:PHE:CD1	1:A:530:ASN:HB3	2.51	0.46
1:A:597:THR:HG23	1:A:600:ASP:H	1.80	0.46
1:B:515:GLN:O	1:B:629:LYS:NZ	2.47	0.45
1:B:597:THR:HG23	1:B:600:ASP:H	1.80	0.45
1:B:544:THR:O	1:B:597:THR:OG1	2.35	0.45
1:A:559:ARG:HG3	1:A:582:VAL:O	2.16	0.45
1:B:90:ARG:HH12	1:B:139:GLN:HG3	1.81	0.44
1:A:121:THR:HG23	1:A:124:ARG:HH22	1.82	0.44
1:A:62:ILE:HG23	1:A:71:MET:HE3	1.98	0.44
1:B:558:ALA:HB3	1:B:623:GLN:HA	1.99	0.44
1:B:93:ILE:O	1:B:97:GLU:HG2	2.18	0.43
1:A:393:SER:OG	1:A:394:PRO:HD3	2.19	0.43
1:B:386:ASP:HB3	1:B:403:LYS:HB3	2.00	0.43
1:A:93:ILE:O	1:A:97:GLU:HG2	2.19	0.43
1:B:393:SER:OG	1:B:394:PRO:HD3	2.18	0.43
1:B:452:THR:O	1:B:454:ASN:N	2.51	0.43
1:A:502:THR:HG22	1:A:533:LYS:HB2	2.01	0.43
1:B:210:LEU:HD13	1:B:217:GLU:HG2	2.00	0.43
1:A:295:SER:HB3	1:A:364:ALA:HB3	2.00	0.43
1:A:384:THR:HG22	1:A:424:LYS:HB2	2.00	0.43
1:A:121:THR:HG23	1:A:124:ARG:NH2	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:449:LEU:HD22	1:A:456:LEU:HD12	2.01	0.42
1:A:631:ILE:HG22	1:A:633:VAL:HG13	2.00	0.42
1:B:414:VAL:HG12	1:B:416:SER:H	1.83	0.42
1:A:558:ALA:HB3	1:A:623:GLN:HA	2.01	0.42
1:B:451:LEU:HD21	1:B:509:LEU:HB2	2.02	0.42
1:B:551:ILE:HB	1:B:631:ILE:HB	2.01	0.42
1:B:279:ALA:O	1:B:431:ARG:NH1	2.47	0.42
1:B:297:THR:HA	1:B:363:ALA:HB2	2.02	0.42
1:B:631:ILE:HG22	1:B:633:VAL:HG13	2.01	0.42
1:A:118:GLU:OE1	1:A:121:THR:N	2.53	0.42
1:A:478:PHE:O	1:A:482:ASN:N	2.46	0.42
1:B:128:ASP:O	1:B:132:GLN:HB2	2.20	0.42
1:B:424:LYS:NZ	1:B:426:ASP:OD1	2.51	0.42
1:B:288:LYS:HB2	1:B:310:LEU:HB3	2.00	0.42
1:B:588:TRP:CZ3	1:B:628:SER:HB2	2.55	0.41
1:B:53:GLN:O	1:B:57:LYS:HG3	2.20	0.41
1:B:425:TYR:HB2	1:B:428:TYR:HB2	2.01	0.41
1:A:382:PHE:HB2	1:A:430:ILE:HB	2.03	0.41
1:A:414:VAL:HG12	1:A:416:SER:H	1.85	0.41
1:A:572:GLY:HA2	1:B:567:ARG:NH2	2.35	0.41
1:B:245:LYS:HB2	1:B:492:ALA:HB3	2.02	0.41
1:A:70:THR:HG22	1:A:71:MET:HE2	2.02	0.41
1:A:441:SER:HB3	1:A:444:TYR:CE2	2.56	0.40
1:B:213:LYS:HE3	1:B:217:GLU:OE1	2.21	0.40
1:A:321:GLY:HA3	1:A:457:GLU:HB3	2.02	0.40
1:B:230:PHE:CE2	1:B:513:PRO:HB2	2.56	0.40
1:A:230:PHE:CE2	1:A:513:PRO:HB2	2.57	0.40
1:A:254:ASN:HD22	1:A:423:LEU:HD22	1.87	0.40
1:B:555:LYS:HE3	1:B:588:TRP:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	620/724 (86%)	604 (97%)	15 (2%)	1 (0%)	43	60
1	B	615/724 (85%)	597 (97%)	18 (3%)	0	100	100
All	All	1235/1448 (85%)	1201 (97%)	33 (3%)	1 (0%)	48	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	453	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	543/635 (86%)	538 (99%)	5 (1%)	70	82
1	B	540/635 (85%)	536 (99%)	4 (1%)	76	86
All	All	1083/1270 (85%)	1074 (99%)	9 (1%)	73	85

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	LEU
1	A	166	ASP
1	A	389	TRP
1	A	417	ILE
1	A	447	ASP
1	B	417	ILE
1	B	424	LYS
1	B	447	ASP
1	B	526	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	123	ASN
1	A	220	ASN
1	A	609	ASN
1	B	113	ASN
1	B	160	HIS
1	B	286	ASN
1	B	290	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	802	-	5,5,5	0.96	0	5,5,5	1.08	0
2	GOL	A	801	-	5,5,5	0.95	0	5,5,5	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	802	-	-	0/4/4/4	-
2	GOL	A	801	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	626/724 (86%)	0.00	9 (1%) 73 69	26, 42, 77, 132	0
1	B	621/724 (85%)	-0.07	5 (0%) 82 79	24, 42, 67, 113	0
All	All	1247/1448 (86%)	-0.03	14 (1%) 78 75	24, 42, 73, 132	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	361	VAL	5.1
1	B	17	PHE	4.1
1	A	655	THR	3.3
1	A	329	SER	3.2
1	A	17	PHE	3.1
1	B	655	THR	2.9
1	B	394	PRO	2.7
1	A	12	ASN	2.6
1	A	14	MET	2.5
1	A	657	PHE	2.3
1	B	361	VAL	2.3
1	B	297	THR	2.2
1	A	595	SER	2.2
1	A	351	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	801	6/6	0.84	0.12	55,64,67,71	0
2	GOL	A	802	6/6	0.86	0.12	44,54,56,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.