



Full wwPDB NMR Structure Validation Report ⓘ

Mar 19, 2026 – 07:54 PM UTC

PDB ID : 1R2N / pdb_00001r2n
Title : NMR structure of the all-trans retinal in dark-adapted Bacteriorhodopsin
Authors : Patzelt, H.; Simon, B.; terLaak, A.; Kessler, B.; Kuhne, R.; Schmieder, P.; Oesterhaelt, D.; Oschkinat, H.
Deposited on : 2003-09-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	249	

2 Ensemble composition and analysis

This entry contains 12 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:119, A:124-A:139, A:144-A:182, A:188-A:213, A:218-A:232 (215)	0.06	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 10, 11, 12
2	2, 8, 9


3 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3706 atoms, of which 1890 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Bacteriorhodopsin.

Mol	Chain	Residues	Atoms						Trace
1	A	232	Total	C	H	N	O	S	0
			3646	1202	1854	275	306	9	

- Molecule 2 is RETINAL (CCD ID: RET) (formula: $C_{20}H_{28}O$).



Mol	Chain	Residues	Atoms		
2	A	1	Total	C	H
			48	20	28

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		
3	A	4	Total	H	O
			12	8	4

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Bacteriorhodopsin




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Bacteriorhodopsin



4.2.2 Score per residue for model 2

- Molecule 1: Bacteriorhodopsin






4.2.3 Score per residue for model 3

- Molecule 1: Bacteriorhodopsin


Chain A: 71% 12% 7% 7%



4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: Bacteriorhodopsin


Chain A: 73% 11% 7% 7%



4.2.5 Score per residue for model 5


- Molecule 1: Bacteriorhodopsin

Chain A: 73% 11% 7% 7%




4.2.6 Score per residue for model 6

- Molecule 1: Bacteriorhodopsin




4.2.7 Score per residue for model 7

- Molecule 1: Bacteriorhodopsin




4.2.8 Score per residue for model 8

- Molecule 1: Bacteriorhodopsin




4.2.9 Score per residue for model 9

- Molecule 1: Bacteriorhodopsin




4.2.10 Score per residue for model 10

- Molecule 1: Bacteriorhodopsin




4.2.11 Score per residue for model 11

- Molecule 1: Bacteriorhodopsin



4.2.12 Score per residue for model 12

- Molecule 1: Bacteriorhodopsin



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 12 were deposited, based on the following criterion: *rms distance between observed and calculated chemical shifts*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	structure solution	4.1
Amber	refinement	4.1

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RET

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.75±0.00	1±0/1725 (0.1± 0.0%)	1.56±0.01	15±1/2355 (0.7± 0.1%)
All	All	0.75	12/20700 (0.1%)	1.56	185/28260 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.6±0.6
All	All	0	43

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	96	ASP	CG-OD2	5.12	1.35	1.25	5	12

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	85	ASP	CA-CB-CG	7.17	119.77	112.60	6	9
1	A	49	VAL	N-CA-CB	6.51	114.88	110.52	7	3
1	A	230	PHE	CB-CA-C	6.37	123.09	110.42	5	12
1	A	165	PRO	CA-C-N	6.30	129.05	120.54	2	12
1	A	165	PRO	C-N-CA	6.30	129.05	120.54	2	12
1	A	230	PHE	CA-C-N	6.18	129.78	121.01	4	12
1	A	230	PHE	C-N-CA	6.18	129.78	121.01	4	12
1	A	128	THR	CA-C-N	6.01	128.25	120.44	12	12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	128	THR	C-N-CA	6.01	128.25	120.44	12	12
1	A	177	VAL	CA-C-N	5.89	128.43	120.65	4	10
1	A	177	VAL	C-N-CA	5.89	128.43	120.65	4	10
1	A	193	SER	CA-C-N	5.67	128.44	120.28	3	5
1	A	193	SER	C-N-CA	5.67	128.44	120.28	3	5
1	A	148	ILE	CA-C-N	5.48	127.94	120.54	11	1
1	A	148	ILE	C-N-CA	5.48	127.94	120.54	11	1
1	A	49	VAL	CA-CB-CG2	5.46	119.68	110.40	1	1
1	A	211	LEU	CA-C-N	5.43	127.82	120.65	12	3
1	A	211	LEU	C-N-CA	5.43	127.82	120.65	12	3
1	A	31	GLY	CA-C-N	5.19	127.19	120.44	8	12
1	A	31	GLY	C-N-CA	5.19	127.19	120.44	8	12
1	A	49	VAL	CA-C-N	5.15	124.59	119.24	4	1
1	A	49	VAL	C-N-CA	5.15	124.59	119.24	4	1
1	A	110	ALA	CA-C-N	5.14	127.44	120.65	12	12
1	A	110	ALA	C-N-CA	5.14	127.44	120.65	12	12

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	7	ARG	Sidechain	12
1	A	43	TYR	Sidechain	12
1	A	96	ASP	Sidechain	12
1	A	83	TYR	Sidechain	3
1	A	115	ASP	Sidechain	3
1	A	82	ARG	Sidechain	1

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1679	1737	1733	7±0
2	A	20	28	27	0±0
All	All	20436	21276	21120	89

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:166:GLU:CD	1:A:227:ARG:HD3	0.63	2.18	3	12
2:A:999:RET:H8	2:A:999:RET:H161	0.57	1.75	1	1
1:A:49:VAL:HG23	1:A:50:PRO:HD3	0.51	1.82	1	3
1:A:5:THR:HG21	1:A:77:PRO:HB3	0.49	1.84	2	12
1:A:129:LYS:H	1:A:129:LYS:CD	0.45	2.25	2	12
1:A:20:MET:SD	1:A:53:ALA:HB1	0.45	2.52	12	1
1:A:129:LYS:H	1:A:129:LYS:HD3	0.43	1.72	2	12
1:A:12:TRP:CD2	1:A:206:LEU:HD23	0.43	2.48	4	12
1:A:10:TRP:CG	1:A:11:ILE:N	0.42	2.88	2	12
1:A:15:LEU:HB3	1:A:209:MET:HE2	0.42	1.92	2	12

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/249 (86%)	184±1 (86±0%)	24±1 (12±0%)	5±0 (2±0%)	7	45
All	All	2556/2988 (86%)	2202 (86%)	294 (12%)	60 (2%)	7	45

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	2	ALA	12
1	A	7	ARG	12
1	A	29	VAL	12
1	A	34	VAL	12
1	A	229	ILE	12

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	173/195 (89%)	161±1 (93±1%)	12±1 (7±1%)	15 64
All	All	2076/2340 (89%)	1928 (93%)	148 (7%)	15 64

All 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	38	ASP	12
1	A	66	LEU	12
1	A	68	MET	12
1	A	97	LEU	12
1	A	99	LEU	12
1	A	111	LEU	12
1	A	129	LYS	12
1	A	151	VAL	12
1	A	166	GLU	12
1	A	206	LEU	12
1	A	229	ILE	12
1	A	89	THR	5
1	A	85	ASP	4
1	A	204	GLU	2
1	A	194	GLU	1
1	A	94	LEU	1
1	A	93	LEU	1
1	A	119	ILE	1
1	A	145	MET	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	RET	A	999	1	20,20,21	3.11±0.05	8±0 (40±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	RET	A	999	1	27,27,28	1.72±0.07	8±2 (27±5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RET	A	999	1	-	0±0,13,30,31	0±0,1,1,1

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	999	RET	C14-C13	11.92	1.41	1.33	2	12
2	A	999	RET	C5-C6	4.85	1.42	1.34	2	12
2	A	999	RET	C8-C7	2.80	1.41	1.33	4	12
2	A	999	RET	C12-C13	2.65	1.40	1.46	12	12
2	A	999	RET	C15-C14	2.53	1.40	1.49	12	12
2	A	999	RET	C11-C12	2.40	1.40	1.34	7	12
2	A	999	RET	C8-C9	2.28	1.41	1.46	12	12
2	A	999	RET	C10-C9	2.27	1.41	1.35	4	12

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	A	999	RET	C18-C5-C6	3.61	120.55	124.48	1	12
2	A	999	RET	C19-C9-C8	3.50	123.43	118.09	12	12
2	A	999	RET	C1-C6-C5	3.24	118.21	122.64	11	12
2	A	999	RET	C10-C11-C12	3.10	114.21	123.20	12	12
2	A	999	RET	C1-C6-C7	3.05	123.91	115.65	2	12
2	A	999	RET	C8-C9-C10	2.96	114.35	119.01	12	12
2	A	999	RET	C2-C1-C6	2.76	114.45	110.44	10	2
2	A	999	RET	C2-C3-C4	2.68	105.39	111.28	10	3
2	A	999	RET	C17-C1-C6	2.64	114.38	110.24	8	2
2	A	999	RET	C16-C1-C6	2.30	113.85	110.24	2	4
2	A	999	RET	C20-C13-C12	2.15	121.36	118.09	8	1
2	A	999	RET	C16-C1-C2	2.13	100.78	108.95	6	5
2	A	999	RET	C17-C1-C2	2.01	101.23	108.95	8	1


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided