



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:30 AM UTC

PDB ID : 1R2T / pdb_00001r2t
Title : CRYSTAL STRUCTURE OF RABBIT MUSCLE TRIOSEPHOSPHATE ISOMERASE
Authors : Aparicio, R.; Ferreira, S.T.; Polikarpov, I.
Deposited on : 2003-09-29
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	3% 85% 12% ..
1	B	248	2% 83% 13% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Triosephosphate isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	8	4	0
			1874	1183	324	360	7			
1	B	247	Total	C	N	O	S	14	3	0
			1881	1192	324	358	7			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	166	Total	O	0	0
			166	166		
2	B	169	Total	O	0	0
			169	169		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Triosephosphate isomerase



• Molecule 1: Triosephosphate isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.16Å 72.03Å 93.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.54 – 2.25 15.54 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.5 (15.54-2.25) 97.1 (15.54-2.25)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.25Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.182 , 0.220 0.178 , 0.219	Depositor DCC
R_{free} test set	1032 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4090	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	3/1928 (0.2%)	1.03	8/2609 (0.3%)
1	B	0.82	3/1929 (0.2%)	1.04	6/2611 (0.2%)
All	All	0.86	6/3857 (0.2%)	1.04	14/5220 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	179	GLN	CG-CD	-23.27	0.93	1.52
1	B	214	GLY	C-N	-15.94	1.10	1.33
1	B	32	LYS	CB-CG	-13.58	1.11	1.52
1	A	145	GLU	CD-OE1	-12.76	1.01	1.25
1	A	145	GLU	CD-OE2	8.64	1.41	1.25
1	B	141	LYS	CD-CE	-7.43	1.30	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	32	LYS	CA-CB-CG	15.69	145.47	114.10
1	A	145	GLU	CB-CG-CD	12.11	133.19	112.60
1	A	179	GLN	CB-CG-CD	10.89	131.12	112.60
1	B	32	LYS	CB-CG-CD	10.67	135.85	111.30
1	A	179	GLN	OE1-CD-NE2	-9.59	113.01	122.60
1	A	145	GLU	CG-CD-OE2	-8.87	97.99	118.40
1	B	216	THR	OG1-CB-CG2	-6.31	96.68	109.30
1	A	187	LYS	CG-CD-CE	-5.76	98.06	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	246	ALA	N-CA-C	5.63	118.15	111.33
1	A	179	GLN	CG-CD-NE2	5.60	124.79	116.40
1	A	216	THR	OG1-CB-CG2	-5.51	98.29	109.30
1	B	58	LYS	CA-CB-CG	5.50	125.09	114.10
1	B	58	LYS	CB-CG-CD	5.43	123.80	111.30
1	B	141	LYS	CD-CE-NZ	-5.03	95.81	111.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	213	THR	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1874	0	1886	13	0
1	B	1881	0	1897	23	0
2	A	166	0	0	1	1
2	B	169	0	0	5	1
All	All	4090	0	3783	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:218:LYS:NZ	1:B:248:GLN:O	1.88	1.06
1:A:52:ARG:HD3	1:A:52:ARG:O	1.74	0.87
1:B:52:ARG:O	1:B:52:ARG:HD3	1.79	0.82
1:B:2:PRO:HB3	2:B:417:HOH:O	1.84	0.78
1:B:213:THR:OG1	1:B:216:THR:HG22	1.86	0.75
1:B:143:VAL:HG13	1:B:188:LEU:HD21	1.68	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:LYS:NZ	2:A:371:HOH:O	2.22	0.72
1:A:143:VAL:HG13	1:A:188:LEU:HD21	1.76	0.67
1:A:52:ARG:HD3	1:A:52:ARG:C	2.24	0.61
1:B:52:ARG:HD3	1:B:52:ARG:C	2.24	0.61
1:B:2:PRO:HA	1:B:3:SER:O	2.01	0.60
1:B:223:GLN:O	2:B:396:HOH:O	2.17	0.59
1:A:134:ARG:HD3	1:A:168:TRP:CD2	2.39	0.57
1:A:167:VAL:HA	1:A:170:ILE:HD12	1.86	0.57
1:B:40:VAL:HG22	1:B:60:ALA:HB3	1.87	0.56
1:B:127:ILE:HD12	1:B:162[B]:LEU:HD12	1.86	0.56
1:B:134:ARG:HD3	1:B:168:TRP:CD2	2.41	0.56
1:B:12:TRP:CZ2	1:B:24[B]:LEU:HD21	2.43	0.54
1:A:80:PRO:HB3	1:A:121:LEU:HD22	1.91	0.53
1:B:80:PRO:HB3	1:B:121:LEU:HD22	1.91	0.52
1:B:84:LYS:HG2	1:B:121:LEU:HD13	1.93	0.51
1:B:175:THR:HG23	2:B:315:HOH:O	2.11	0.50
1:A:40:VAL:HG22	1:A:60:ALA:HB3	1.96	0.48
1:B:19:LYS:NZ	2:B:383:HOH:O	2.47	0.48
1:A:139:THR:O	1:A:143:VAL:HB	2.17	0.45
1:B:24[B]:LEU:HD23	1:B:25:ILE:N	2.32	0.43
1:A:5:LYS:HE2	1:A:38:GLU:HB2	1.99	0.43
1:B:139:THR:O	1:B:143:VAL:HB	2.19	0.42
1:B:5:LYS:HE2	1:B:38:GLU:HB2	2.01	0.42
1:A:9:GLY:HA2	1:A:40:VAL:O	2.20	0.42
1:A:33:VAL:HA	1:A:34:PRO:HD3	1.96	0.41
1:B:179:GLN:NE2	1:B:179:GLN:H	2.17	0.41
1:B:19:LYS:HE2	2:B:385:HOH:O	2.19	0.41
1:A:125:ALA:HB1	1:A:150:ILE:HD13	2.02	0.41
1:B:127:ILE:HD12	1:B:162[B]:LEU:CD1	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:323:HOH:O	2:B:341:HOH:O[2_564]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/248 (100%)	242 (98%)	6 (2%)	0	100	100
1	B	248/248 (100%)	237 (96%)	9 (4%)	2 (1%)	16	14
All	All	496/496 (100%)	479 (97%)	15 (3%)	2 (0%)	30	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	172	THR
1	B	3	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/197 (102%)	183 (92%)	17 (8%)	10	8
1	B	200/197 (102%)	184 (92%)	16 (8%)	11	9
All	All	400/394 (102%)	367 (92%)	33 (8%)	11	8

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	LYS
1	A	33	VAL
1	A	52	ARG
1	A	69	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	84	LYS
1	A	108	LEU
1	A	121	LEU
1	A	127	ILE
1	A	143	VAL
1	A	155[A]	LYS
1	A	155[B]	LYS
1	A	179	GLN
1	A	183[A]	GLU
1	A	183[B]	GLU
1	A	211	SER
1	A	218	LYS
1	A	248	GLN
1	B	23	GLU
1	B	33	VAL
1	B	52	ARG
1	B	69	VAL
1	B	84	LYS
1	B	108	LEU
1	B	121	LEU
1	B	127	ILE
1	B	143	VAL
1	B	155	LYS
1	B	172	THR
1	B	183	GLU
1	B	194	SER
1	B	202	GLN
1	B	211	SER
1	B	218	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	179	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	214:GLY	C	215:ALA	N	1.10

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	246/248 (99%)	0.22	7 (2%) 55 55	9, 17, 28, 38	7 (2%)
1	B	246/248 (99%)	-0.06	6 (2%) 59 61	8, 17, 30, 39	6 (2%)
All	All	492/496 (99%)	0.08	13 (2%) 57 58	8, 17, 30, 39	13 (2%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	201	ALA	4.0
1	B	175	THR	3.2
1	B	172	THR	3.0
1	B	248	GLN	3.0
1	A	75	THR	2.6
1	B	2	PRO	2.6
1	A	200	VAL	2.4
1	A	173	GLY	2.3
1	B	36	ASP	2.1
1	A	171	GLY	2.1
1	B	75	THR	2.1
1	A	248	GLN	2.1
1	A	202	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.