



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:30 AM UTC

PDB ID : 2R2O / pdb_00002r2o
Title : Crystal structure of the effector domain of human Plexin B1
Authors : Tong, Y.; Tempel, W.; Shen, L.; Arrowsmith, C.H.; Edwards, A.M.; Sundstrom, M.; Weigelt, J.; Bochkarev, A.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2007-08-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	
1	B	138	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UNX	A	203	-	-	-	X
2	UNX	A	205	-	-	-	X
2	UNX	A	207	-	-	-	X
2	UNX	A	208	-	-	-	X
2	UNX	A	210	-	-	-	X
2	UNX	A	212	-	-	-	X
2	UNX	B	201	-	-	-	X
2	UNX	B	202	-	-	-	X
2	UNX	B	204	-	-	-	X
2	UNX	B	209	-	-	-	X
2	UNX	B	211	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1573 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plexin-B1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	Se	0	1	0
			754	482	128	141	2	1			
1	B	103	Total	C	N	O	S	Se	0	0	0
			759	488	127	141	2	1			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1725	MSE	-	expression tag	UNP O43157
A	1726	HIS	-	expression tag	UNP O43157
A	1727	HIS	-	expression tag	UNP O43157
A	1728	HIS	-	expression tag	UNP O43157
A	1729	HIS	-	expression tag	UNP O43157
A	1730	HIS	-	expression tag	UNP O43157
A	1731	HIS	-	expression tag	UNP O43157
A	1732	SER	-	expression tag	UNP O43157
A	1733	SER	-	expression tag	UNP O43157
A	1734	GLY	-	expression tag	UNP O43157
A	1735	ARG	-	expression tag	UNP O43157
A	1736	GLU	-	expression tag	UNP O43157
A	1737	ASN	-	expression tag	UNP O43157
A	1738	LEU	-	expression tag	UNP O43157
A	1739	TYR	-	expression tag	UNP O43157
A	1740	PHE	-	expression tag	UNP O43157
A	1741	GLN	-	expression tag	UNP O43157
A	1742	GLY	-	expression tag	UNP O43157
B	1725	MSE	-	expression tag	UNP O43157
B	1726	HIS	-	expression tag	UNP O43157
B	1727	HIS	-	expression tag	UNP O43157
B	1728	HIS	-	expression tag	UNP O43157
B	1729	HIS	-	expression tag	UNP O43157
B	1730	HIS	-	expression tag	UNP O43157
B	1731	HIS	-	expression tag	UNP O43157

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1732	SER	-	expression tag	UNP O43157
B	1733	SER	-	expression tag	UNP O43157
B	1734	GLY	-	expression tag	UNP O43157
B	1735	ARG	-	expression tag	UNP O43157
B	1736	GLU	-	expression tag	UNP O43157
B	1737	ASN	-	expression tag	UNP O43157
B	1738	LEU	-	expression tag	UNP O43157
B	1739	TYR	-	expression tag	UNP O43157
B	1740	PHE	-	expression tag	UNP O43157
B	1741	GLN	-	expression tag	UNP O43157
B	1742	GLY	-	expression tag	UNP O43157

- Molecule 2 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	7	Total X 7 7	0	0
2	B	5	Total X 5 5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	24	Total O 24 24	0	0
3	B	24	Total O 24 24	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Plexin-B1



• Molecule 1: Plexin-B1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.98Å 56.57Å 99.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 20.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.2 (20.00-2.00) 95.0 (20.00-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.01Å)	Xtriage
Refinement program	REFMAC refmac_5.3.0037	Depositor
R, R_{free}	0.222 , 0.269 (Not available) , 0.276	Depositor DCC
R_{free} test set	842 reflections (4.20%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 73.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1573	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	2/769 (0.3%)	0.85	0/1050
1	B	1.10	0/772	0.90	0/1055
All	All	1.03	2/1541 (0.1%)	0.87	0/2105

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1767	VAL	CA-C	6.14	1.57	1.52
1	A	1767	VAL	CA-CB	6.06	1.59	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	754	0	759	1	0
1	B	759	0	755	4	0
2	A	7	0	0	0	0
2	B	5	0	0	0	0
3	A	24	0	0	0	0
3	B	24	0	0	0	0
All	All	1573	0	1514	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1754:LEU:HD23	1:B:1766:GLY:HA2	1.86	0.56
1:A:1789:TYR:CZ	1:A:1798:PRO:HD2	2.49	0.48
1:B:1754:LEU:HD23	1:B:1766:GLY:CA	2.44	0.47
1:B:1756:ALA:HB2	1:B:1850:VAL:HG22	1.97	0.47
1:B:1756:ALA:CB	1:B:1850:VAL:HG22	2.47	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/138 (70%)	96 (99%)	1 (1%)	0	100	100
1	B	99/138 (72%)	99 (100%)	0	0	100	100
All	All	196/276 (71%)	195 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/117 (69%)	80 (99%)	1 (1%)	63	70
1	B	78/117 (67%)	77 (99%)	1 (1%)	61	68
All	All	159/234 (68%)	157 (99%)	2 (1%)	61	68

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1767	VAL
1	B	1852	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 12 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/138 (71%)	0.41	6 (6%) 27 26	24, 39, 51, 56	1 (1%)
1	B	102/138 (73%)	0.19	5 (4%) 35 34	31, 36, 50, 60	0
All	All	201/276 (72%)	0.30	11 (5%) 30 29	24, 38, 51, 60	1 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1746	TYR	5.2
1	A	1746	TYR	4.4
1	A	1789	TYR	4.3
1	B	1764	ALA	3.3
1	B	1756	ALA	2.9
1	B	1852	CYS	2.8
1	A	1764	ALA	2.6
1	A	1852	CYS	2.6
1	A	1826	VAL	2.3
1	B	1759	PRO	2.3
1	A	1815	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	UNX	B	209	1/1	-0.20	1.50	2,2,2,2	1
2	UNX	B	202	1/1	-0.09	1.54	2,2,2,2	1
2	UNX	A	207	1/1	0.05	1.62	2,2,2,2	1
2	UNX	A	203	1/1	0.11	1.16	13,13,13,13	1
2	UNX	B	204	1/1	0.14	1.43	2,2,2,2	1
2	UNX	A	210	1/1	0.18	1.56	2,2,2,2	1
2	UNX	A	205	1/1	0.23	1.71	2,2,2,2	1
2	UNX	A	208	1/1	0.26	1.69	2,2,2,2	1
2	UNX	A	212	1/1	0.27	1.83	2,2,2,2	1
2	UNX	B	201	1/1	0.32	1.14	2,2,2,2	1
2	UNX	B	211	1/1	0.55	1.46	2,2,2,2	1
2	UNX	A	206	1/1	0.85	2.01	8,8,8,8	1

6.5 Other polymers [i](#)

There are no such residues in this entry.