



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 03:13 PM UTC

PDB ID : 3R2G / pdb_00003r2g
Title : Crystal structure of Inosine 5' monophosphate dehydrogenase from Legionella pneumophila
Authors : Agarwal, R.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRC)
Deposited on : 2011-03-14
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	5% 79% 8% 11%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inosine 5'-monophosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2405	1500	425	461	19			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q5ZRN7
A	2	VAL	-	expression tag	UNP Q5ZRN7
A	340	ALA	-	expression tag	UNP Q5ZRN7
A	341	GLU	-	expression tag	UNP Q5ZRN7
A	342	ASN	-	expression tag	UNP Q5ZRN7
A	343	LEU	-	expression tag	UNP Q5ZRN7
A	344	TYR	-	expression tag	UNP Q5ZRN7
A	345	PHE	-	expression tag	UNP Q5ZRN7
A	346	GLN	-	expression tag	UNP Q5ZRN7
A	347	SER	-	expression tag	UNP Q5ZRN7
A	348	HIS	-	expression tag	UNP Q5ZRN7
A	349	HIS	-	expression tag	UNP Q5ZRN7
A	350	HIS	-	expression tag	UNP Q5ZRN7
A	351	HIS	-	expression tag	UNP Q5ZRN7
A	352	HIS	-	expression tag	UNP Q5ZRN7
A	353	HIS	-	expression tag	UNP Q5ZRN7
A	354	TRP	-	expression tag	UNP Q5ZRN7
A	355	SER	-	expression tag	UNP Q5ZRN7
A	356	HIS	-	expression tag	UNP Q5ZRN7
A	357	PRO	-	expression tag	UNP Q5ZRN7
A	358	GLN	-	expression tag	UNP Q5ZRN7
A	359	PHE	-	expression tag	UNP Q5ZRN7
A	360	GLU	-	expression tag	UNP Q5ZRN7
A	361	LYS	-	expression tag	UNP Q5ZRN7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	217	Total 217	O 217	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Inosine 5'-monophosphate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	115.13Å 115.13Å 63.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.70 – 1.94 40.70 – 1.94	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.70-1.94) 97.6 (40.70-1.94)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.179 , 0.207 0.180 , 0.175	Depositor DCC
R_{free} test set	1516 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	27.2	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2622	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	2/2441 (0.1%)	0.91	1/3289 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	149	VAL	C-O	-6.30	1.17	1.23
1	A	146	ALA	C-O	-5.82	1.17	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	171	GLY	N-CA-C	-5.30	100.62	113.18

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2405	0	2399	28	0
2	A	217	0	0	3	0
All	All	2622	0	2399	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:MET:HE1	1:A:83:GLU:HG2	1.52	0.92
1:A:80:ASN:HD22	1:A:104:ARG:HH11	1.41	0.68
1:A:45:VAL:CG1	1:A:227:ILE:HD12	2.25	0.67
1:A:35:ARG:HG3	1:A:35:ARG:NH1	2.09	0.66
1:A:35:ARG:HH11	1:A:35:ARG:CG	2.14	0.60
1:A:35:ARG:HG3	1:A:35:ARG:HH11	1.66	0.60
1:A:150:ALA:HB2	1:A:170:ILE:HG12	1.84	0.59
1:A:45:VAL:HG13	1:A:227:ILE:HD12	1.87	0.56
1:A:35:ARG:NH1	1:A:35:ARG:CG	2.71	0.53
1:A:175:VAL:HG22	1:A:178:THR:H	1.73	0.53
1:A:245:ASP:O	1:A:246:ASP:CB	2.57	0.52
1:A:31:THR:HG22	2:A:448:HOH:O	2.10	0.51
1:A:143:CYS:C	1:A:144:ILE:HD13	2.37	0.49
1:A:171:GLY:HA3	1:A:175:VAL:HG13	1.95	0.49
1:A:35:ARG:NH2	2:A:415:HOH:O	2.46	0.48
1:A:172:GLY:HA2	1:A:179:ARG:CD	2.44	0.47
1:A:173:GLY:O	1:A:174:SER:HB2	2.17	0.45
1:A:45:VAL:CG1	1:A:227:ILE:CD1	2.94	0.45
1:A:32:SER:HB3	1:A:44:PRO:HB3	1.99	0.44
1:A:188:MET:O	1:A:192:ILE:HG12	2.17	0.44
1:A:227:ILE:H	1:A:227:ILE:HD13	1.81	0.44
1:A:51:ASP:HA	1:A:72:HIS:CD2	2.52	0.44
1:A:81:ILE:CD1	1:A:108:LEU:HD23	2.49	0.42
1:A:126:LYS:HE3	1:A:130:LYS:HE3	2.02	0.42
1:A:144:ILE:HD13	1:A:144:ILE:N	2.36	0.40
1:A:150:ALA:HB2	1:A:170:ILE:CG1	2.51	0.40
1:A:181:LYS:HE3	2:A:442:HOH:O	2.21	0.40
1:A:8:ILE:HG21	1:A:322:ILE:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/361 (89%)	303 (94%)	14 (4%)	4 (1%)	10 3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	174	SER
1	A	246	ASP
1	A	171	GLY
1	A	172	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	251/289 (87%)	243 (97%)	8 (3%)	34 20

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ARG
1	A	46	ILE
1	A	144	ILE
1	A	166	ILE
1	A	175	VAL
1	A	177	SER
1	A	227	ILE
1	A	245	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	80	ASN
1	A	262	GLN
1	A	268	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/361 (89%)	0.19	18 (5%) 30 33	16, 25, 50, 64	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	173	GLY	11.8
1	A	175	VAL	7.4
1	A	174	SER	6.4
1	A	171	GLY	6.4
1	A	172	GLY	6.1
1	A	274	THR	4.1
1	A	27	VAL	3.5
1	A	176	CYS	3.5
1	A	25	ARG	3.3
1	A	328	ARG	3.3
1	A	327	GLY	3.1
1	A	273	LYS	2.7
1	A	170	ILE	2.6
1	A	28	VAL	2.5
1	A	242	PHE	2.3
1	A	309	ASP	2.2
1	A	246	ASP	2.1
1	A	275	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.