



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:33 AM UTC

PDB ID : 8R2W / pdb_00008r2w
Title : Crystal structure of hydroxyquinol-1,2-dioxygenase from *Trametes versicolor* (TvHDX1)
Authors : Zahn, M.; Kuatsjah, E.; Salvachua, D.
Deposited on : 2023-11-07
Resolution : 1.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	8% 78% 5% 16%
1	B	347	3% 77% 5% 16%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 5235 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydroxyquinol 1,2-dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	1	0
			2278	1442	391	438	7			
1	B	290	Total	C	N	O	S	0	1	0
			2281	1444	394	436	7			

- Molecule 2 is PHOSPHATIDYLETHANOLAMINE (CCD ID: PTY) (formula: $C_{40}H_{80}NO_8P$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			50	40	1	8	1		
2	B	1	Total	C	N	O	P	0	0
			50	40	1	8	1		

- Molecule 3 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		
3	B	1	Total	Fe	0	0
			1	1		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	260	Total	O	0	0
			260	260		
5	B	306	Total	O	0	0
			306	306		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hydroxyquinol 1,2-dioxygenase



• Molecule 1: Hydroxyquinol 1,2-dioxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	177.36Å 54.12Å 69.49Å 90.00° 108.39° 90.00°	Depositor
Resolution (Å)	84.16 – 1.46 84.16 – 1.46	Depositor EDS
% Data completeness (in resolution range)	80.8 (84.16-1.46) 80.8 (84.16-1.46)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 1.46Å)	Xtriage
Refinement program	REFMAC 5.8.0411	Depositor
R, R_{free}	0.172 , 0.203 0.178 , 0.210	Depositor DCC
R_{free} test set	4338 reflections (3.98%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5235	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FE, TRS, PTY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2339	1.01	4/3183 (0.1%)
1	B	0.70	2/2342 (0.1%)	0.98	3/3186 (0.1%)
All	All	0.68	2/4681 (0.0%)	1.00	7/6369 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
All	All	0	6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	25	GLU	CD-OE1	-5.81	1.14	1.25
1	B	142	GLU	CD-OE2	5.57	1.35	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	106	ASP	CA-CB-CG	7.63	120.23	112.60
1	A	235	ASN	CA-CB-CG	6.52	119.12	112.60
1	B	235	ASN	CA-CB-CG	6.19	118.79	112.60
1	A	127	THR	CA-CB-OG1	-5.73	101.00	109.60
1	A	240	MET	CG-SD-CE	-5.57	88.64	100.90
1	A	302	ARG	CG-CD-NE	-5.55	99.79	112.00
1	B	97	ASP	CA-CB-CG	5.21	117.81	112.60

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	185	ARG	Sidechain
1	A	281	ARG	Sidechain
1	A	302	ARG	Sidechain
1	A	62	ARG	Sidechain
1	B	247	ARG	Sidechain
1	B	62	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2278	0	2214	9	0
1	B	2281	0	2223	15	0
2	A	50	0	79	6	0
2	B	50	0	79	5	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	B	8	0	12	1	0
5	A	260	0	0	3	0
5	B	306	0	0	5	0
All	All	5235	0	4607	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:401:TRS:N	5:B:502:HOH:O	2.00	0.94
1:B:52:ASN:OD1	5:B:501:HOH:O	1.92	0.86
2:A:401:PTY:H211	2:B:402:PTY:H291	1.61	0.82
1:B:241:VAL:HG21	1:B:304:ILE:HD13	1.68	0.74
1:B:247:ARG:HH12	1:B:307:VAL:HG21	1.53	0.73
1:A:283:GLU:O	1:A:287:ARG:HG2	1.88	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:LYS:HE3	5:A:683:HOH:O	1.89	0.72
1:A:279:GLU:OE1	5:A:501:HOH:O	2.10	0.70
2:A:401:PTY:C21	2:B:402:PTY:H291	2.23	0.68
1:B:287:ARG:NH1	1:B:292:PRO:O	2.29	0.66
2:B:402:PTY:H361	2:B:402:PTY:H251	1.77	0.65
1:B:253:TRP:CZ2	1:B:302[B]:ARG:HG2	2.36	0.59
1:B:253:TRP:CH2	1:B:302[B]:ARG:HG2	2.41	0.55
1:A:113:VAL:HA	5:A:702:HOH:O	2.06	0.55
2:A:401:PTY:H211	2:B:402:PTY:C29	2.38	0.51
2:A:401:PTY:H242	1:B:53:LEU:HD12	1.93	0.51
1:A:104:LEU:HD11	1:B:62:ARG:HD3	1.94	0.49
1:A:108:ILE:HD11	1:B:62:ARG:HD2	1.95	0.49
1:B:247:ARG:HH12	1:B:307:VAL:CG2	2.25	0.48
1:B:94:LEU:HG	1:B:220:VAL:HG11	1.96	0.47
2:A:401:PTY:H201	2:A:401:PTY:H171	1.77	0.45
1:B:52:ASN:HB3	5:B:501:HOH:O	2.16	0.45
1:B:158:ASP:HB3	5:B:626:HOH:O	2.16	0.45
1:A:94:LEU:HG	1:A:220:VAL:HG11	1.99	0.43
1:B:222:ASN:HB3	2:B:402:PTY:O12	2.19	0.42
1:A:111:PRO:O	1:A:113:VAL:N	2.52	0.42
1:B:309:GLU:HG2	5:B:559:HOH:O	2.19	0.42
1:A:95:LEU:HG	2:A:401:PTY:H392	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/347 (83%)	283 (98%)	6 (2%)	0	100	100
1	B	289/347 (83%)	286 (99%)	3 (1%)	0	100	100
All	All	578/694 (83%)	569 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/293 (85%)	246 (99%)	3 (1%)	63	33
1	B	249/293 (85%)	247 (99%)	2 (1%)	73	50
All	All	498/586 (85%)	493 (99%)	5 (1%)	68	42

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	235	ASN
1	A	283	GLU
1	B	121	VAL
1	B	235	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	HIS
1	B	175	HIS
1	B	222	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PTY	B	402	-	49,49,49	0.66	0	52,54,54	0.72	2 (3%)
2	PTY	A	401	-	49,49,49	0.72	0	52,54,54	0.71	3 (5%)
4	TRS	B	401	-	7,7,7	0.46	0	9,9,9	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTY	B	402	-	-	26/53/53/53	-
2	PTY	A	401	-	-	17/53/53/53	-
4	TRS	B	401	-	-	2/9/9/9	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	402	PTY	O7-C6-C5	-2.55	99.19	108.34
2	B	402	PTY	O11-C3-C2	2.34	117.73	109.17
2	A	401	PTY	O7-C6-C1	2.14	116.04	108.34
2	A	401	PTY	O11-C3-C2	2.11	116.89	109.17
2	A	401	PTY	O7-C6-C5	-2.07	100.91	108.34

There are no chirality outliers.

All (45) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	402	PTY	N1-C2-C3-O11
2	B	402	PTY	C3-O11-P1-O12
2	B	402	PTY	C3-O11-P1-O14
2	A	401	PTY	C17-C18-C19-C20
2	A	401	PTY	C6-C5-O14-P1
2	A	401	PTY	C15-C16-C17-C18
2	B	402	PTY	C24-C25-C26-C27
2	A	401	PTY	C22-C23-C24-C25
2	B	402	PTY	C11-C12-C13-C14
2	B	402	PTY	C37-C38-C39-C40
2	B	402	PTY	C17-C18-C19-C20
2	B	402	PTY	C14-C15-C16-C17
2	B	402	PTY	C21-C22-C23-C24
2	A	401	PTY	C26-C27-C28-C29
2	A	401	PTY	C20-C21-C22-C23
2	A	401	PTY	C36-C37-C38-C39
2	B	402	PTY	C23-C24-C25-C26
2	B	402	PTY	C25-C26-C27-C28
2	B	402	PTY	C39-C40-C41-C42
2	B	402	PTY	C22-C23-C24-C25
2	A	401	PTY	C12-C13-C14-C15
2	B	402	PTY	C41-C42-C43-C44
2	B	402	PTY	C6-C5-O14-P1
2	B	402	PTY	C38-C39-C40-C41
2	A	401	PTY	C39-C40-C41-C42
2	B	402	PTY	C12-C13-C14-C15
2	A	401	PTY	C14-C15-C16-C17
2	B	402	PTY	C16-C17-C18-C19
2	A	401	PTY	C37-C38-C39-C40
2	B	402	PTY	C33-C34-C35-C36
2	B	402	PTY	C36-C37-C38-C39
2	A	401	PTY	C5-O14-P1-O13
2	B	402	PTY	C3-O11-P1-O13
2	B	402	PTY	C32-C33-C34-C35
2	A	401	PTY	C21-C22-C23-C24
4	B	401	TRS	C1-C-C2-O2
2	A	401	PTY	C16-C17-C18-C19
2	B	402	PTY	C35-C36-C37-C38
2	A	401	PTY	C35-C36-C37-C38
2	A	401	PTY	C25-C26-C27-C28
2	B	402	PTY	O4-C1-C6-O7
2	A	401	PTY	C38-C39-C40-C41

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	402	PTY	C18-C19-C20-C21
4	B	401	TRS	N-C-C2-O2
2	B	402	PTY	C15-C16-C17-C18


There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	402	PTY	5	0
2	A	401	PTY	6	0
4	B	401	TRS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/347 (83%)	0.42	27 (9%) 14 14	15, 25, 44, 74	1 (0%)
1	B	290/347 (83%)	0.26	12 (4%) 41 44	14, 23, 38, 73	1 (0%)
All	All	580/694 (83%)	0.34	39 (6%) 24 23	14, 24, 42, 74	2 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	VAL	8.1
1	A	114	HIS	4.8
1	B	20	LYS	4.7
1	A	20	LYS	3.4
1	A	294	GLY	3.4
1	A	160	THR	3.2
1	A	157	THR	3.1
1	B	23	TYR	3.0
1	A	115	GLY	2.8
1	A	88	LEU	2.8
1	B	247	ARG	2.8
1	A	307	VAL	2.7
1	B	21	LEU	2.7
1	B	293	LYS	2.6
1	A	306	LEU	2.5
1	A	184	VAL	2.5
1	B	262	GLU	2.4
1	A	198	LYS	2.4
1	A	293	LYS	2.4
1	B	295	GLY	2.4
1	A	112	PRO	2.4
1	A	309	GLU	2.4
1	A	199	ASP	2.3
1	B	121	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	308	PRO	2.2
1	B	225	LEU	2.2
1	A	163	PRO	2.2
1	A	244	PRO	2.1
1	A	162	VAL	2.1
1	A	305	ILE	2.1
1	A	246	PHE	2.1
1	A	121	VAL	2.1
1	A	295	GLY	2.1
1	B	26	SER	2.1
1	A	125	PHE	2.0
1	B	113	VAL	2.0
1	A	23	TYR	2.0
1	B	108	ILE	2.0
1	A	158	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PTY	B	402	50/50	0.83	0.18	26,41,57,62	0
2	PTY	A	401	50/50	0.84	0.17	28,36,47,56	0
4	TRS	B	401	8/8	0.87	0.12	22,26,31,37	0
3	FE	B	403	1/1	1.00	0.01	15,15,15,15	0
3	FE	A	402	1/1	1.00	0.03	18,18,18,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around TRS B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.