



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:20 PM UTC

PDB ID : 1R3F / pdb_00001r3f
Title : Crystal Structure of tRNA Pseudouridine Synthase TruB and Its RNA Complex: RNA-protein Recognition Through a Combination of Rigid Docking and Induced Fit
Authors : Pan, H.; Agarwalla, S.; Moustakas, D.T.; Finer-Moore, J.; Stroud, R.M.
Deposited on : 2003-10-01
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	14% 74% 14% • 11%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA pseudouridine synthase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	278	2158	1354	377	418	9	0	0	0

- Molecule 1: tRNA pseudouridine synthase B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.03Å 71.01Å 48.67Å 90.00° 117.87° 90.00°	Depositor
Resolution (Å)	50.00 – 1.85 43.02 – 1.85	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.85) 96.0 (43.02-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.76 (at 1.84Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.228 , 0.257 0.250 , 0.261	Depositor DCC
R_{free} test set	2306 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 26.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.004 for -h-l,k,h 0.004 for l,k,-h-l 0.023 for h,-k,-h-l 0.023 for -h-l,-k,l 0.027 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2158	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2193	0.90	3/2971 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	242	MET	N-CA-C	-6.23	102.01	110.36
1	A	54	MET	N-CA-C	5.34	117.92	109.96
1	A	77	ARG	CD-NE-CZ	-5.06	117.32	124.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2158	0	2163	29	0
All	All	2158	0	2163	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:ASP:HB3	1:A:181:ARG:NH1	2.00	0.76
1:A:271:ARG:NH1	1:A:302:ARG:HG2	2.13	0.63
1:A:79:ILE:HB	1:A:199:TYR:HB3	1.82	0.61
1:A:43:HIS:HD2	1:A:45:GLY:H	1.49	0.61
1:A:70:LEU:HD23	1:A:70:LEU:O	2.01	0.61
1:A:299:ASP:HB2	1:A:300:GLU:OE1	2.00	0.60
1:A:95:ILE:HD12	1:A:95:ILE:N	2.17	0.60
1:A:97:GLU:HG2	1:A:99:ARG:HD2	1.85	0.58
1:A:43:HIS:CD2	1:A:45:GLY:H	2.22	0.57
1:A:86:THR:HG23	1:A:89:SER:H	1.71	0.55
1:A:84:GLN:O	1:A:196:HIS:HD2	1.91	0.53
1:A:48:ASP:OD2	1:A:181:ARG:NH2	2.43	0.52
1:A:70:LEU:HD23	1:A:70:LEU:C	2.36	0.51
1:A:86:THR:HG21	1:A:90:ASP:O	2.12	0.49
1:A:97:GLU:CG	1:A:99:ARG:HD2	2.42	0.49
1:A:95:ILE:HD12	1:A:95:ILE:H	1.77	0.48
1:A:252:PRO:HG2	1:A:281:VAL:HG12	1.96	0.48
1:A:104:SER:OG	1:A:107:GLN:HG3	2.14	0.47
1:A:88:THR:O	1:A:89:SER:HB2	2.14	0.47
1:A:86:THR:HG23	1:A:89:SER:N	2.30	0.47
1:A:296:GLU:HB2	1:A:306:ARG:HG3	1.96	0.47
1:A:87:ASP:O	1:A:88:THR:HG23	2.15	0.47
1:A:278:GLU:HG2	1:A:279:GLY:N	2.32	0.45
1:A:262:VAL:HG13	1:A:263:TYR:N	2.33	0.44
1:A:69:LEU:HG	1:A:247:PRO:HA	1.99	0.44
1:A:300:GLU:OE1	1:A:300:GLU:N	2.52	0.43
1:A:159:LEU:HB3	1:A:171:GLU:HB2	2.01	0.42
1:A:121:GLN:O	1:A:122:ILE:HG23	2.21	0.41
1:A:158:LEU:HD23	1:A:158:LEU:C	2.45	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	274/314 (87%)	269 (98%)	3 (1%)	2 (1%)	18 8

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	88	THR
1	A	276	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	234/265 (88%)	229 (98%)	5 (2%)	47 33

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	ASP
1	A	94	GLN
1	A	219	LEU
1	A	273	SER
1	A	300	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	HIS
1	A	84	GLN
1	A	166	ASN
1	A	196	HIS
1	A	229	GLN
1	A	255	ASN
1	A	266	ASN
1	A	268	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	278/314 (88%)	0.95	43 (15%) 5 5	20, 31, 51, 59	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	91	ALA	6.9
1	A	96	VAL	5.7
1	A	122	ILE	5.6
1	A	123	PRO	5.6
1	A	92	ASP	5.4
1	A	275	ALA	5.0
1	A	272	THR	4.6
1	A	88	THR	4.5
1	A	87	ASP	4.2
1	A	277	LEU	3.9
1	A	274	GLY	3.9
1	A	93	GLY	3.8
1	A	273	SER	3.7
1	A	153	ILE	3.5
1	A	95	ILE	3.4
1	A	302	ARG	3.2
1	A	310	VAL	3.2
1	A	70	LEU	3.2
1	A	86	THR	3.1
1	A	94	GLN	3.1
1	A	312	TYR	3.1
1	A	8	GLY	2.9
1	A	90	ASP	2.8
1	A	301	GLY	2.8
1	A	40	ARG	2.8
1	A	276	PRO	2.7
1	A	268	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	297	ILE	2.6
1	A	306	ARG	2.5
1	A	271	ARG	2.5
1	A	278	GLU	2.5
1	A	270	VAL	2.5
1	A	299	ASP	2.3
1	A	89	SER	2.3
1	A	263	TYR	2.3
1	A	98	GLU	2.3
1	A	298	ASP	2.3
1	A	300	GLU	2.2
1	A	269	PRO	2.2
1	A	266	ASN	2.2
1	A	313	PRO	2.1
1	A	267	GLY	2.1
1	A	314	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.