



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 27, 2026 – 07:56 PM UTC

PDB ID : 7R3V / pdb_00007r3v
Title : Crystal structure of bovine Cytochrome bc1 in complex with inhibitor CK-2-67.
Authors : Pinthong, N.; Ampornpanai, K.; O'Neill, P.M.; Hasnain, S.S.; Antonyuk, S.
Deposited on : 2022-02-07
Resolution : 3.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	2% 76% 22% .
2	B	418	2% 71% 26% ..
3	C	379	4% 72% 26% ..
4	D	239	3% 76% 23% .
5	E	196	21% 66% 30% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	99	
7	G	74	
8	H	65	
9	I	46	
10	J	59	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
16	PEE	C	406	X	-	-	-
16	PEE	E	204	X	-	-	-

2 Entry composition [i](#)

There are 23 unique types of molecules in this entry. The entry contains 16382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome b-c1 complex subunit 1, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	445	Total	C	N	O	S	0	0	0
			3449	2157	608	664	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	226	GLU	ASP	conflict	UNP P31800
A	227	THR	ALA	conflict	UNP P31800

- Molecule 2 is a protein called Cytochrome b-c1 complex subunit 2, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	413	Total	C	N	O	S	0	0	0
			3093	1943	546	597	7			

- Molecule 3 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	379	Total	C	N	O	S	0	0	0
			3004	2012	472	501	19			

- Molecule 4 is a protein called Cytochrome c1, heme protein, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	239	Total	C	N	O	S	0	0	0
			1884	1205	321	343	15			

- Molecule 5 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	196	Total	C	N	O	S	0	0	0
			1519	957	263	291	8			

- Molecule 6 is a protein called Cytochrome b-c1 complex subunit 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	99	Total	C	N	O	S	0	0	0
			859	545	156	156	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	56	ASP	ASN	conflict	UNP P00129

- Molecule 7 is a protein called Cytochrome b-c1 complex subunit 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	74	Total	C	N	O	S	0	0	0
			620	406	116	97	1			

- Molecule 8 is a protein called Cytochrome b-c1 complex subunit 6, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	65	Total	C	N	O	S	0	0	0
			525	318	95	107	5			

- Molecule 9 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	46	Total	C	N	O	S	0	0	0
			338	209	64	64	1			

- Molecule 10 is a protein called Cytochrome b-c1 complex subunit 9.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
10	J	59	Total	C	N	O	0	0	0
			487	320	84	83			

- Molecule 11 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			13	8	5		
11	C	1	Total	C	O	0	0
			13	8	5		
11	C	1	Total	C	O	0	0
			13	8	5		

- Molecule 12 is 1,2-DIHEXANOYL-SN-GLYCERO-3-PHOSPHOETHANOLAMINE (CCD ID: 6PE) (formula: C₁₇H₃₃NO₈P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	A	1	Total	C	N	O	P	0	0
			23	13	1	8	1		

- Molecule 13 is CARDIOLIPIN (CCD ID: CDL) (formula: $C_{81}H_{156}O_{17}P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
13	A	1	Total	C	O	P	0	0
			34	17	15	2		
13	D	1	Total	C	O	P	0	0
			54	35	17	2		
13	G	1	Total	C	O	P	0	0
			44	25	17	2		

- Molecule 14 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
14	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
14	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 15 is DODECYL-BETA-D-MALTOSE (CCD ID: LMT) (formula: $C_{24}H_{46}O_{11}$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	C	1	Total	C O	0	0
			35	24 11		


- Molecule 16 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (CCD ID: PEE) (formula:

C₄₁H₇₈NO₈P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
16	C	1	Total	C	N	O	P	0	0
			40	30	1	8	1		
16	E	1	Total	C	N	O	P	0	0
			41	31	1	8	1		

- Molecule 17 is 3-methyl-2-[4-[[4-(trifluoromethoxy)phenyl]methyl]phenyl]-1H-quinolin-4-one (CCD ID: I2Q) (formula: C₂₄H₁₈F₃NO₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
17	C	1	Total	C	F	N	O	0	0
			30	24	3	1	2		

- Molecule 18 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
18	C	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 19 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
19	C	1	Total	O	P	0	0
			5	4	1		
19	C	1	Total	O	P	0	0
			5	4	1		
19	D	1	Total	O	P	0	0
			5	4	1		
19	E	1	Total	O	P	0	0
			5	4	1		
19	F	1	Total	O	P	0	0
			5	4	1		
19	F	1	Total	O	P	0	0
			5	4	1		
19	G	1	Total	O	P	0	0
			5	4	1		
19	G	1	Total	O	P	0	0
			5	4	1		
19	G	1	Total	O	P	0	0
			5	4	1		

- Molecule 20 is HEME C (CCD ID: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
20	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 21 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
21	E	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 22 is 1,2-DIMYRISTOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PX4) (formula: C₃₆H₇₃NO₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
22	E	1	Total	C	N	O	P	0	0
			37	27	1	8	1		


- Molecule 23 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
23	A	10	Total 10	O 10	0	0
23	B	9	Total 9	O 9	0	0
23	C	10	Total 10	O 10	0	0
23	D	4	Total 4	O 4	0	0
23	E	4	Total 4	O 4	0	0
23	F	5	Total 5	O 5	0	0
23	G	2	Total 2	O 2	0	0
23	J	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome b-c1 complex subunit 1, mitochondrial




- Molecule 2: Cytochrome b-c1 complex subunit 2, mitochondrial





- Molecule 3: Cytochrome b








- Molecule 4: Cytochrome c1, heme protein, mitochondrial




- Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial



- Molecule 6: Cytochrome b-c1 complex subunit 7




- Molecule 7: Cytochrome b-c1 complex subunit 8







- Molecule 8: Cytochrome b-c1 complex subunit 6, mitochondrial



- Molecule 9: Cytochrome b-c1 complex subunit Rieske, mitochondrial



- Molecule 10: Cytochrome b-c1 complex subunit 9



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	209.59Å 209.59Å 342.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.86 – 3.20 49.86 – 3.20	Depositor EDS
% Data completeness (in resolution range)	95.1 (49.86-3.20) 95.1 (49.86-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.205 , 0.256 0.207 , 0.258	Depositor DCC
R_{free} test set	3444 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	85.7	Xtriage
Anisotropy	0.158	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 62.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	16382	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEE, CDL, LMT, PO4, I2Q, HEC, PX4, 6PE, PG4, HEM, FES, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.60	0/3522	1.31	9/4781 (0.2%)
2	B	0.60	0/3148	1.28	7/4270 (0.2%)
3	C	0.57	0/3100	1.21	6/4241 (0.1%)
4	D	0.61	0/1943	1.23	6/2642 (0.2%)
5	E	0.62	0/1553	1.21	2/2100 (0.1%)
6	F	0.55	0/878	1.33	1/1181 (0.1%)
7	G	0.61	0/641	1.39	3/868 (0.3%)
8	H	0.51	0/530	1.25	0/714
9	I	0.74	0/342	1.50	4/464 (0.9%)
10	J	0.55	0/500	1.24	1/675 (0.1%)
All	All	0.60	0/16157	1.27	39/21936 (0.2%)

There are no bond length outliers.

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	139	THR	CB-CA-C	7.29	121.82	109.72
3	C	368	THR	CA-CB-OG1	-6.82	99.38	109.60
1	A	445	ARG	N-CA-C	-6.37	106.47	114.56
3	C	345	HIS	CB-CA-C	6.29	118.54	109.11
3	C	219	PRO	N-CA-C	-6.19	101.34	111.68

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3449	0	3346	83	0
2	B	3093	0	3064	75	0
3	C	3004	0	3070	74	0
4	D	1884	0	1814	58	0
5	E	1519	0	1503	45	0
6	F	859	0	836	14	0
7	G	620	0	624	16	0
8	H	525	0	500	22	0
9	I	338	0	356	25	0
10	J	487	0	487	8	0
11	A	13	0	18	0	0
11	C	26	0	36	1	0
12	A	23	0	19	4	0
13	A	34	0	24	3	0
13	D	54	0	52	1	0
13	G	44	0	32	1	0
14	C	86	0	60	12	0
15	C	35	0	46	2	0
16	C	40	0	54	1	0
16	E	41	0	59	3	0
17	C	30	0	0	1	0
18	C	4	0	6	1	0
19	C	10	0	0	0	0
19	D	5	0	0	0	0
19	E	5	0	0	0	0
19	F	10	0	0	0	0
19	G	15	0	0	0	0
20	D	43	0	32	11	0
21	E	4	0	0	0	0
22	E	37	0	51	4	0
23	A	10	0	0	0	0
23	B	9	0	0	0	0
23	C	10	0	0	0	0
23	D	4	0	0	0	0
23	E	4	0	0	1	0
23	F	5	0	0	0	0
23	G	2	0	0	0	0
23	J	1	0	0	0	0
All	All	16382	0	16089	377	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 377 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
14:C:402:HEM:HBC2	14:C:402:HEM:HMC2	1.41	1.00
7:G:50:PRO:HB2	7:G:51:PRO:HD3	1.47	0.97
5:E:16:PRO:HA	5:E:19:LEU:HD12	1.51	0.93
1:A:86:LEU:HB3	2:B:285:VAL:HG22	1.54	0.89
1:A:4:TYR:HB3	2:B:114:ASP:OD2	1.75	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/445 (100%)	417 (94%)	25 (6%)	1 (0%)	43	73
2	B	409/418 (98%)	395 (97%)	14 (3%)	0	100	100
3	C	377/379 (100%)	354 (94%)	22 (6%)	1 (0%)	36	68
4	D	237/239 (99%)	224 (94%)	10 (4%)	3 (1%)	9	40
5	E	194/196 (99%)	187 (96%)	5 (3%)	2 (1%)	12	45
6	F	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
7	G	72/74 (97%)	69 (96%)	3 (4%)	0	100	100
8	H	63/65 (97%)	56 (89%)	7 (11%)	0	100	100
9	I	44/46 (96%)	42 (96%)	2 (4%)	0	100	100
10	J	57/59 (97%)	54 (95%)	3 (5%)	0	100	100
All	All	1993/2020 (99%)	1893 (95%)	93 (5%)	7 (0%)	30	62

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	THR
5	E	120	PRO
3	C	172	LYS
5	E	121	GLN
4	D	162	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/370 (100%)	351 (95%)	18 (5%)	22	55
2	B	322/327 (98%)	307 (95%)	15 (5%)	23	57
3	C	326/327 (100%)	303 (93%)	23 (7%)	13	44
4	D	200/204 (98%)	192 (96%)	8 (4%)	28	61
5	E	168/168 (100%)	152 (90%)	16 (10%)	8	32
6	F	88/91 (97%)	80 (91%)	8 (9%)	9	34
7	G	65/66 (98%)	62 (95%)	3 (5%)	24	58
8	H	61/62 (98%)	55 (90%)	6 (10%)	7	31
9	I	38/38 (100%)	27 (71%)	11 (29%)	0	1
10	J	49/49 (100%)	44 (90%)	5 (10%)	7	29
All	All	1686/1702 (99%)	1573 (93%)	113 (7%)	15	47

5 of 113 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	17	LEU
10	J	9	LEU
5	E	74	ILE
10	J	6	THR
9	I	47	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	161	HIS
8	H	67	HIS
6	F	27	ASN
8	H	23	GLN
1	A	363	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
19	PO4	C	409	-	4,4,4	0.54	0	6,6,6	0.69	0
11	PG4	C	405	-	12,12,12	0.30	0	11,11,11	0.29	0
19	PO4	F	502	-	4,4,4	0.70	0	6,6,6	0.49	0
11	PG4	A	501	-	12,12,12	0.24	0	11,11,11	0.16	0
13	CDL	A	503	-	33,33,99	0.52	0	37,43,111	0.87	0
11	PG4	C	404	-	12,12,12	0.42	0	11,11,11	0.36	0
19	PO4	F	501	-	4,4,4	1.26	1 (25%)	6,6,6	0.52	0
19	PO4	G	104	-	4,4,4	1.42	1 (25%)	6,6,6	0.55	0
13	CDL	G	101	-	43,43,99	0.67	1 (2%)	49,55,111	1.02	5 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
17	I2Q	C	407	-	33,33,33	0.81	1 (3%)	47,48,48	0.75	0
19	PO4	D	502	-	4,4,4	0.74	0	6,6,6	0.57	0
19	PO4	C	410	-	4,4,4	0.93	0	6,6,6	0.38	0
14	HEM	C	401	3	50,50,50	1.63	10 (20%)	67,82,82	1.74	16 (23%)
22	PX4	E	202	-	36,36,45	0.63	0	42,44,53	0.93	2 (4%)
18	DMS	C	408	-	3,3,3	0.35	0	3,3,3	0.27	0
19	PO4	G	103	-	4,4,4	0.98	0	6,6,6	0.48	0
12	6PE	A	502	-	22,22,26	0.51	0	25,27,31	0.90	1 (4%)
13	CDL	D	503	-	53,53,99	0.64	1 (1%)	59,65,111	0.83	1 (1%)
16	PEE	E	204	-	40,40,50	0.81	1 (2%)	43,45,55	0.89	1 (2%)
19	PO4	E	203	-	4,4,4	0.96	0	6,6,6	0.42	0
14	HEM	C	402	3	50,50,50	1.52	10 (20%)	67,82,82	1.66	13 (19%)
16	PEE	C	406	-	39,39,50	0.82	1 (2%)	42,44,55	0.75	0
21	FES	E	201	5	0,4,4	-	-	-	-	-
15	LMT	C	403	-	36,36,36	0.90	1 (2%)	47,47,47	1.66	13 (27%)
19	PO4	G	102	-	4,4,4	1.17	1 (25%)	6,6,6	0.41	0
20	HEC	D	501	4	46,50,50	2.70	25 (54%)	58,82,82	2.41	24 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	HEM	C	401	3	-	4/14/54/54	-
22	PX4	E	202	-	-	16/40/40/49	-
11	PG4	C	405	-	-	5/10/10/10	-
12	6PE	A	502	-	-	12/26/26/30	-
13	CDL	D	503	-	-	26/63/63/110	-
13	CDL	G	101	-	-	15/52/52/110	-
16	PEE	E	204	-	1/1/4/8	25/44/44/54	-
14	HEM	C	402	3	-	3/14/54/54	-
11	PG4	A	501	-	-	7/10/10/10	-
16	PEE	C	406	-	1/1/4/8	18/43/43/54	-
13	CDL	A	503	-	-	15/41/41/110	-
11	PG4	C	404	-	-	8/10/10/10	-
17	I2Q	C	407	-	-	2/13/13/13	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
21	FES	E	201	5	-	-	0/1/1/1
15	LMT	C	403	-	-	14/21/61/61	0/2/2/2
20	HEC	D	501	4	-	0/14/54/54	-

The worst 5 of 54 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
20	D	501	HEC	C2A-C3A	5.08	1.47	1.36
20	D	501	HEC	CHD-C4C	4.94	1.48	1.38
20	D	501	HEC	CHB-C4A	4.93	1.48	1.38
20	D	501	HEC	CAC-C3C	4.73	1.50	1.35
14	C	401	HEM	FE-NB	4.59	2.09	1.94

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
20	D	501	HEC	C2A-C1A-NA	7.33	117.40	110.32
14	C	402	HEM	CHC-C4B-NB	5.47	130.31	124.42
20	D	501	HEC	C3D-C4D-ND	5.41	116.16	110.15
20	D	501	HEC	C1A-C2A-C3A	-5.26	100.18	107.11
14	C	402	HEM	CMD-C2D-C1D	4.27	131.71	125.03

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
16	C	406	PEE	C2
16	E	204	PEE	C2

5 of 170 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
12	A	502	6PE	C1-O3-P1-O1
12	A	502	6PE	C1-O3-P1-O2
12	A	502	6PE	C1-O3-P1-O8
12	A	502	6PE	C11-C10-O6-C2
12	A	502	6PE	O8-C16-C17-N1


There are no ring outliers.


14 monomers are involved in 42 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	C	405	PG4	1	0
13	A	503	CDL	3	0
13	G	101	CDL	1	0
17	C	407	I2Q	1	0
14	C	401	HEM	5	0
22	E	202	PX4	4	0
18	C	408	DMS	1	0
12	A	502	6PE	4	0
13	D	503	CDL	1	0
16	E	204	PEE	3	0
14	C	402	HEM	7	0
16	C	406	PEE	1	0
15	C	403	LMT	2	0
20	D	501	HEC	11	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









Ligand 6PE A 502




Ligand CDL D 503






Ligand PEE E 204



Ligand HEM C 402







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/445 (100%)	-0.06	10 (2%) 62 42	47, 76, 111, 231	0
2	B	413/418 (98%)	-0.25	7 (1%) 69 49	45, 80, 110, 175	0
3	C	379/379 (100%)	-0.02	14 (3%) 45 28	44, 68, 102, 148	0
4	D	239/239 (100%)	0.22	7 (2%) 53 35	58, 95, 125, 159	0
5	E	196/196 (100%)	1.04	41 (20%) 2 2	52, 132, 194, 249	1 (0%)
6	F	99/99 (100%)	-0.11	8 (8%) 18 11	50, 71, 103, 139	0
7	G	74/74 (100%)	0.24	6 (8%) 18 11	48, 79, 119, 138	0
8	H	65/65 (100%)	0.35	3 (4%) 37 23	86, 121, 152, 168	0
9	I	46/46 (100%)	1.24	10 (21%) 2 2	63, 119, 155, 184	0
10	J	59/59 (100%)	-0.09	1 (1%) 69 49	61, 82, 118, 145	0
All	All	2015/2020 (99%)	0.10	107 (5%) 32 20	44, 81, 144, 249	1 (0%)

The worst 5 of 107 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	115	SER	9.2
5	E	157	TYR	6.8
5	E	114	VAL	5.2
3	C	263	ASN	5.1
5	E	163	SER	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	LMT	C	403	35/35	0.68	0.18	77,130,187,224	0
19	PO4	E	203	5/5	0.72	0.17	158,174,183,211	0
19	PO4	C	410	5/5	0.77	0.11	144,157,159,166	0
19	PO4	G	104	5/5	0.80	0.15	124,124,147,165	0
17	I2Q	C	407	30/30	0.83	0.25	96,131,213,222	0
19	PO4	F	501	5/5	0.85	0.13	93,118,139,145	0
19	PO4	G	103	5/5	0.86	0.13	124,127,136,140	0
19	PO4	C	409	5/5	0.88	0.14	103,143,165,191	0
13	CDL	D	503	54/100	0.88	0.19	52,102,135,143	0
19	PO4	D	502	5/5	0.90	0.11	141,167,186,220	0
19	PO4	F	502	5/5	0.90	0.11	161,163,191,198	0
11	PG4	A	501	13/13	0.91	0.13	72,108,128,129	0
21	FES	E	201	4/4	0.91	0.14	144,185,205,211	2
11	PG4	C	405	13/13	0.92	0.19	70,88,99,106	0
13	CDL	A	503	34/100	0.92	0.12	81,108,137,148	0
19	PO4	G	102	5/5	0.93	0.21	97,105,113,125	0
13	CDL	G	101	44/100	0.93	0.15	63,95,135,146	0
22	PX4	E	202	37/46	0.93	0.17	55,88,141,175	0
16	PEE	E	204	41/51	0.94	0.14	60,78,90,103	0
18	DMS	C	408	4/4	0.95	0.26	129,139,144,167	0
12	6PE	A	502	23/27	0.95	0.13	46,100,122,148	0
16	PEE	C	406	40/51	0.95	0.12	56,74,92,97	0
11	PG4	C	404	13/13	0.96	0.12	72,86,107,117	0
20	HEC	D	501	43/43	0.96	0.12	72,91,113,127	0
14	HEM	C	401	43/43	0.98	0.09	54,68,73,75	0
14	HEM	C	402	43/43	0.99	0.07	49,57,66,71	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LMT C 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around I2Q C 407:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CDL D 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CDL A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CDL G 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PX4 E 202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PEE E 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 6PE A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PEE C 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around HEC D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.