



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 27, 2026 – 08:26 PM UTC

PDB ID : 8R38 / pdb_00008r38
Title : BIIG2 anti-integrin Fab
Authors : Cordara, G.; Heim, J.B.; Johannesen, H.; Krengel, U.
Deposited on : 2023-11-08
Resolution : 1.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4403 (1.40-1.36)
Clashscore	190562	4528 (1.40-1.36)
Ramachandran outliers	187476	4459 (1.40-1.36)
Sidechain outliers	187428	4458 (1.40-1.36)
RSRZ outliers	180081	4399 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	220	
2	L	214	
3	S	4	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4185 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BIIG2 Fab, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	217	Total	C	N	O	S	0	46	0
			1974	1222	332	405	15			

- Molecule 2 is a protein called BIIG2 Fab, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	32	0
			1906	1179	331	383	13			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	S	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	116	Total	O	0	0
			116	116		
6	L	108	Total	O	0	0
			108	108		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BIIG2 Fab, heavy chain



- Molecule 2: BIIG2 Fab, light chain



- Molecule 3: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.93Å 80.88Å 87.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.38 – 1.38 59.38 – 1.38	Depositor EDS
% Data completeness (in resolution range)	99.3 (59.38-1.38) 99.3 (59.38-1.38)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 1.38Å)	Xtriage
Refinement program	REFMAC 5.8.0415	Depositor
R, R_{free}	0.195 , 0.217 0.200 , 0.222	Depositor DCC
R_{free} test set	4717 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 25.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4185	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, PCA, GOL, NAG, XRE, CL, MLZ, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.63	0/2000	0.97	1/2726 (0.0%)
2	L	0.64	0/1930	0.95	2/2601 (0.1%)
All	All	0.63	0/3930	0.96	3/5327 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	2
2	L	0	3
All	All	0	5

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	L	83	PHE	CB-CA-C	5.69	119.12	109.84
1	H	1	PCA	O-C-N	5.63	132.01	123.00
2	L	64	GLY	O-C-N	5.20	127.95	123.37

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	42[A]	GLY	Peptide
1	H	66[A]	ARG	Sidechain
2	L	142[A]	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	L	211	ARG	Sidechain
2	L	28[A]	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1974	0	1888	34	0
2	L	1906	0	1858	29	0
3	S	50	0	43	0	0
4	H	6	0	8	1	0
4	L	24	0	32	1	0
5	L	1	0	0	0	0
6	H	116	0	0	2	0
6	L	108	0	0	0	0
All	All	4185	0	3829	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:142[B]:ARG:O	2:L:142[B]:ARG:HD2	1.48	1.13
1:H:41[B]:PRO:HG3	1:H:87[B]:THR:OG1	1.60	1.02
1:H:40:PRO:HA	1:H:88[B]:THR:HG22	1.47	0.95
2:L:136[B]:MET:SD	2:L:146:VAL:HG21	2.10	0.91
2:L:142[B]:ARG:HH11	2:L:142[B]:ARG:HG2	1.35	0.91
1:H:41[B]:PRO:HG3	1:H:87[B]:THR:HG1	1.40	0.86
1:H:88[B]:THR:HG21	6:H:467:HOH:O	1.76	0.84
2:L:136[B]:MET:SD	2:L:146:VAL:CG2	2.68	0.81
2:L:142[B]:ARG:HG2	2:L:142[B]:ARG:NH1	1.93	0.79
1:H:107[B]:VAL:HG22	6:H:419:HOH:O	1.82	0.78
2:L:142[B]:ARG:HB2	2:L:173:TYR:CE2	2.21	0.75
2:L:142[B]:ARG:NH1	2:L:142[B]:ARG:CG	2.49	0.75
1:H:16[B]:GLU:O	1:H:82(C)[B]:LEU:HD13	1.88	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:4:MET:CE	2:L:23[B]:CYS:SG	2.77	0.72
2:L:142[B]:ARG:HD2	2:L:142[B]:ARG:C	2.16	0.70
2:L:195[A]:GLU:HG2	2:L:206:VAL:HG22	1.75	0.69
1:H:14[B]:PRO:HB3	1:H:83[B]:GLN:OE1	1.93	0.69
1:H:13[A]:GLN:HG2	1:H:14[A]:PRO:HD2	1.74	0.68
1:H:83[B]:GLN:O	1:H:84[B]:SER:OG	2.15	0.65
2:L:4:MET:HE3	2:L:23[B]:CYS:SG	2.38	0.62
2:L:150:ILE:HD12	2:L:155[B]:ARG:HG3	1.82	0.62
2:L:61:ARG:HB2	2:L:76[B]:ARG:O	2.02	0.60
2:L:4:MET:HE2	2:L:23[B]:CYS:SG	2.42	0.59
1:H:177[B]:LEU:HD12	1:H:177[B]:LEU:C	2.29	0.58
1:H:16[B]:GLU:O	1:H:82(C)[B]:LEU:CD1	2.51	0.58
1:H:83[B]:GLN:O	1:H:84[B]:SER:CB	2.50	0.57
1:H:83[B]:GLN:C	1:H:84[B]:SER:OG	2.46	0.57
1:H:86[B]:ASP:C	1:H:87[B]:THR:HG23	2.31	0.56
1:H:67:LEU:HD21	1:H:82[B]:MET:HE3	1.89	0.55
1:H:41[B]:PRO:HG3	1:H:87[B]:THR:O	2.08	0.54
2:L:142[B]:ARG:O	2:L:142[B]:ARG:CD	2.40	0.53
1:H:13[A]:GLN:HG3	1:H:113[A]:SER:HA	1.89	0.53
2:L:110:ASP:OD2	2:L:199:MLZ:HCM3	2.08	0.53
2:L:136[B]:MET:CE	2:L:146:VAL:HG21	2.38	0.53
2:L:47[B]:LEU:C	2:L:48[B]:ILE:HG13	2.34	0.52
2:L:47[B]:LEU:O	2:L:48[B]:ILE:HG13	2.11	0.51
1:H:107[B]:VAL:O	1:H:107[B]:VAL:HG13	2.11	0.50
1:H:165:THR:HG23	1:H:177[A]:LEU:HD21	1.94	0.49
2:L:8:PRO:HG2	2:L:11[B]:LEU:HG	1.94	0.49
1:H:18[B]:LEU:HD12	1:H:19[B]:SER:H	1.79	0.48
2:L:136[B]:MET:SD	2:L:146:VAL:HG22	2.51	0.48
1:H:145:TYR:CE2	1:H:150[B]:VAL:HG23	2.49	0.48
2:L:150:ILE:HD11	2:L:179:LEU:HD21	1.96	0.48
1:H:177[B]:LEU:HD12	1:H:177[B]:LEU:O	2.14	0.47
2:L:181:LEU:HD23	2:L:181:LEU:N	2.30	0.46
2:L:142[B]:ARG:HB2	2:L:173:TYR:CD2	2.51	0.46
1:H:41[B]:PRO:CG	1:H:87[B]:THR:OG1	2.49	0.46
1:H:128[A]:CYS:SG	2:L:119:PRO:HG2	2.55	0.46
1:H:41[B]:PRO:HD3	1:H:88[B]:THR:HG22	1.99	0.45
1:H:82(A)[A]:ASN:O	1:H:82(B)[A]:SER:HB2	2.16	0.45
2:L:69:THR:OG1	4:L:303:GOL:H12	2.17	0.45
2:L:136[B]:MET:HE1	2:L:146:VAL:HG21	1.98	0.45
1:H:177[B]:LEU:HA	4:H:301:GOL:O3	2.17	0.44
1:H:63:LEU:HD23	1:H:66[B]:ARG:NH1	2.33	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:83[A]:GLN:O	1:H:84[A]:SER:C	2.60	0.44
2:L:8:PRO:HG3	2:L:11[B]:LEU:HD21	2.00	0.44
2:L:11[A]:LEU:HD13	2:L:19:VAL:HB	1.99	0.43
1:H:82(A)[B]:ASN:HA	1:H:82(C)[B]:LEU:CD1	2.49	0.42
1:H:18[A]:LEU:HD23	1:H:82[A]:MET:SD	2.60	0.42
1:H:82(A)[B]:ASN:HA	1:H:82(C)[B]:LEU:HD12	2.01	0.41
1:H:100(B):THR:HB	1:H:100(C):XRE:OD	2.21	0.41
1:H:14[B]:PRO:O	1:H:82(C)[B]:LEU:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	260/220 (118%)	250 (96%)	8 (3%)	2 (1%)	16	3
2	L	242/214 (113%)	238 (98%)	4 (2%)	0	100	100
All	All	502/434 (116%)	488 (97%)	12 (2%)	2 (0%)	43	10

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	82(C)[A]	LEU
1	H	82(C)[B]	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	230/192 (120%)	229 (100%)	1 (0%)	84	68
2	L	219/187 (117%)	218 (100%)	1 (0%)	81	62
All	All	449/379 (118%)	447 (100%)	2 (0%)	81	68

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	204	VAL
2	L	156	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	XRE	H	100(C)	1	7,8,9	1.05	0	7,10,12	1.84	2 (28%)
2	MLZ	L	199	2	8,9,10	0.43	0	4,9,11	0.72	0
1	PCA	H	1	1	7,8,9	0.84	0	9,10,12	1.02	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '–' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	XRE	H	100(C)	1	-	0/0/11/13	0/1/1/1
2	MLZ	L	199	2	-	3/7/8/10	-
1	PCA	H	1	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	100(C)	XRE	CB-CA-C	3.02	116.80	112.66
1	H	100(C)	XRE	CG-CB-CA	-2.67	100.09	103.75
1	H	1	PCA	O-C-CA	-2.26	118.95	124.77

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	L	199	MLZ	CD-CE-NZ-CM
2	L	199	MLZ	CA-CB-CG-CD
2	L	199	MLZ	CE-CD-CG-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	H	100(C)	XRE	1	0
2	L	199	MLZ	1	0

5.5 Carbohydrates

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	S	1	1,3	14,14,15	0.65	0	17,19,21	0.94	2 (11%)
3	NAG	S	2	3	14,14,15	0.52	0	17,19,21	0.53	0
3	BMA	S	3	3	11,11,12	0.56	0	15,15,17	1.00	1 (6%)
3	MAN	S	4	3	11,11,12	0.79	0	15,15,17	1.06	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	S	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	S	2	3	-	0/6/23/26	0/1/1/1
3	BMA	S	3	3	-	0/2/19/22	0/1/1/1
3	MAN	S	4	3	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	S	3	BMA	O2-C2-C3	2.96	116.28	110.15
3	S	4	MAN	O2-C2-C3	2.35	115.01	110.15
3	S	4	MAN	C1-O5-C5	2.27	115.23	112.19
3	S	1	NAG	C1-C2-N2	-2.25	106.89	110.43
3	S	1	NAG	O5-C1-C2	-2.04	108.13	111.29

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	S	4	MAN	C4-C5-C6-O6
3	S	4	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	L	304	-	5,5,5	0.12	0	5,5,5	0.24	0
4	GOL	L	301	-	5,5,5	0.11	0	5,5,5	0.47	0
4	GOL	L	303	-	5,5,5	0.14	0	5,5,5	0.59	0
4	GOL	L	302	-	5,5,5	0.41	0	5,5,5	0.69	0
4	GOL	H	301	-	5,5,5	0.27	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	304	-	-	4/4/4/4	-
4	GOL	L	301	-	-	1/4/4/4	-
4	GOL	L	303	-	-	0/4/4/4	-
4	GOL	L	302	-	-	2/4/4/4	-
4	GOL	H	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	302	GOL	O1-C1-C2-C3
4	L	304	GOL	O1-C1-C2-C3
4	L	304	GOL	O1-C1-C2-O2
4	L	301	GOL	C1-C2-C3-O3
4	L	304	GOL	C1-C2-C3-O3
4	L	304	GOL	O2-C2-C3-O3
4	L	302	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	303	GOL	1	0
4	H	301	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	215/220 (97%)	1.04	40 (18%) 3 4	8, 21, 49, 94	46 (21%)
2	L	213/214 (99%)	0.68	24 (11%) 10 12	7, 21, 37, 65	32 (15%)
All	All	428/434 (98%)	0.86	64 (14%) 5 7	7, 21, 43, 94	78 (18%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	84[A]	SER	6.1
1	H	132	THR	5.3
1	H	99	GLY	5.2
1	H	82(C)[A]	LEU	5.1
1	H	204	VAL	4.7
1	H	131	THR	3.9
1	H	82(B)[A]	SER	3.9
1	H	159	LEU	3.6
1	H	210	ARG	3.5
2	L	212	ASN	3.5
2	L	201	SER	3.5
1	H	128[A]	CYS	3.5
1	H	187	PRO	3.5
1	H	186	TRP	3.5
1	H	158	ALA	3.4
2	L	156	ARG	3.4
1	H	190	THR	3.2
1	H	133	SER	3.2
2	L	122[A]	THR	3.2
1	H	130[A]	ASP	3.2
1	H	98	MET	3.1
2	L	127	THR	3.0
1	H	136	VAL	3.0
1	H	82(A)[A]	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	214[A]	CYS	3.0
1	H	185	THR	2.9
2	L	152	GLY	2.8
2	L	129	GLY	2.8
1	H	188	SER	2.7
2	L	197	VAL	2.7
2	L	147[A]	LYS	2.7
1	H	83[A]	GLN	2.7
1	H	196	ALA	2.7
1	H	135	THR	2.6
1	H	208	VAL	2.5
1	H	97	THR	2.5
1	H	160	SER	2.4
1	H	161	SER	2.4
1	H	134	SER	2.4
2	L	188	SER	2.4
1	H	189[A]	GLN	2.4
2	L	150	ILE	2.4
1	H	157	GLY	2.4
2	L	130	ALA	2.4
2	L	184	ALA	2.4
1	H	191	VAL	2.3
1	H	184	SER	2.3
2	L	157[A]	ASP	2.3
2	L	158	GLY	2.3
1	H	85[A]	GLU	2.3
2	L	202[A]	SER	2.2
2	L	40	PRO	2.2
1	H	201	SER	2.2
2	L	200	THR	2.2
1	H	121	VAL	2.2
1	H	207	LYS	2.1
2	L	160	LEU	2.1
1	H	42[A]	GLY	2.1
2	L	3	GLN	2.1
1	H	62	THR	2.1
2	L	126	ALA	2.1
1	H	5[A]	MET	2.1
2	L	182	THR	2.0
2	L	41	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MLZ	L	199	10/11	0.84	0.18	28,37,53,54	0
1	XRE	H	100(C)	8/9	0.95	0.09	18,20,23,31	1
1	PCA	H	1	8/9	0.97	0.06	20,21,23,25	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	S	1	14/15	-	-	19,22,26,28	0
3	NAG	S	2	14/15	-	-	28,33,42,43	0
3	BMA	S	3	11/12	-	-	33,44,51,53	0
3	MAN	S	4	11/12	-	-	46,51,54,55	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	L	304	6/6	0.85	0.13	38,41,48,49	0
4	GOL	L	303	6/6	0.88	0.13	23,39,41,47	0
4	GOL	L	301	6/6	0.89	0.12	22,34,35,38	0
4	GOL	L	302	6/6	0.91	0.12	22,30,36,39	0
4	GOL	H	301	6/6	0.94	0.09	22,25,29,30	0
5	CL	L	305	1/1	0.98	0.06	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.