



wwPDB EM Validation Summary Report ⓘ

Jul 14, 2026 – 05:20 pm BST

PDB ID : 9R3S / pdb_00009r3s
EMDB ID : EMD-53558
Title : pro-TGF-beta1 in complex with the third TB Domain from Latent Transforming Growth Factor-beta Binding Protein-1
Authors : Biggin, G.; Snee, M.; Godwin, A.; Roseman, A.; Baldock, C.
Deposited on : 2025-05-06
Resolution : 3.06 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13976 (2.56 - 3.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	 72% 18% 10%
1	B	369	 72% 17% 11%
2	C	435	 11% 86%
3	D	3	 67% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	3	 33% 67%
4	E	4	 25% 50% 25%
4	G	4	 75% 25%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11803 atoms, of which 5800 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transforming growth factor beta-1 proprotein.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	B	328	Total	C	H	N	O	S	0	0
			5250	1678	2613	463	478	18		
1	A	332	Total	C	H	N	O	S	2	0
			5323	1704	2646	471	484	18		

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	22	ASP	-	expression tag	UNP P01137
B	23	TYR	-	expression tag	UNP P01137
B	24	LYS	-	expression tag	UNP P01137
B	25	ASP	-	expression tag	UNP P01137
B	26	ASP	-	expression tag	UNP P01137
B	27	ASP	-	expression tag	UNP P01137
B	28	ASP	-	expression tag	UNP P01137
B	29	LYS	-	expression tag	UNP P01137
A	22	ASP	-	expression tag	UNP P01137
A	23	TYR	-	expression tag	UNP P01137
A	24	LYS	-	expression tag	UNP P01137
A	25	ASP	-	expression tag	UNP P01137
A	26	ASP	-	expression tag	UNP P01137
A	27	ASP	-	expression tag	UNP P01137
A	28	ASP	-	expression tag	UNP P01137
A	29	LYS	-	expression tag	UNP P01137

- Molecule 2 is a protein called Latent-transforming growth factor beta-binding protein 1.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	C	63	Total	C	H	N	O	S	0	0
			869	284	411	73	92	9		

There are 49 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	1331	ALA	-	expression tag	UNP Q14766
C	1332	PRO	-	expression tag	UNP Q14766
C	1333	LEU	-	expression tag	UNP Q14766
C	1334	ALA	-	expression tag	UNP Q14766
C	1721	THR	-	expression tag	UNP Q14766
C	1722	GLY	-	expression tag	UNP Q14766
C	1723	GLY	-	expression tag	UNP Q14766
C	1724	GLY	-	expression tag	UNP Q14766
C	1725	GLY	-	expression tag	UNP Q14766
C	1726	SER	-	expression tag	UNP Q14766
C	1727	GLY	-	expression tag	UNP Q14766
C	1728	GLY	-	expression tag	UNP Q14766
C	1729	GLY	-	expression tag	UNP Q14766
C	1730	GLY	-	expression tag	UNP Q14766
C	1731	SER	-	expression tag	UNP Q14766
C	1732	GLY	-	expression tag	UNP Q14766
C	1733	GLY	-	expression tag	UNP Q14766
C	1734	GLY	-	expression tag	UNP Q14766
C	1735	GLY	-	expression tag	UNP Q14766
C	1736	SER	-	expression tag	UNP Q14766
C	1737	ALA	-	expression tag	UNP Q14766
C	1738	TRP	-	expression tag	UNP Q14766
C	1739	SER	-	expression tag	UNP Q14766
C	1740	HIS	-	expression tag	UNP Q14766
C	1741	PRO	-	expression tag	UNP Q14766
C	1742	GLN	-	expression tag	UNP Q14766
C	1743	PHE	-	expression tag	UNP Q14766
C	1744	GLU	-	expression tag	UNP Q14766
C	1745	LYS	-	expression tag	UNP Q14766
C	1746	GLY	-	expression tag	UNP Q14766
C	1747	GLY	-	expression tag	UNP Q14766
C	1748	GLY	-	expression tag	UNP Q14766
C	1749	SER	-	expression tag	UNP Q14766
C	1750	GLY	-	expression tag	UNP Q14766
C	1751	GLY	-	expression tag	UNP Q14766
C	1752	GLY	-	expression tag	UNP Q14766
C	1753	SER	-	expression tag	UNP Q14766
C	1754	GLY	-	expression tag	UNP Q14766
C	1755	GLY	-	expression tag	UNP Q14766
C	1756	SER	-	expression tag	UNP Q14766
C	1757	ALA	-	expression tag	UNP Q14766
C	1758	TRP	-	expression tag	UNP Q14766
C	1759	SER	-	expression tag	UNP Q14766

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	1760	HIS	-	expression tag	UNP Q14766
C	1761	PRO	-	expression tag	UNP Q14766
C	1762	GLN	-	expression tag	UNP Q14766
C	1763	PHE	-	expression tag	UNP Q14766
C	1764	GLU	-	expression tag	UNP Q14766
C	1765	LYS	-	expression tag	UNP Q14766

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	3	Total	C	H	N	O	0	0
			67	22	28	2	15		
3	F	3	Total	C	H	N	O	0	0
			67	22	28	2	15		

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	4	Total	C	H	N	O	0	0
			80	28	31	2	19		
4	G	4	Total	C	H	N	O	0	0
			80	28	31	2	19		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					AltConf
5	C	1	Total	C	H	N	O	0
			26	8	12	1	5	


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		AltConf
6	B	18	Total	O	0
			18	18	
6	A	22	Total	O	0
			22	22	
6	C	1	Total	O	0
			1	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transforming growth factor beta-1 proprotein



- Molecule 1: Transforming growth factor beta-1 proprotein



- Molecule 2: Latent-transforming growth factor beta-binding protein 1



GLY	GLY	ASN	ALA	GLN	THR	ASP	GLN	ASN
GLY	GLY	GLU	ASP	THR	THR	THR	THR	THR
SER	SER	CYS	PRO	THR	GLU	GLY	THR	THR
GLY	GLY	ASP	TYR	PHE	THR	THR	THR	SER
GLY	GLY	LEU	ILE	ILE	GLU	TYR	GLY	TYR
GLY	GLY	LEU	GLN	GLN	CYS	CYS	CYS	ASN
SER	SER	ASN	ASP	ASP	CYS	CYS	CYS	ASN
GLY	GLY	ARG	ARG	ARG	CYS	CYS	CYS	PHE
GLY	GLY	MET	PHE	PHE	LEU	LEU	LEU	CYS
GLY	GLY	SER	LEU	LEU	GLY	GLY	GLY	THR
GLY	GLY	LEU	ASN	ASN	TYR	TYR	ARG	ASP
SER	SER	CYS	SER	SER	GLU	LYS	ARG	LEU
ALA	ALA	LYS	PHE	PHE	ALA	ASP	PRO	CYS
TRP	TRP	ASN	GLU	GLU	TRP	GLY	PRO	TRP
SER	SER	ALA	GLU	GLU	GLY	MET	VAL	TYR
HIS	HIS	LYS	LEU	LEU	GLN	GLN	GLN	HIS
PRO	PRO	CYS	GLN	GLN	CYS	CYS	CYS	LEU
GLN	GLN	ILE	ALA	ALA	CYS	CYS	CYS	ASP
PHE	PHE	ASN	GLU	GLU	ALA	ALA	ALA	ALA
GLU	GLU	THR	GLU	GLU	LEU	LEU	LEU	SER
LYS	LYS	ASP	CYS	CYS	CYS	CYS	CYS	GLY
GLY	GLY	GLY	GLY	GLY	PRO	PRO	PRO	ILE
GLY	GLY	SER	SER	SER	LEU	LEU	LEU	ARG
SER	SER	LYS	CYS	CYS	ASN	ASN	ASN	ARG
GLY	GLY	TYR	LEU	LEU	LYS	LYS	LYS	THR
SER	SER	GLY	GLY	GLY	ASP	SER	ASP	PRO
GLY	GLY	CYS	CYS	CYS	GLU	GLU	GLU	ALA
GLY	GLY	LEU	ASN	ASN	GLY	GLY	GLY	GLU
SER	SER	PRO	ARG	ARG	ALA	ALA	ALA	GLN
GLY	GLY	TYR	VAL	VAL	VAL	VAL	VAL	THR
TRP	TRP	PRO	SER	SER	ASN	ASN	ASN	ASP
SER	SER	ASP	GLN	GLN	PRO	PRO	PRO	THR
HIS	HIS	LYS	GLU	GLU	VAL	VAL	VAL	CYS
PRO	PRO	GLY	GLY	GLY	THR	THR	THR	GLN
GLN	GLN	PRO	GLY	GLY	THR	THR	THR	TYR
PHE	PHE	ASN	TYR	TYR	GLY	GLY	GLY	ASP
GLU	GLU	TYR	THR	THR	ARG	ARG	ARG	LEU
LYS	LYS	CYS	CYS	CYS	ILE	ILE	ILE	LEU
		THR	ASP	THR	THR	THR	THR	CYS
		ASP	GLN	GLN	PRO	PRO	PRO	TRP
		LYS	VAL	VAL	VAL	VAL	VAL	TYR
		GLY	THR	THR	GLY	GLY	GLY	GLY
		ASP	ARG	ARG	ASP	ASP	ASP	ASP
		LEU	CYS	CYS	LEU	LEU	LEU	LEU
		GLN	ASP	ASP	GLN	GLN	GLN	GLN
		THR	ASP	ASP	THR	THR	THR	THR
		CYS	CYS	CYS	CYS	CYS	CYS	CYS
		SER	LYS	LYS	SER	SER	SER	SER
		ASP	MET	MET	GLU	GLU	GLU	GLU
		THR	THR	THR	THR	THR	THR	THR
		VAL	VAL	VAL	VAL	VAL	VAL	VAL
		GLY	ASP	PRO	GLY	GLY	GLY	GLY

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 67% 33%

NAG1
NAG2
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  33% 67%


NAG1	NAG2	BMA3
------	------	------

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:

NAG1	NAG2	BMA3	FUC4
------	------	------	------

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  75% 25%

NAG1	NAG2	BMA3	FUC4
------	------	------	------

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	124397	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50.0	Depositor
Minimum defocus (nm)	750	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	165000	Depositor
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.231	Depositor
Minimum map value	-0.564	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.020	Depositor
Recommended contour level	0.0821	Depositor
Map size (Å)	276.48, 276.48, 276.48	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.08, 1.08, 1.08	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, FUC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/2748	0.31	0/3721
1	B	0.17	0/2700	0.30	0/3656
2	C	0.18	0/467	0.42	0/634
All	All	0.17	0/5915	0.32	0/8011

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2677	2646	2650	45	0
1	B	2637	2613	2617	43	0
2	C	458	411	411	13	0
3	D	39	28	34	1	0
3	F	39	28	34	1	0
4	E	49	31	43	4	0
4	G	49	31	43	2	0
5	C	14	12	13	0	0
6	A	22	0	0	0	0
6	B	18	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	1	0	0	0	0
All	All	6003	5800	5845	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 98 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1351:TYR:O	2:C:1361:ASN:ND2	2.16	0.78
1:B:133:MET:SD	1:B:157:LEU:HD11	2.29	0.72
4:G:1:NAG:H83	4:G:1:NAG:H3	1.72	0.71
4:E:2:NAG:H83	4:E:2:NAG:H3	1.72	0.71
1:B:112:LEU:HD23	1:B:257:PHE:HB3	1.76	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/369 (88%)	295 (90%)	31 (10%)	0	100	100
1	B	320/369 (87%)	288 (90%)	32 (10%)	0	100	100
2	C	61/435 (14%)	48 (79%)	13 (21%)	0	100	100
All	All	707/1173 (60%)	631 (89%)	76 (11%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/330 (90%)	294 (99%)	2 (1%)	76	80
1	B	293/330 (89%)	290 (99%)	3 (1%)	68	77
2	C	50/367 (14%)	49 (98%)	1 (2%)	48	69
All	All	639/1027 (62%)	633 (99%)	6 (1%)	68	78

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	CYS
1	A	293	CYS
2	C	1374	CYS
1	B	285	CYS
1	B	174	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	237	ASN
1	A	129	HIS
1	B	345	GLN
1	B	233	GLN
1	B	346	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	1	1,3	14,14,15	0.24	0	17,19,21	0.71	1 (5%)
3	NAG	D	2	3	14,14,15	0.19	0	17,19,21	0.39	0
3	BMA	D	3	3	11,11,12	0.57	0	15,15,17	0.78	0
4	NAG	E	1	4,1	14,14,15	0.23	0	17,19,21	0.40	0
4	NAG	E	2	4	14,14,15	0.39	0	17,19,21	1.18	2 (11%)
4	BMA	E	3	4	11,11,12	0.58	0	15,15,17	0.93	0
4	FUC	E	4	4	10,10,11	0.58	0	14,14,16	0.83	0
3	NAG	F	1	1,3	14,14,15	0.22	0	17,19,21	0.57	0
3	NAG	F	2	3	14,14,15	0.23	0	17,19,21	0.41	0
3	BMA	F	3	3	11,11,12	0.54	0	15,15,17	0.80	0
4	NAG	G	1	4,1	14,14,15	0.25	0	17,19,21	0.93	1 (5%)
4	NAG	G	2	4	14,14,15	0.25	0	17,19,21	0.45	0
4	BMA	G	3	4	11,11,12	0.55	0	15,15,17	0.71	0
4	FUC	G	4	4	10,10,11	0.67	0	14,14,16	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	1/2/19/22	0/1/1/1
4	NAG	E	1	4,1	-	4/6/23/26	0/1/1/1
4	NAG	E	2	4	-	5/6/23/26	0/1/1/1
4	BMA	E	3	4	-	1/2/19/22	0/1/1/1
4	FUC	E	4	4	-	-	0/1/1/1
3	NAG	F	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
4	NAG	G	1	4,1	-	3/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	FUC	G	4	4	-	-	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	2	NAG	C1-O5-C5	2.79	115.98	112.19
4	G	1	NAG	C2-N2-C7	2.67	126.70	122.90
4	E	2	NAG	C2-N2-C7	2.64	126.67	122.90
3	D	1	NAG	C1-O5-C5	2.54	115.63	112.19

There are no chirality outliers.

5 of 20 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	G	2	NAG	O5-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6
4	E	2	NAG	O5-C5-C6-O6
4	E	1	NAG	O5-C5-C6-O6
3	F	2	NAG	C4-C5-C6-O6


There are no ring outliers.


7 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	4	FUC	1	0
4	E	2	NAG	4	0
4	G	1	NAG	2	0
3	F	3	BMA	1	0
4	E	3	BMA	1	0
3	F	2	NAG	1	0
3	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	C	1801	2	14,14,15	0.19	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1801	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	1801	NAG	C8-C7-N2-C2
5	C	1801	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-53558. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 131




Y Index: 114




Z Index: 139


6.3.2 Raw map



X Index: 131



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map




6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0821. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 51 nm^3 ; this corresponds to an approximate mass of 46 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.327 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.327 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.06	-	-
Author-provided FSC curve	3.09	3.62	3.11
Unmasked-calculated*	3.93	7.18	4.00

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.93 differs from the reported value 3.06 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-53558 and PDB model 9R3S. Per-residue inclusion information can be found in [section 3](#) on [page 8](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0821 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0821).

9.4 Atom inclusion [i](#)



At the recommended contour level, 88% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0821) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8770	0.5190
A	0.8870	0.5310
B	0.9000	0.5300
C	0.8020	0.4500
D	0.7690	0.3880
E	0.7350	0.3780
F	0.6150	0.3670
G	0.7140	0.3550

1.0

0.0

<0.0