



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:24 AM UTC

PDB ID : 2R4G / pdb_00002r4g
Title : The high resolution structure of the RNA-binding domain of telomerase
Authors : Rouda, S.; Skordalakes, E.
Deposited on : 2007-08-31
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	15% 83% 9% • 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2364 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Telomerase reverse transcriptase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			2145	1401	354	383	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	253	MET	-	initiating methionine	UNP O77448

- Molecule 2 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	7	Total	Br	0	0
			7	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	212	Total	O	0	0
			212	212		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Telomerase reverse transcriptase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.35Å 67.22Å 51.53Å 90.00° 90.66° 90.00°	Depositor
Resolution (Å)	20.00 – 1.71 20.00 – 1.71	Depositor EDS
% Data completeness (in resolution range)	99.9 (20.00-1.71) 99.8 (20.00-1.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.200 , 0.239 (Not available) , 0.236	Depositor DCC
R_{free} test set	1472 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.034 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2364	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2187	0.65	1/2931 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	482	HIS	CB-CA-C	-5.02	109.21	116.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2145	0	2210	22	0
2	A	7	0	0	2	0
3	A	212	0	0	3	0
All	All	2364	0	2210	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:HIS:HD2	1:A:282:GLN:H	1.20	0.88
1:A:481:LYS:HB3	1:A:489:PHE:CE2	2.12	0.84
1:A:349:GLN:HA	1:A:350:SER:HB2	1.73	0.70
1:A:424:CYS:HB3	3:A:718:HOH:O	1.94	0.68
1:A:371:ILE:HD11	1:A:492:ARG:HG2	1.76	0.66
1:A:280:HIS:CD2	1:A:282:GLN:H	2.12	0.58
1:A:264:LYS:HE2	1:A:408:PHE:O	2.04	0.57
1:A:481:LYS:HD2	1:A:489:PHE:CZ	2.40	0.56
1:A:280:HIS:CD2	1:A:282:GLN:HG2	2.46	0.51
1:A:481:LYS:HB3	1:A:489:PHE:CD2	2.45	0.50
1:A:487:GLN:HB3	1:A:489:PHE:CE2	2.49	0.48
1:A:487:GLN:HB3	1:A:489:PHE:HE2	1.78	0.47
1:A:332:LYS:HE3	3:A:724:HOH:O	2.15	0.46
1:A:412:LYS:HA	1:A:488:ILE:HG12	1.98	0.45
1:A:343:GLN:O	1:A:346:GLN:HG3	2.17	0.45
1:A:277:LYS:HD2	1:A:278:PRO:HD3	2.00	0.44
1:A:370:GLU:HG3	1:A:489:PHE:CD1	2.54	0.42
1:A:332:LYS:HD2	3:A:619:HOH:O	2.19	0.42
1:A:500:SER:O	1:A:504:ILE:HG12	2.19	0.42
1:A:447:LYS:HG3	2:A:4:BR:BR	2.75	0.41
1:A:427:ASP:HA	2:A:3:BR:BR	2.75	0.41
1:A:503:THR:O	1:A:507:LEU:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	246/261 (94%)	242 (98%)	4 (2%)	55 34

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	264	LYS
1	A	352	ASN
1	A	481	LYS
1	A	507	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	280	HIS
1	A	294	GLN
1	A	297	ASN
1	A	343	GLN
1	A	352	ASN
1	A	375	GLN
1	A	385	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/267 (93%)	0.80	41 (16%) 4 7	13, 20, 64, 85	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	515	VAL	6.3
1	A	354	ALA	5.1
1	A	519	LEU	5.0
1	A	512	LEU	4.9
1	A	511	ASN	4.5
1	A	518	LYS	4.5
1	A	359	CYS	4.2
1	A	514	LYS	4.1
1	A	350	SER	4.1
1	A	507	LEU	4.0
1	A	496	TRP	3.7
1	A	505	VAL	3.6
1	A	353	LYS	3.4
1	A	516	GLU	3.4
1	A	362	LEU	3.4
1	A	367	TYR	3.3
1	A	504	ILE	3.2
1	A	357	GLN	3.1
1	A	265	LEU	3.0
1	A	358	SER	3.0
1	A	517	GLU	2.8
1	A	503	THR	2.8
1	A	502	LEU	2.7
1	A	363	PHE	2.6
1	A	347	ILE	2.6
1	A	336	ASN	2.6
1	A	278	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	506	LYS	2.4
1	A	482	HIS	2.4
1	A	360	GLU	2.3
1	A	489	PHE	2.3
1	A	355	ASN	2.3
1	A	509	GLU	2.3
1	A	510	GLU	2.3
1	A	499	VAL	2.3
1	A	442	PHE	2.2
1	A	481	LYS	2.2
1	A	424	CYS	2.1
1	A	348	VAL	2.1
1	A	349	GLN	2.0
1	A	260	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BR	A	3	1/1	0.92	0.12	59,59,59,59	0
2	BR	A	1	1/1	0.95	0.09	28,28,28,28	0
2	BR	A	4	1/1	0.96	0.09	49,49,49,49	0
2	BR	A	6	1/1	0.97	0.18	42,42,42,42	0
2	BR	A	5	1/1	0.98	0.17	46,46,46,46	0
2	BR	A	7	1/1	0.98	0.05	28,28,28,28	0
2	BR	A	2	1/1	0.99	0.17	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.