



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 03:08 AM UTC

PDB ID : 3R40 / pdb_00003r40
Title : Crystal Structure of the Fluoroacetate Dehalogenase RPA1163 - Asp110Asn/apo
Authors : Chan, P.W.Y.; Yakunin, A.F.; Edwards, E.A.; Pai, E.F.
Deposited on : 2011-03-16
Resolution : 1.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1182 (1.08-1.04)
Clashscore	190562	1204 (1.08-1.04)
Ramachandran outliers	187476	1175 (1.08-1.04)
Sidechain outliers	187428	1176 (1.08-1.04)
RSRZ outliers	180081	1181 (1.08-1.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
1	B	306	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fluoroacetate dehalogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	S	0	13	0
			2438	1572	428	425	13			
1	B	287	Total	C	N	O	S	0	8	0
			2365	1523	414	415	13			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q6NAM1
A	0	HIS	-	expression tag	UNP Q6NAM1
A	110	ASN	ASP	engineered mutation	UNP Q6NAM1
A	303	GLY	-	expression tag	UNP Q6NAM1
A	304	SER	-	expression tag	UNP Q6NAM1
B	-1	GLY	-	expression tag	UNP Q6NAM1
B	0	HIS	-	expression tag	UNP Q6NAM1
B	110	ASN	ASP	engineered mutation	UNP Q6NAM1
B	303	GLY	-	expression tag	UNP Q6NAM1
B	304	SER	-	expression tag	UNP Q6NAM1

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Cl	0	0
			4	4		
3	B	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	253	Total 253	O 253	0	0
4	B	144	Total 144	O 144	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fluoroacetate dehalogenase



• Molecule 1: Fluoroacetate dehalogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.93Å 79.27Å 85.08Å 90.00° 103.22° 90.00°	Depositor
Resolution (Å)	41.41 – 1.05 41.41 – 1.05	Depositor EDS
% Data completeness (in resolution range)	96.8 (41.41-1.05) 96.8 (41.41-1.05)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 1.05Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.144 , 0.168 0.152 , 0.174	Depositor DCC
R_{free} test set	12237 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	9.5	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 35.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5206	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	13/2516 (0.5%)	1.22	12/3416 (0.4%)
1	B	1.49	13/2441 (0.5%)	1.21	3/3315 (0.1%)
All	All	1.42	26/4957 (0.5%)	1.21	15/6731 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
All	All	0	5

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	79	GLN	C-O	9.39	1.36	1.24
1	B	82	PRO	N-CA	9.14	1.59	1.47
1	B	81	THR	C-O	-9.07	1.15	1.24
1	A	50	VAL	CA-CB	-8.96	1.43	1.54
1	B	74	PRO	CA-C	8.41	1.64	1.52
1	A	247	LEU	CB-CG	-7.94	1.37	1.53
1	B	144[A]	ARG	N-CA	-7.06	1.36	1.46
1	B	144[B]	ARG	N-CA	-7.06	1.36	1.46
1	A	130	LEU	CB-CG	-6.93	1.39	1.53
1	A	242	VAL	CA-CB	6.48	1.64	1.53
1	B	81	THR	CA-C	6.30	1.59	1.52
1	B	81	THR	C-N	6.00	1.40	1.34
1	B	75	GLU	C-O	5.97	1.31	1.24
1	A	232	ILE	CG1-CD1	-5.79	1.29	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	252	GLY	CA-C	5.78	1.60	1.51
1	A	235	GLU	CA-CB	5.77	1.62	1.53
1	B	249	GLY	C-O	5.74	1.29	1.24
1	A	252	GLY	C-N	5.72	1.41	1.33
1	A	187	ARG	CD-NE	5.61	1.54	1.46
1	B	60	VAL	C-O	-5.52	1.18	1.24
1	A	252	GLY	C-O	5.47	1.31	1.23
1	B	54	LEU	N-CA	-5.44	1.39	1.46
1	A	94	GLU	CD-OE1	-5.39	1.15	1.25
1	A	271	VAL	CA-CB	5.29	1.61	1.54
1	A	130	LEU	CA-CB	5.21	1.64	1.53
1	B	74	PRO	N-CA	5.15	1.53	1.47

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	244	MET	CG-SD-CE	-6.29	87.07	100.90
1	A	143	GLN	CA-C-O	-6.00	112.53	119.79
1	A	265	ARG	NE-CZ-NH1	-5.85	115.65	121.50
1	B	40	PHE	CA-C-O	-5.53	115.54	120.19
1	A	153[A]	ILE	O-C-N	5.38	129.30	122.57
1	A	153[B]	ILE	O-C-N	5.38	129.30	122.57
1	A	187	ARG	NE-CZ-NH1	-5.27	116.23	121.50
1	A	153[A]	ILE	CA-C-N	5.19	130.23	121.14
1	A	153[A]	ILE	C-N-CA	5.19	130.23	121.14
1	A	153[B]	ILE	CA-C-N	5.19	130.23	121.14
1	A	153[B]	ILE	C-N-CA	5.19	130.23	121.14
1	A	218	ASP	CA-CB-CG	5.19	117.79	112.60
1	B	220	ARG	NE-CZ-NH2	5.17	123.85	119.20
1	B	276	ILE	N-CA-C	-5.06	101.19	108.48
1	A	86	ARG	NE-CZ-NH2	5.00	123.70	119.20

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	ARG	Sidechain
1	A	154	TYR	Sidechain
1	B	114	ARG	Sidechain
1	B	141	TYR	Sidechain
1	B	154	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2438	0	2361	23	0
1	B	2365	0	2283	19	0
2	A	1	0	0	0	0
3	A	4	0	0	0	0
3	B	1	0	0	0	0
4	A	253	0	0	7	0
4	B	144	0	0	2	0
All	All	5206	0	4644	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144[B]:ARG:O	1:A:149:TYR:CD1	2.13	1.02
1:A:168[B]:ASN:ND2	4:A:366:HOH:O	1.97	0.96
4:A:464:HOH:O	1:B:176[A]:PHE:CE2	2.20	0.95
1:B:4:LEU:HD21	1:B:204:ILE:HD12	1.46	0.94
4:A:464:HOH:O	1:B:176[A]:PHE:HE2	1.57	0.82
1:A:210[A]:MET:HE2	4:A:472:HOH:O	1.86	0.76
1:A:184[B]:SER:HG	1:A:185:TRP:CD1	2.06	0.74
1:A:153[B]:ILE:HG22	1:A:156:TRP:CD1	2.23	0.73
1:A:144[B]:ARG:O	1:A:149:TYR:HD1	1.64	0.72
1:A:144[A]:ARG:HH21	1:A:144[A]:ARG:HG3	1.55	0.71
1:B:145[A]:MET:HE1	1:B:153:ILE:HD11	1.76	0.68
1:B:144[B]:ARG:NE	4:B:416:HOH:O	2.28	0.67
1:A:210[A]:MET:CE	4:A:430:HOH:O	2.48	0.62
1:B:136:LEU:HB2	1:B:141:TYR:CE2	2.35	0.62
1:B:73:MET:HE1	1:B:210:MET:HB3	1.81	0.61
1:A:153[B]:ILE:CG2	1:A:156:TRP:CD1	2.84	0.60
1:A:210[A]:MET:HE2	4:A:430:HOH:O	2.02	0.58
1:B:232:ILE:HD12	4:B:383:HOH:O	2.03	0.57
1:B:136:LEU:HB2	1:B:141:TYR:HE2	1.70	0.55
1:A:152[A]:LYS:O	1:A:152[A]:LYS:HE2	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:145[A]:MET:HE1	1:B:153:ILE:CD1	2.38	0.54
1:A:248:TRP:HB2	1:A:253:ILE:HG12	1.91	0.53
1:A:46[B]:MET:C	1:A:46[B]:MET:SD	2.93	0.51
1:A:263:VAL:HA	1:A:266:LYS:HE2	1.92	0.51
1:B:46[C]:MET:SD	1:B:46[C]:MET:C	2.95	0.50
1:A:144[A]:ARG:HG3	1:A:144[A]:ARG:NH2	2.21	0.50
1:A:139:TYR:CE2	1:A:143:GLN:HG3	2.47	0.49
1:A:153[B]:ILE:O	1:A:154:TYR:C	2.54	0.47
1:A:46[A]:MET:CE	1:A:283:PRO:HG2	2.46	0.46
1:A:141:TYR:CD2	1:A:156:TRP:CE3	3.03	0.46
1:B:51:ALA:N	1:B:52:PRO:CD	2.79	0.46
1:B:130:LEU:O	1:B:244[A]:MET:HA	2.15	0.46
1:B:277:GLU:HB2	1:B:289:GLN:HE22	1.81	0.45
1:A:210[A]:MET:HE1	4:A:430:HOH:O	2.16	0.44
1:B:46[A]:MET:HE2	1:B:47:TRP:CE2	2.53	0.43
1:B:43:THR:O	1:B:46[B]:MET:HG3	2.19	0.42
1:B:130:LEU:CD2	1:B:244[B]:MET:HE3	2.49	0.42
1:B:234:VAL:CG2	1:B:267:TRP:CZ2	3.03	0.42
1:A:49[B]:ARG:NH2	1:A:193:ALA:O	2.45	0.41
1:A:46[B]:MET:HE3	1:A:47:TRP:CE2	2.56	0.41
1:A:144[A]:ARG:NH2	1:A:144[A]:ARG:CG	2.82	0.40
1:B:181:LYS:HD2	1:B:185:TRP:CZ2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/306 (98%)	293 (97%)	8 (3%)	0	100	100
1	B	292/306 (95%)	284 (97%)	8 (3%)	0	100	100
All	All	593/612 (97%)	577 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/242 (102%)	243 (98%)	4 (2%)	55	17
1	B	240/242 (99%)	239 (100%)	1 (0%)	84	64
All	All	487/484 (101%)	482 (99%)	5 (1%)	70	36

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	TRP
1	A	53	LYS
1	A	152[A]	LYS
1	A	152[B]	LYS
1	B	57	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	B	17	ASN
1	B	79	GLN
1	B	289	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	291/306 (95%)	0.20	8 (2%) 56 68	3, 10, 21, 28	15 (5%)
1	B	287/306 (93%)	0.82	36 (12%) 8 9	4, 15, 28, 35	8 (2%)
All	All	578/612 (94%)	0.51	44 (7%) 20 23	3, 12, 26, 35	23 (3%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	5	ALA	7.3
1	A	300	ALA	7.0
1	B	10	GLY	5.1
1	B	15	TRP	4.8
1	A	15	TRP	4.7
1	B	185	TRP	4.7
1	A	149	TYR	4.6
1	A	253	ILE	4.0
1	B	236	ALA	3.7
1	B	149	TYR	3.6
1	B	52	PRO	3.5
1	B	299	SER	3.4
1	B	276	ILE	3.3
1	B	248	TRP	3.3
1	B	261	LEU	3.2
1	B	234	VAL	3.1
1	A	144[A]	ARG	3.1
1	B	80	HIS	3.0
1	B	4	LEU	3.0
1	A	9	PRO	2.9
1	B	9	PRO	2.8
1	B	260	PRO	2.8
1	B	263	VAL	2.8
1	A	252	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	77	ASP	2.7
1	B	295	VAL	2.6
1	B	81	THR	2.4
1	B	78	GLU	2.4
1	B	271	VAL	2.4
1	B	176[A]	PHE	2.3
1	B	74	PRO	2.3
1	B	188	ALA	2.3
1	A	299	SER	2.3
1	B	70	TRP	2.3
1	B	277	GLU	2.2
1	B	267	TRP	2.2
1	B	250	ALA	2.2
1	B	141	TYR	2.2
1	B	288	ASP	2.2
1	B	275	PRO	2.2
1	B	246	ALA	2.1
1	B	298	PHE	2.0
1	B	184	SER	2.0
1	B	49	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	A	305	1/1	0.98	0.12	17,17,17,17	1
3	CL	A	309	1/1	0.99	0.05	13,13,13,13	1
3	CL	B	305	1/1	0.99	0.06	12,12,12,12	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	A	308	1/1	1.00	0.05	15,15,15,15	0
3	CL	A	306	1/1	1.00	0.02	6,6,6,6	1
3	CL	A	307	1/1	1.00	0.02	11,11,11,11	1

6.5 Other polymers [i](#)

There are no such residues in this entry.