



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:59 AM UTC

PDB ID : 6R4J / pdb_00006r4j
Title : Crystal structure of human GFAT-1 G451E in complex with UDP-GlcNAc
Authors : Ruegenberg, S.; Horn, M.; Pichlo, C.; Allmeroth, K.; Baumann, U.; Denzel, M.S.
Deposited on : 2019-03-22
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	687	3% 92% 6%
1	B	687	25% 93% 5%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 21010 atoms, of which 10440 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutamine--fructose-6-phosphate aminotransferase [isomerizing] 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	649	Total	C	H	N	O	S	0	0	0
			10304	3248	5161	895	968	32			
1	B	654	Total	C	H	N	O	S	0	0	0
			10377	3270	5198	899	978	32			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	299A	HIS	-	insertion	UNP Q06210
A	299B	HIS	-	insertion	UNP Q06210
A	299C	HIS	-	insertion	UNP Q06210
A	299D	HIS	-	insertion	UNP Q06210
A	299E	HIS	-	insertion	UNP Q06210
A	299F	HIS	-	insertion	UNP Q06210
A	451	GLU	GLY	engineered mutation	UNP Q06210
B	299A	HIS	-	insertion	UNP Q06210
B	299B	HIS	-	insertion	UNP Q06210
B	299C	HIS	-	insertion	UNP Q06210
B	299D	HIS	-	insertion	UNP Q06210
B	299E	HIS	-	insertion	UNP Q06210
B	299F	HIS	-	insertion	UNP Q06210
B	451	GLU	GLY	engineered mutation	UNP Q06210

- Molecule 2 is URIDINE-DIPHOSPHATE-N-ACETYLGLUCOSAMINE (CCD ID: UD1) (formula: C₁₇H₂₇N₃O₁₇P₂).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			64	17	25	3	17	2		
2	B	1	Total	C	H	N	O	P	0	0
			64	17	25	3	17	2		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GLUCOSE-6-PHOSPHATE (CCD ID: G6Q) (formula: C₆H₁₃O₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	O	P	0	0
			29	6	13	9	1		
4	B	1	Total	C	H	O	P	0	0
			29	6	13	9	1		

- Molecule 5 is GLUTAMIC ACID (CCD ID: GLU) (formula: $C_5H_9NO_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	0	0
			15	5	5	1	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	79	Total 79	O 79	0	0
6	B	47	Total 47	O 47	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamine--fructose-6-phosphate aminotransferase [isomerizing] 1



- Molecule 1: Glutamine--fructose-6-phosphate aminotransferase [isomerizing] 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	153.23Å 153.23Å 162.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.73 – 2.42 48.73 – 2.42	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.73-2.42) 96.4 (48.73-2.42)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.42Å)	Xtriage
Refinement program	PHENIX (dev_2499: ???)	Depositor
R, R_{free}	0.194 , 0.223 0.196 , 0.224	Depositor DCC
R_{free} test set	1991 reflections (2.69%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21010	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, UD1, G6Q

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/5232	0.27	0/7061
1	B	0.10	0/5267	0.25	0/7109
All	All	0.10	0/10499	0.26	0/14170

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5143	5161	5160	11	0
1	B	5179	5198	5197	7	0
2	A	39	25	25	0	0
2	B	39	25	25	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	16	13	11	2	0
4	B	16	13	11	1	0
5	A	10	5	5	1	0
6	A	79	0	0	0	0
6	B	47	0	0	0	0
All	All	10570	10440	10434	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:558:LYS:NZ	4:A:703:G6Q:O1	2.15	0.79
1:A:422:GLN:N	4:A:703:G6Q:O2P	2.27	0.68
1:A:621:ASP:OD1	1:A:622:LYS:N	2.34	0.59
1:B:210:GLY:O	1:B:258:TYR:N	2.42	0.51
1:B:334:GLU:OE1	1:B:334:GLU:N	2.47	0.48
1:A:672:ARG:NH1	1:B:581:ALA:O	2.47	0.47
1:A:29:ARG:NH1	1:A:317:MET:SD	2.88	0.47
1:A:544:GLY:N	1:A:552:CYS:SG	2.89	0.46
1:B:422:GLN:N	4:B:703:G6Q:O3P	2.40	0.46
1:A:332:GLN:N	1:A:333:PRO:CD	2.81	0.43
1:B:419:PHE:CZ	1:B:433:LEU:HA	2.53	0.43
1:A:419:PHE:CZ	1:A:433:LEU:HA	2.53	0.42
1:A:2:CYS:N	5:A:704:GLU:OE1	2.52	0.42
1:B:642:ASP:OD1	1:B:643:CYS:N	2.52	0.42
1:A:385:ARG:NH2	1:B:399:GLU:OE2	2.42	0.41
1:A:554:GLU:OE2	1:A:558:LYS:HE3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	643/687 (94%)	629 (98%)	13 (2%)	1 (0%)	43	57
1	B	648/687 (94%)	628 (97%)	18 (3%)	2 (0%)	36	49
All	All	1291/1374 (94%)	1257 (97%)	31 (2%)	3 (0%)	43	57

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	250	VAL
1	B	677	SER
1	A	250	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	567/602 (94%)	567 (100%)	0	100	100
1	B	572/602 (95%)	571 (100%)	1 (0%)	87	94
All	All	1139/1204 (95%)	1138 (100%)	1 (0%)	88	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	79	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	162	ASN
1	A	215	HIS
1	A	422	GLN
1	A	448	ASN
1	B	106	ASN
1	B	162	ASN
1	B	269	HIS
1	B	422	GLN
1	B	448	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	G6Q	A	703	-	14,15,15	0.44	0	19,21,21	0.89	2 (10%)
5	GLU	A	704	-	8,9,9	1.10	1 (12%)	8,11,11	1.20	1 (12%)
2	UD1	A	701	3	40,41,41	3.36	21 (52%)	59,62,62	1.40	6 (10%)
4	G6Q	B	703	-	14,15,15	0.44	0	19,21,21	0.78	0
2	UD1	B	701	3	40,41,41	3.34	21 (52%)	59,62,62	1.44	7 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	G6Q	A	703	-	-	6/19/20/20	-
5	GLU	A	704	-	-	2/9/9/9	-
2	UD1	A	701	3	-	1/26/63/63	0/3/3/3
4	G6Q	B	703	-	-	6/19/20/20	-
2	UD1	B	701	3	-	7/26/63/63	0/3/3/3

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	UD1	C3B-C2B	-8.30	1.30	1.53
2	A	701	UD1	C3B-C2B	-8.25	1.31	1.53
2	A	701	UD1	C2-N1	7.17	1.49	1.38
2	B	701	UD1	C2-N3	7.14	1.50	1.38
2	B	701	UD1	C2-N1	7.14	1.49	1.38
2	A	701	UD1	C2-N3	7.08	1.50	1.38
2	A	701	UD1	PA-O3A	6.26	1.66	1.59
2	A	701	UD1	PB-O3A	6.04	1.66	1.59
2	B	701	UD1	PA-O3A	5.92	1.65	1.59
2	B	701	UD1	C6-C5	5.70	1.48	1.35
2	B	701	UD1	PB-O3A	5.65	1.65	1.59
2	A	701	UD1	C6-C5	5.64	1.48	1.35
2	A	701	UD1	O4B-C1B	-5.07	1.30	1.42
2	B	701	UD1	O4B-C1B	-5.03	1.30	1.42
2	B	701	UD1	O4B-C4B	4.42	1.54	1.45
2	B	701	UD1	C4-N3	4.40	1.46	1.38
2	A	701	UD1	O4B-C4B	4.40	1.54	1.45
2	A	701	UD1	C4-N3	4.34	1.46	1.38
2	B	701	UD1	C5B-C4B	-3.61	1.40	1.51
2	A	701	UD1	C5B-C4B	-3.55	1.40	1.51
2	B	701	UD1	C7'-N2'	3.42	1.45	1.34
2	A	701	UD1	C7'-N2'	3.32	1.45	1.34
2	B	701	UD1	O5'-C1'	3.32	1.50	1.41
2	A	701	UD1	O5'-C1'	3.28	1.50	1.41
2	A	701	UD1	C6-N1	2.96	1.45	1.38
2	B	701	UD1	C6-N1	2.92	1.45	1.38
2	A	701	UD1	C2B-C1B	2.87	1.62	1.53
2	A	701	UD1	O3B-C3B	2.86	1.50	1.43
2	B	701	UD1	O3B-C3B	2.85	1.50	1.43
2	B	701	UD1	C2B-C1B	2.85	1.62	1.53
2	B	701	UD1	O4-C4	-2.78	1.19	1.24
2	A	701	UD1	O4-C4	-2.70	1.19	1.24
2	B	701	UD1	C2'-N2'	2.49	1.49	1.45
2	B	701	UD1	C5-C4	2.40	1.48	1.43
2	A	701	UD1	C5-C4	2.36	1.48	1.43
2	A	701	UD1	C3'-C2'	-2.33	1.48	1.53
2	A	701	UD1	O2-C2	-2.33	1.18	1.23
2	A	701	UD1	C2'-N2'	2.32	1.49	1.45
2	B	701	UD1	O2-C2	-2.30	1.19	1.23
2	B	701	UD1	C3'-C2'	-2.24	1.49	1.53
5	A	704	GLU	OXT-C	-2.21	1.23	1.30
2	B	701	UD1	PB-O1'	2.13	1.65	1.59
2	A	701	UD1	PB-O1'	2.00	1.65	1.59

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	UD1	C4-N3-C2	-5.49	119.80	126.61
2	A	701	UD1	C4-N3-C2	-5.38	119.94	126.61
2	B	701	UD1	N3-C2-N1	3.62	119.60	114.89
2	B	701	UD1	C5-C4-N3	3.58	119.82	114.80
2	A	701	UD1	C5-C4-N3	3.57	119.80	114.80
2	A	701	UD1	N3-C2-N1	3.53	119.49	114.89
2	A	701	UD1	O4-C4-C5	-3.02	119.96	125.16
2	B	701	UD1	O4-C4-C5	-2.95	120.07	125.16
2	B	701	UD1	C8'-C7'-N2'	2.82	120.80	116.12
5	A	704	GLU	OXT-C-O	-2.70	117.95	124.08
2	A	701	UD1	C8'-C7'-N2'	2.21	119.78	116.12
4	A	703	G6Q	O2-C2-C3	2.17	114.54	109.46
2	B	701	UD1	O2-C2-N1	-2.09	120.08	122.80
2	B	701	UD1	C3B-C2B-C1B	2.07	105.38	101.46
4	A	703	G6Q	O6-P-O1P	2.04	111.95	106.44
2	A	701	UD1	O2-C2-N1	-2.00	120.19	122.80

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	UD1	PA-O3A-PB-O1'
2	B	701	UD1	C5B-O5B-PA-O1A
4	A	703	G6Q	O1-C1-C2-C3
4	A	703	G6Q	O2-C2-C3-C4
4	A	703	G6Q	O2-C2-C3-O3
4	A	703	G6Q	C4-C5-C6-O6
4	B	703	G6Q	O1-C1-C2-C3
4	B	703	G6Q	C1-C2-C3-C4
4	B	703	G6Q	C1-C2-C3-O3
4	B	703	G6Q	O2-C2-C3-C4
4	B	703	G6Q	O2-C2-C3-O3
4	B	703	G6Q	C4-C5-C6-O6
2	B	701	UD1	C8'-C7'-N2'-C2'
2	B	701	UD1	O7'-C7'-N2'-C2'
2	B	701	UD1	C1'-C2'-N2'-C7'
2	B	701	UD1	O5'-C5'-C6'-O6'
4	A	703	G6Q	C1-C2-C3-C4
4	A	703	G6Q	C1-C2-C3-O3
5	A	704	GLU	OE2-CD-CG-CB
2	B	701	UD1	C1'-O1'-PB-O2B

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	704	GLU	OE1-CD-CG-CB
2	B	701	UD1	C3'-C2'-N2'-C7'


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	703	G6Q	2	0
5	A	704	GLU	1	0
4	B	703	G6Q	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand G6Q A 703





Ligand G6Q B 703





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	649/687 (94%)	-0.01	20 (3%)	51	48	36, 58, 110, 167	0
1	B	654/687 (95%)	0.96	170 (25%)	1	1	39, 83, 207, 268	0
All	All	1303/1374 (94%)	0.48	190 (14%)	6	4	36, 64, 189, 268	0

All (190) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	304	ARG	7.0
1	B	227	ARG	6.6
1	B	305	ALA	5.9
1	B	20	LEU	5.5
1	A	227	ARG	5.4
1	B	94	THR	5.0
1	B	155	LEU	5.0
1	B	19	ILE	5.0
1	B	259	PHE	5.0
1	B	680	VAL	4.9
1	A	301	HIS	4.9
1	B	54	ALA	4.8
1	A	294	ILE	4.7
1	B	250	VAL	4.7
1	A	244	THR	4.6
1	B	255	VAL	4.4
1	B	198	ALA	4.3
1	B	120	VAL	4.3
1	B	173	LEU	4.3
1	B	125	ILE	4.1
1	B	170	PHE	4.1
1	B	50	TRP	4.1
1	B	152	ILE	4.1
1	B	124	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	293	ARG	4.0
1	B	309	LEU	3.9
1	B	674	LEU	3.9
1	A	367	ARG	3.9
1	B	282	ALA	3.9
1	B	257	TYR	3.9
1	B	273	VAL	3.9
1	B	64	GLY	3.8
1	B	222	ILE	3.8
1	B	225	LEU	3.8
1	B	24	ILE	3.8
1	B	208	LEU	3.7
1	A	54	ALA	3.7
1	B	48	LYS	3.7
1	B	11	HIS	3.6
1	B	119	ILE	3.6
1	B	291	ILE	3.6
1	B	258	TYR	3.6
1	B	308	THR	3.4
1	B	59	LEU	3.4
1	B	224	ILE	3.4
1	A	50	TRP	3.4
1	B	216	LYS	3.4
1	B	254	ALA	3.3
1	B	221	HIS	3.3
1	B	7	TYR	3.3
1	B	73	VAL	3.3
1	B	244	THR	3.3
1	B	129	TYR	3.3
1	A	671	PRO	3.2
1	B	174	VAL	3.2
1	A	672	ARG	3.2
1	B	81	LEU	3.2
1	B	159	MET	3.2
1	B	226	TYR	3.2
1	B	285	VAL	3.2
1	B	195	PRO	3.1
1	A	33	ARG	3.1
1	B	153	ALA	3.1
1	B	223	PRO	3.1
1	B	274	ILE	3.1
1	B	670	PHE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	103	SER	3.1
1	A	302	PRO	3.0
1	B	284	VAL	3.0
1	B	530	LEU	3.0
1	B	100	GLY	3.0
1	B	283	ALA	3.0
1	A	349	TYR	3.0
1	B	281	VAL	2.9
1	B	78	ASP	2.9
1	B	55	CYS	2.9
1	B	69	LEU	2.9
1	B	49	ASP	2.9
1	B	306	VAL	2.9
1	B	253	LYS	2.8
1	B	90	GLY	2.8
1	B	74	HIS	2.8
1	B	200	GLY	2.8
1	B	169	SER	2.8
1	B	673	ASN	2.8
1	B	127	THR	2.8
1	B	25	LYS	2.8
1	B	135	PHE	2.8
1	B	37	SER	2.7
1	B	102	PRO	2.7
1	B	141	TYR	2.7
1	A	55	CYS	2.7
1	B	98	THR	2.7
1	B	139	LYS	2.7
1	B	245	THR	2.7
1	B	56	LYS	2.7
1	B	134	LYS	2.7
1	B	187	LEU	2.7
1	B	204	GLY	2.7
1	B	91	ILE	2.7
1	A	75	LYS	2.7
1	B	23	LEU	2.7
1	B	156	VAL	2.7
1	B	199	VAL	2.7
1	A	310	GLN	2.7
1	B	310	GLN	2.7
1	B	251	GLU	2.6
1	A	96	TRP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	67	LYS	2.6
1	B	17	ARG	2.6
1	B	132	LEU	2.6
1	B	38	ALA	2.6
1	B	272	ARG	2.6
1	B	367	ARG	2.5
1	B	275	PHE	2.5
1	B	472	VAL	2.5
1	B	292	HIS	2.5
1	B	349	TYR	2.5
1	A	79	MET	2.5
1	B	209	ILE	2.5
1	B	270	THR	2.5
1	B	290	SER	2.5
1	B	26	GLY	2.5
1	B	32	TYR	2.5
1	B	99	HIS	2.5
1	B	87	VAL	2.5
1	B	164	GLU	2.4
1	B	611	ALA	2.4
1	B	123	ASN	2.4
1	B	51	GLU	2.4
1	B	677	SER	2.4
1	B	247	LEU	2.4
1	B	189	PHE	2.4
1	B	22	THR	2.4
1	B	105	VAL	2.4
1	B	271	ASN	2.4
1	B	264	SER	2.3
1	B	86	ASP	2.3
1	B	157	LYS	2.3
1	B	268	GLU	2.3
1	B	671	PRO	2.3
1	B	307	GLN	2.3
1	B	33	ARG	2.3
1	B	44	GLY	2.3
1	B	79	MET	2.3
1	B	118	PHE	2.3
1	B	62	LYS	2.3
1	B	280	ASP	2.3
1	B	13	PRO	2.3
1	A	184	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	201	THR	2.2
1	B	114	LYS	2.2
1	B	681	GLU	2.2
1	B	668	VAL	2.2
1	A	251	GLU	2.2
1	B	42	PHE	2.2
1	B	158	TYR	2.2
1	B	41	GLY	2.2
1	B	146	GLU	2.2
1	B	52	ALA	2.2
1	B	93	HIS	2.2
1	B	89	LEU	2.2
1	B	148	ASP	2.2
1	B	65	LYS	2.2
1	B	177	VAL	2.1
1	B	450	VAL	2.1
1	A	52	ALA	2.1
1	B	46	ASN	2.1
1	B	45	GLY	2.1
1	B	276	LEU	2.1
1	B	112	SER	2.1
1	B	167	ASP	2.1
1	B	12	VAL	2.1
1	B	188	VAL	2.1
1	B	192	VAL	2.1
1	B	68	ALA	2.1
1	B	220	ASP	2.1
1	B	136	LEU	2.1
1	B	217	LEU	2.1
1	B	186	ALA	2.1
1	B	106	ASN	2.1
1	B	95	ARG	2.1
1	B	84	GLU	2.1
1	B	219	THR	2.1
1	B	287	GLY	2.0
1	B	172	THR	2.0
1	B	246	CYS	2.0
1	B	678	VAL	2.0
1	B	10	TYR	2.0
1	B	215	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UD1	B	701	39/39	0.93	0.10	48,72,108,125	0
5	GLU	A	704	10/10	0.93	0.11	76,80,96,96	0
4	G6Q	B	703	16/16	0.94	0.09	40,49,57,63	0
4	G6Q	A	703	16/16	0.95	0.09	34,42,51,56	0
2	UD1	A	701	39/39	0.96	0.07	41,55,72,79	0
3	MG	B	702	1/1	0.97	0.08	40,40,40,40	0
3	MG	A	702	1/1	0.97	0.04	31,31,31,31	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UD1 B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around G6Q B 703:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around G6Q A 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UD1 A 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.