



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:08 AM UTC

PDB ID : 4R5W / pdb_00004r5w
Title : Human artd1 (parp1) - catalytic domain in complex with inhibitor xav939
Authors : Karlberg, T.; Thorsell, A.G.; Schuler, H.
Deposited on : 2014-08-22
Resolution : 2.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1520 (2.86-2.82)
Clashscore	190562	1559 (2.86-2.82)
Ramachandran outliers	187476	1517 (2.86-2.82)
Sidechain outliers	187428	1518 (2.86-2.82)
RSRZ outliers	180081	1521 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	2% 82% 15% ..
1	B	358	5% 81% 17% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	351	Total	C	N	O	S	0	0	0
			2758	1756	468	522	12			
1	B	355	Total	C	N	O	S	0	0	0
			2798	1780	480	526	12			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	661	MET	-	initiating methionine	UNP P09874
A	762	ALA	VAL	variant	UNP P09874
A	1012	ALA	-	expression tag	UNP P09874
A	1013	HIS	-	expression tag	UNP P09874
A	1014	HIS	-	expression tag	UNP P09874
A	1015	HIS	-	expression tag	UNP P09874
A	1016	HIS	-	expression tag	UNP P09874
A	1017	HIS	-	expression tag	UNP P09874
A	1018	HIS	-	expression tag	UNP P09874
B	661	MET	-	initiating methionine	UNP P09874
B	762	ALA	VAL	variant	UNP P09874
B	1012	ALA	-	expression tag	UNP P09874
B	1013	HIS	-	expression tag	UNP P09874
B	1014	HIS	-	expression tag	UNP P09874
B	1015	HIS	-	expression tag	UNP P09874
B	1016	HIS	-	expression tag	UNP P09874
B	1017	HIS	-	expression tag	UNP P09874
B	1018	HIS	-	expression tag	UNP P09874

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 2-[4-(trifluoromethyl)phenyl]-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol (CCD ID: XAV) (formula: C₁₄H₁₁F₃N₂OS).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	0	0
			21	14	3	2	1	1		
3	B	1	Total	C	F	N	O	S	0	0
			21	14	3	2	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		
4	B	6	Total	O	0	0
			6	6		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.53Å 67.66Å 91.17Å 90.00° 111.28° 90.00°	Depositor
Resolution (Å)	48.46 – 2.84 48.46 – 2.84	Depositor EDS
% Data completeness (in resolution range)	99.2 (48.46-2.84) 99.9 (48.46-2.84)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.86Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.192 , 0.245 0.200 , 0.250	Depositor DCC
R_{free} test set	1013 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	46.5	Xtriage
Anisotropy	0.350	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5644	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XAV, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/2810	1.33	14/3790 (0.4%)
1	B	0.87	0/2854	1.35	15/3850 (0.4%)
All	All	0.86	0/5664	1.34	29/7640 (0.4%)

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	979	VAL	CA-C-N	8.92	137.75	121.70
1	A	979	VAL	C-N-CA	8.92	137.75	121.70
1	B	979	VAL	CA-C-N	8.91	137.73	121.70
1	B	979	VAL	C-N-CA	8.91	137.73	121.70
1	A	978	GLY	CA-C-N	6.42	130.14	120.77
1	A	978	GLY	C-N-CA	6.42	130.14	120.77
1	B	978	GLY	CA-C-N	6.38	130.09	120.77
1	B	978	GLY	C-N-CA	6.38	130.09	120.77
1	B	766	ASP	CA-CB-CG	6.19	118.79	112.60
1	A	766	ASP	CA-CB-CG	6.01	118.61	112.60
1	B	899	ASP	CA-CB-CG	5.85	118.45	112.60
1	A	852	LYS	CA-C-N	5.71	128.20	120.44
1	A	852	LYS	C-N-CA	5.71	128.20	120.44
1	B	852	LYS	CA-C-N	5.58	128.03	120.44
1	B	852	LYS	C-N-CA	5.58	128.03	120.44
1	A	899	ASP	CA-CB-CG	5.51	118.11	112.60
1	A	987	ASN	CA-CB-CG	5.30	117.90	112.60
1	B	987	ASN	CA-CB-CG	5.28	117.88	112.60
1	A	738	THR	CA-C-N	5.21	127.53	120.44
1	A	738	THR	C-N-CA	5.21	127.53	120.44
1	B	791	ASP	CA-CB-CG	5.20	117.80	112.60
1	B	738	THR	CA-C-N	5.18	127.48	120.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	738	THR	C-N-CA	5.18	127.48	120.44
1	A	828	ALA	N-CA-C	-5.12	105.93	113.61
1	B	912	GLN	CA-C-N	5.11	125.62	120.00
1	B	912	GLN	C-N-CA	5.11	125.62	120.00
1	B	828	ALA	N-CA-C	-5.07	106.00	113.61
1	A	823	ALA	CA-C-N	5.05	127.99	120.31
1	A	823	ALA	C-N-CA	5.05	127.99	120.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2758	0	2803	21	0
1	B	2798	0	2831	22	0
2	A	25	0	0	0	0
2	B	10	0	0	0	0
3	A	21	0	10	0	0
3	B	21	0	10	1	0
4	A	5	0	0	0	0
4	B	6	0	0	0	0
All	All	5644	0	5654	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:834:ILE:HD11	1:A:1006:LYS:HB2	1.76	0.68
1:B:834:ILE:HD11	1:B:1006:LYS:HB2	1.76	0.67
1:A:717:GLN:HG2	1:A:887:THR:OG1	1.99	0.61
1:B:675:MET:HE1	1:B:1004:LYS:HD2	1.86	0.58
1:A:686:MET:HE3	1:A:693:LEU:CD2	2.37	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:686:MET:HE3	1:B:693:LEU:CD2	2.39	0.53
1:A:686:MET:HE3	1:A:693:LEU:HD21	1.92	0.52
1:B:717:GLN:HG2	1:B:887:THR:OG1	2.10	0.51
1:A:675:MET:HE1	1:A:1004:LYS:HD2	1.93	0.51
1:A:927:GLY:HA3	1:A:946:HIS:ND1	2.26	0.51
1:B:686:MET:HE3	1:B:693:LEU:HD21	1.92	0.51
1:A:696:MET:HB2	1:A:741:PRO:HG2	1.93	0.50
1:A:933:LYS:HE2	1:A:979:VAL:HG13	1.93	0.50
1:B:841:ARG:HD2	1:B:873:LEU:O	2.10	0.50
1:B:933:LYS:HE2	1:B:979:VAL:HG13	1.93	0.50
1:A:841:ARG:HD2	1:A:873:LEU:O	2.12	0.49
1:B:697:PRO:HD2	1:B:700:LYS:HD3	1.96	0.48
1:B:696:MET:HB2	1:B:741:PRO:HG2	1.96	0.48
1:A:865:ARG:HG2	1:A:909:HIS:HB3	1.96	0.48
1:A:780:GLY:HA2	1:B:780:GLY:HA2	1.96	0.47
1:B:916:ILE:HG21	1:B:1004:LYS:HE3	1.97	0.47
1:A:697:PRO:HD2	1:A:700:LYS:HD3	1.97	0.47
1:B:900:MET:HE3	1:B:956:PRO:HG3	1.98	0.46
1:A:821:THR:HB	1:A:900:MET:HA	1.98	0.46
1:B:821:THR:HB	1:B:900:MET:HA	1.99	0.45
1:A:891:PHE:HA	1:A:936:SER:O	2.16	0.44
1:B:891:PHE:HA	1:B:936:SER:O	2.17	0.44
1:A:916:ILE:HG21	1:A:1004:LYS:HE3	1.98	0.44
1:A:900:MET:HE3	1:A:956:PRO:HG3	2.00	0.43
1:A:689:TYR:CG	1:A:764:MET:HG3	2.54	0.42
1:A:826:HIS:CE1	1:A:986:TYR:CE2	3.07	0.42
1:B:872:ILE:HG21	1:B:920:LEU:HD11	2.00	0.42
1:B:863:GLY:O	3:B:1103:XAV:HAE	2.19	0.42
1:A:826:HIS:ND1	1:A:902:SER:HB2	2.34	0.42
1:B:826:HIS:ND1	1:B:902:SER:HB2	2.34	0.42
1:A:872:ILE:HG21	1:A:920:LEU:HD11	2.02	0.42
1:B:831:LEU:HD22	1:B:1005:LEU:HD13	2.03	0.41
1:A:821:THR:HA	1:A:900:MET:HE2	2.03	0.41
1:B:689:TYR:CG	1:B:764:MET:HG3	2.56	0.41
1:B:821:THR:HA	1:B:900:MET:HE2	2.02	0.41
1:B:661:MET:O	1:B:788:ASP:HB2	2.21	0.40
1:B:826:HIS:CE1	1:B:986:TYR:CE2	3.09	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/358 (97%)	340 (98%)	7 (2%)	0	100	100
1	B	351/358 (98%)	343 (98%)	8 (2%)	0	100	100
All	All	698/716 (98%)	683 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/314 (98%)	288 (94%)	19 (6%)	16	34
1	B	311/314 (99%)	287 (92%)	24 (8%)	12	26
All	All	618/628 (98%)	575 (93%)	43 (7%)	14	30

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	661	MET
1	A	674	LYS
1	A	684	LYS
1	A	686	MET
1	A	724	SER
1	A	729	ILE
1	A	732	LEU
1	A	740	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	758	VAL
1	A	763	GLU
1	A	769	LEU
1	A	806	ARG
1	A	900	MET
1	A	939	SER
1	A	961	ASN
1	A	962	ILE
1	A	979	VAL
1	A	985	LEU
1	A	988	GLU
1	B	661	MET
1	B	664	LYS
1	B	674	LYS
1	B	684	LYS
1	B	686	MET
1	B	695	LYS
1	B	724	SER
1	B	729	ILE
1	B	732	LEU
1	B	740	ILE
1	B	758	VAL
1	B	763	GLU
1	B	769	LEU
1	B	806	ARG
1	B	845	CYS
1	B	865	ARG
1	B	900	MET
1	B	903	LYS
1	B	962	ILE
1	B	979	VAL
1	B	985	LEU
1	B	988	GLU
1	B	1010	LYS
1	B	1017	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	820	ASN
1	A	868	ASN
1	A	906	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	998	ASN
1	B	868	ASN
1	B	875	GLN
1	B	906	ASN
1	B	998	ASN
1	B	1014	HIS
1	B	1016	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	XAV	A	1106	-	23,23,23	1.62	4 (17%)	30,34,34	2.61	16 (53%)
2	SO4	A	1101	-	4,4,4	0.40	0	6,6,6	0.20	0
3	XAV	B	1103	-	23,23,23	1.32	3 (13%)	30,34,34	2.17	10 (33%)
2	SO4	A	1102	-	4,4,4	0.26	0	6,6,6	0.32	0
2	SO4	A	1104	-	4,4,4	0.31	0	6,6,6	0.41	0
2	SO4	B	1101	-	4,4,4	0.21	0	6,6,6	0.20	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	1102	-	4,4,4	0.37	0	6,6,6	0.32	0
2	SO4	A	1103	-	4,4,4	0.27	0	6,6,6	0.22	0
2	SO4	A	1105	-	4,4,4	0.27	0	6,6,6	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	XAV	B	1103	-	-	4/10/17/17	0/3/3/3
3	XAV	A	1106	-	-	4/10/17/17	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1106	XAV	CAP-C2	-4.15	1.36	1.48
3	B	1103	XAV	CAP-C2	-3.21	1.39	1.48
3	A	1106	XAV	C5-C4	2.92	1.45	1.40
3	A	1106	XAV	C4-N3	-2.47	1.30	1.34
3	B	1103	XAV	CAU-CAQ	2.13	1.54	1.49
3	A	1106	XAV	FAC-CAU	2.08	1.40	1.33
3	B	1103	XAV	CAI-SAN	2.05	1.86	1.80

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1106	XAV	CAJ-C4-C5	6.34	127.88	121.40
3	A	1106	XAV	CAP-C2-N1	5.38	126.30	117.40
3	B	1103	XAV	CAJ-C4-C5	5.08	126.59	121.40
3	B	1103	XAV	C5-C4-N3	-4.74	119.83	123.48
3	B	1103	XAV	FAB-CAU-CAQ	-4.20	103.90	112.90
3	A	1106	XAV	FAB-CAU-CAQ	-3.83	104.70	112.90
3	A	1106	XAV	CAI-CAJ-C4	-3.70	108.28	113.64
3	B	1103	XAV	C2-N1-C6	3.60	121.47	115.98
3	A	1106	XAV	C5-C4-N3	-3.48	120.80	123.48
3	B	1103	XAV	N3-C2-N1	-3.41	119.52	125.13
3	A	1106	XAV	CAJ-C4-N3	-3.34	111.32	115.85
3	A	1106	XAV	FAD-CAU-CAQ	-3.21	106.02	112.90
3	A	1106	XAV	FAD-CAU-FAC	3.04	116.74	105.77
3	A	1106	XAV	CAP-C2-N3	-3.04	112.38	117.40
3	A	1106	XAV	C2-N1-C6	2.69	120.08	115.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1103	XAV	C2-N3-C4	2.61	121.34	116.82
3	A	1106	XAV	CAJ-CAI-SAN	-2.60	106.46	112.39
3	B	1103	XAV	FAC-CAU-FAB	2.37	114.33	105.77
3	A	1106	XAV	N3-C2-N1	-2.32	121.31	125.13
3	A	1106	XAV	C2-N3-C4	2.11	120.48	116.82
3	A	1106	XAV	FAC-CAU-CAQ	-2.10	108.40	112.90
3	B	1103	XAV	CAP-C2-N1	2.10	120.87	117.40
3	A	1106	XAV	CAH-CAQ-CAG	2.07	121.11	118.03
3	B	1103	XAV	CAE-CAG-CAQ	-2.06	118.32	121.17
3	A	1106	XAV	O6-C6-N1	2.05	122.60	117.24
3	B	1103	XAV	CAE-CAP-C2	2.01	124.08	120.78

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1106	XAV	N1-C2-CAP-CAE
3	A	1106	XAV	N3-C2-CAP-CAE
3	A	1106	XAV	N3-C2-CAP-CAF
3	A	1106	XAV	N1-C2-CAP-CAF
3	B	1103	XAV	N1-C2-CAP-CAF
3	B	1103	XAV	N1-C2-CAP-CAE
3	B	1103	XAV	N3-C2-CAP-CAF
3	B	1103	XAV	N3-C2-CAP-CAE


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1103	XAV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	351/358 (98%)	-0.03	7 (1%) 65 57	25, 43, 82, 117	0
1	B	355/358 (99%)	0.11	18 (5%) 33 26	24, 45, 84, 111	0
All	All	706/716 (98%)	0.04	25 (3%) 47 37	24, 44, 84, 117	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	824	THR	4.9
1	A	826	HIS	4.4
1	B	823	ALA	4.2
1	B	824	THR	4.0
1	B	782	SER	3.7
1	A	823	ALA	3.6
1	A	825	THR	3.6
1	B	825	THR	3.5
1	B	827	ASN	3.0
1	B	964	LEU	2.9
1	B	661	MET	2.8
1	B	855	HIS	2.8
1	A	661	MET	2.7
1	B	1017	HIS	2.7
1	B	745	GLY	2.4
1	B	788	ASP	2.4
1	A	827	ASN	2.3
1	B	744	PHE	2.3
1	B	938	ILE	2.3
1	B	749	PRO	2.2
1	B	828	ALA	2.1
1	B	781	GLY	2.1
1	B	965	ASP	2.1
1	A	782	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	979	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	1105	5/5	0.83	0.18	88,90,93,93	0
2	SO4	A	1104	5/5	0.90	0.10	77,77,77,78	0
2	SO4	B	1102	5/5	0.92	0.10	62,67,68,70	0
2	SO4	A	1103	5/5	0.93	0.10	78,79,80,80	0
3	XAV	B	1103	21/21	0.93	0.11	44,51,72,76	0
3	XAV	A	1106	21/21	0.95	0.09	42,48,63,65	0
2	SO4	A	1102	5/5	0.95	0.09	57,57,59,60	0
2	SO4	B	1101	5/5	0.98	0.05	35,36,38,39	0
2	SO4	A	1101	5/5	0.98	0.06	31,32,34,39	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around XAV B 1103:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around XAV A 1106:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.