



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:38 PM UTC

PDB ID : 6R58 / pdb_00006r58
Title : Crystal structure of PPEP-1(E143A/Y178F/E184A) in complex with substrate peptide Ac-EVNAPVP-CONH2
Authors : Pichlo, C.; Baumann, U.
Deposited on : 2019-03-24
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	2% 93% 5% 2%
1	B	198	4% 92% 5% 1%
1	C	198	6% 95% 3% 1%
1	D	198	5% 97% 2% 1%
2	E	8	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	8	 100%
2	G	8	 100%
2	I	8	 62%  100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12954 atoms, of which 6207 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pro-Pro endopeptidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	192	Total	C	H	N	O	S	0	2	0
			3013	965	1500	259	287	2			
1	B	192	Total	C	H	N	O	S	0	2	0
			3038	968	1519	262	287	2			
1	C	192	Total	C	H	N	O	S	0	1	0
			2989	957	1486	257	287	2			
1	D	198	Total	C	H	N	O	S	0	1	0
			3037	976	1498	263	297	3			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLY	-	expression tag	UNP Q183R7
A	24	SER	-	expression tag	UNP Q183R7
A	25	HIS	-	expression tag	UNP Q183R7
A	26	MET	-	expression tag	UNP Q183R7
A	143	ALA	GLU	engineered mutation	UNP Q183R7
A	178	PHE	TYR	engineered mutation	UNP Q183R7
A	184	ALA	GLU	engineered mutation	UNP Q183R7
B	23	GLY	-	expression tag	UNP Q183R7
B	24	SER	-	expression tag	UNP Q183R7
B	25	HIS	-	expression tag	UNP Q183R7
B	26	MET	-	expression tag	UNP Q183R7
B	143	ALA	GLU	engineered mutation	UNP Q183R7
B	178	PHE	TYR	engineered mutation	UNP Q183R7
B	184	ALA	GLU	engineered mutation	UNP Q183R7
C	23	GLY	-	expression tag	UNP Q183R7
C	24	SER	-	expression tag	UNP Q183R7
C	25	HIS	-	expression tag	UNP Q183R7
C	26	MET	-	expression tag	UNP Q183R7
C	143	ALA	GLU	engineered mutation	UNP Q183R7
C	178	PHE	TYR	engineered mutation	UNP Q183R7
C	184	ALA	GLU	engineered mutation	UNP Q183R7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	23	GLY	-	expression tag	UNP Q183R7
D	24	SER	-	expression tag	UNP Q183R7
D	25	HIS	-	expression tag	UNP Q183R7
D	26	MET	-	expression tag	UNP Q183R7
D	143	ALA	GLU	engineered mutation	UNP Q183R7
D	178	PHE	TYR	engineered mutation	UNP Q183R7
D	184	ALA	GLU	engineered mutation	UNP Q183R7

- Molecule 2 is a protein called ACE-GLU-VAL-ASN-ALA-PRO-VAL-LPD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	8	Total	C	H	N	O	0	0	0
			105	34	51	9	11			
2	F	8	Total	C	H	N	O	0	0	0
			105	34	51	9	11			
2	I	8	Total	C	H	N	O	0	0	0
			105	34	51	9	11			
2	G	8	Total	C	H	N	O	0	0	0
			105	34	51	9	11			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	Ni	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	131	Total	O	0	0
			131	131		
6	B	106	Total	O	0	0
			106	106		
6	C	112	Total	O	0	0
			112	112		
6	D	85	Total	O	0	0
			85	85		
6	E	6	Total	O	0	0
			6	6		
6	F	4	Total	O	0	0
			4	4		
6	I	2	Total	O	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pro-Pro endopeptidase



- Molecule 1: Pro-Pro endopeptidase



- Molecule 1: Pro-Pro endopeptidase



- Molecule 1: Pro-Pro endopeptidase



- Molecule 2: ACE-GLU-VAL-ASN-ALA-PRO-VAL-LPD



There are no outlier residues recorded for this chain.


- Molecule 2: ACE-GLU-VAL-ASN-ALA-PRO-VAL-LPD



There are no outlier residues recorded for this chain.

- Molecule 2: ACE-GLU-VAL-ASN-ALA-PRO-VAL-LPD

Chain I:  62%
100%



- Molecule 2: ACE-GLU-VAL-ASN-ALA-PRO-VAL-LPD

Chain G:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.99Å 73.32Å 237.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.12 – 1.90 46.12 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.12-1.90) 97.4 (46.12-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 1.90Å)	Xtrriage
Refinement program	PHENIX (dev_3409: ???)	Depositor
R, R_{free}	0.190 , 0.214 0.190 , 0.216	Depositor DCC
R_{free} test set	1994 reflections (3.30%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtrriage
Anisotropy	0.366	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12954	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.2946e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, LPD, ACE, ZN, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/1543	0.33	0/2084
1	B	0.17	0/1549	0.31	0/2091
1	C	0.14	0/1533	0.32	0/2073
1	D	0.12	0/1570	0.31	0/2123
2	E	0.13	0/44	0.28	0/61
2	F	0.15	0/44	0.29	0/61
2	G	0.12	0/44	0.27	0/61
2	I	0.13	0/44	0.23	0/61
All	All	0.15	0/6371	0.32	0/8615

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1513	1500	1504	6	0
1	B	1519	1519	1515	7	2
1	C	1503	1486	1485	2	0
1	D	1539	1498	1504	4	2
2	E	54	51	54	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	54	51	54	0	0
2	G	54	51	54	0	0
2	I	54	51	54	0	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	B	5	0	0	0	0
5	D	1	0	0	0	0
6	A	131	0	0	1	0
6	B	106	0	0	5	0
6	C	112	0	0	0	0
6	D	85	0	0	1	0
6	E	6	0	0	0	0
6	F	4	0	0	0	0
6	G	2	0	0	0	0
6	I	2	0	0	0	0
All	All	6747	6207	6224	19	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:TYR:OH	6:B:401:HOH:O	1.82	0.81
1:B:120:THR:HB	6:B:401:HOH:O	1.81	0.77
1:B:80:LYS:NZ	6:B:401:HOH:O	2.21	0.67
1:D:27:ASP:OD1	1:D:28:SER:N	2.30	0.64
1:A:29:THR:OG1	1:A:34:ASN:OD1	2.17	0.60
1:A:130:LYS:NZ	6:A:401:HOH:O	2.35	0.58
1:C:41:ILE:HD11	1:C:73:LYS:HB2	1.88	0.56
1:D:35:LYS:HE3	6:D:411:HOH:O	2.09	0.52
1:B:130:LYS:HE2	1:B:135:ASP:OD1	2.11	0.50
1:A:219:ALA:O	1:A:220:LYS:HB3	2.12	0.49
1:B:96:LYS:NZ	6:B:408:HOH:O	2.47	0.47
1:B:116:LEU:C	1:B:116:LEU:HD23	2.41	0.46
1:D:116:LEU:HD23	1:D:116:LEU:C	2.41	0.45
1:B:80:LYS:CE	6:B:401:HOH:O	2.61	0.44
1:C:116:LEU:HD23	1:C:116:LEU:C	2.43	0.44
1:D:77:LEU:HD13	1:D:77:LEU:C	2.43	0.44
1:A:116:LEU:C	1:A:116:LEU:HD23	2.44	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:PRO:HG2	1:A:49:TYR:CD2	2.55	0.41
1:A:219:ALA:O	1:A:220:LYS:CB	2.69	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:ASN:OD1	1:D:40:GLN:NE2[3_655]	2.08	0.12
1:B:48:ASN:OD1	1:D:40:GLN:HE22[3_655]	1.57	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/198 (97%)	189 (98%)	3 (2%)	0	100	100
1	B	192/198 (97%)	188 (98%)	4 (2%)	0	100	100
1	C	191/198 (96%)	187 (98%)	4 (2%)	0	100	100
1	D	197/198 (100%)	193 (98%)	4 (2%)	0	100	100
2	E	6/8 (75%)	6 (100%)	0	0	100	100
2	F	6/8 (75%)	6 (100%)	0	0	100	100
2	G	6/8 (75%)	6 (100%)	0	0	100	100
2	I	6/8 (75%)	6 (100%)	0	0	100	100
All	All	796/824 (97%)	781 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/164 (98%)	160 (100%)	0	100	100
1	B	161/164 (98%)	160 (99%)	1 (1%)	78	81
1	C	159/164 (97%)	159 (100%)	0	100	100
1	D	163/164 (99%)	163 (100%)	0	100	100
2	E	5/5 (100%)	5 (100%)	0	100	100
2	F	5/5 (100%)	5 (100%)	0	100	100
2	G	5/5 (100%)	5 (100%)	0	100	100
2	I	5/5 (100%)	5 (100%)	0	100	100
All	All	663/676 (98%)	662 (100%)	1 (0%)	87	90

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	73	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	ASN
1	A	198	GLN
1	B	198	GLN
1	C	217	ASN
1	D	34	ASN
1	D	198	GLN
1	D	210	GLN
1	D	217	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LPD	G	7	2	8,8,8	0.83	0	10,10,10	0.73	0
2	LPD	F	7	2	8,8,8	0.80	0	10,10,10	0.80	0
2	LPD	I	7	2	8,8,8	0.81	0	10,10,10	0.78	0
2	LPD	E	7	2	8,8,8	0.82	0	10,10,10	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LPD	G	7	2	-	0/4/11/11	0/1/1/1
2	LPD	F	7	2	-	0/4/11/11	0/1/1/1
2	LPD	I	7	2	-	0/4/11/11	0/1/1/1
2	LPD	E	7	2	-	0/4/11/11	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	B	301	-	4,4,4	0.97	0	6,6,6	0.45	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/198 (96%)	-0.09	4 (2%) 63 67	16, 28, 50, 79	2 (1%)
1	B	192/198 (96%)	0.02	7 (3%) 46 49	18, 31, 56, 90	2 (1%)
1	C	192/198 (96%)	0.12	11 (5%) 29 31	13, 32, 61, 86	1 (0%)
1	D	198/198 (100%)	0.14	10 (5%) 33 36	18, 35, 68, 102	1 (0%)
2	E	6/8 (75%)	0.34	0 100 100	31, 33, 44, 74	0
2	F	6/8 (75%)	0.26	0 100 100	29, 32, 37, 61	0
2	G	6/8 (75%)	0.19	0 100 100	29, 31, 35, 66	0
2	I	6/8 (75%)	2.91	5 (83%) 0 0	63, 66, 78, 90	0
All	All	798/824 (96%)	0.07	37 (4%) 37 40	13, 32, 61, 102	6 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	29	THR	4.7
1	D	23	GLY	4.5
1	B	29	THR	4.5
1	C	29	THR	4.5
2	I	6	VAL	4.0
1	B	47	GLY	3.8
2	I	2	VAL	3.6
2	I	4	ALA	3.5
1	A	220	LYS	3.2
1	D	26	MET	3.2
1	D	24	SER	3.1
1	C	101	LYS	3.1
1	D	28	SER	2.9
1	C	106	THR	2.8
1	B	83	SER	2.8
1	C	99	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	107	GLY	2.7
1	D	220	LYS	2.7
1	A	30	THR	2.7
1	B	32	GLN	2.6
1	B	48	ASN	2.6
1	C	103	TRP	2.5
1	D	46	THR	2.4
1	C	102	GLY	2.4
1	D	31	ILE	2.3
2	I	5	PRO	2.3
1	A	163[A]	LYS	2.2
1	D	40	GLN	2.2
1	D	86	LEU	2.2
2	I	1	GLU	2.1
1	C	108	LYS	2.1
1	C	213	SER	2.1
1	B	86	LEU	2.1
1	C	36	ASP	2.1
1	D	32	GLN	2.0
1	B	154	ASN	2.0
1	C	105	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	LPD	I	7	8/8	0.76	0.20	64,73,87,87	0
2	LPD	E	7	8/8	0.94	0.07	29,33,40,40	0
2	LPD	G	7	8/8	0.94	0.08	32,37,45,45	0
2	LPD	F	7	8/8	0.95	0.07	28,34,42,42	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	D	301	1/1	0.67	0.20	11,11,11,11	1
3	ZN	A	300	1/1	0.71	0.13	12,12,12,12	1
4	PO4	B	301	5/5	0.76	0.14	94,94,94,95	0
5	NI	D	302	1/1	0.88	0.12	62,62,62,62	1
3	ZN	C	300	1/1	1.00	0.05	15,15,15,15	1

6.5 Other polymers [i](#)

There are no such residues in this entry.