



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:45 PM UTC

PDB ID : 4R6V / pdb_00004r6v
Title : Crystal Structure of FGF Receptor (FGFR) 4 Kinase Harboring the V550L Gate-Keeper Mutation in Complex with FIIN-3, an Irreversible Tyrosine Kinase Inhibitor Capable of Overcoming FGFR kinase Gate-Keeper Mutations
Authors : Huang, Z.; Mohammadi, M.
Deposited on : 2014-08-26
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2227 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fibroblast growth factor receptor 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	0	0
			2126	1370	362	378	16			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	431	MET	-	expression tag	UNP P22455
A	432	GLY	-	expression tag	UNP P22455
A	433	SER	-	expression tag	UNP P22455
A	434	SER	-	expression tag	UNP P22455
A	435	HIS	-	expression tag	UNP P22455
A	436	HIS	-	expression tag	UNP P22455
A	437	HIS	-	expression tag	UNP P22455
A	438	HIS	-	expression tag	UNP P22455
A	439	HIS	-	expression tag	UNP P22455
A	440	HIS	-	expression tag	UNP P22455
A	441	SER	-	expression tag	UNP P22455
A	442	GLN	-	expression tag	UNP P22455
A	443	ASP	-	expression tag	UNP P22455
A	444	PRO	-	expression tag	UNP P22455
A	550	LEU	VAL	engineered mutation	UNP P22455
A	664	GLU	ARG	engineered mutation	UNP P22455

- Molecule 2 is N-[4-({[(2,6-dichloro-3,5-dimethoxyphenyl)carbamoyl](6-{[4-(4-methylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)amino}methyl)phenyl]propanamide (CCD ID: FI3) (formula: C₃₄H₃₈Cl₂N₈O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			48	34	2	8	4		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 5	O 4	S 1	0	0
3	A	1	Total 5	O 4	S 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	43	Total 43	O 43	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fibroblast growth factor receptor 4



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	139.38Å 139.38Å 50.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.54 – 2.35 38.54 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.54-2.35) 95.4 (38.54-2.35)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.37Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.174 , 0.227 0.180 , 0.236	Depositor DCC
R_{free} test set	1504 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	26.9	Xtriage
Anisotropy	0.258	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.029 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2227	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FI3, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/2176	0.87	4/2953 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	443	ASP	CA-C-N	9.20	131.33	119.84
1	A	443	ASP	C-N-CA	9.20	131.33	119.84
1	A	582	GLY	CA-C-N	5.60	125.27	119.05
1	A	582	GLY	C-N-CA	5.60	125.27	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2126	0	2133	18	0
2	A	48	0	35	8	0
3	A	10	0	0	1	0
4	A	43	0	0	2	0
All	All	2227	0	2168	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:802:SO4:O3	4:A:930:HOH:O	2.11	0.68
1:A:446:LEU:O	4:A:901:HOH:O	2.16	0.61
2:A:801:FI3:OAF	2:A:801:FI3:H9	2.05	0.57
1:A:487:PHE:CE2	1:A:497:ALA:HB2	2.40	0.56
2:A:801:FI3:CAQ	2:A:801:FI3:H21	2.36	0.56
1:A:469:LEU:HD13	1:A:504:MET:HE1	1.88	0.56
1:A:683:PHE:CZ	1:A:718:LEU:HD13	2.42	0.55
1:A:653:VAL:HG13	1:A:661:LEU:HD21	1.90	0.52
2:A:801:FI3:CAQ	2:A:801:FI3:C5	2.84	0.51
1:A:520:GLU:OE1	2:A:801:FI3:H20	2.13	0.48
1:A:523:VAL:HG11	1:A:633:LEU:HD22	1.96	0.48
1:A:471:LYS:HG3	1:A:472:PRO:HD2	1.95	0.47
1:A:550:LEU:HD21	2:A:801:FI3:CLI	2.52	0.46
2:A:801:FI3:N3	2:A:801:FI3:NBG	2.60	0.44
1:A:553:ALA:O	2:A:801:FI3:NBH	2.48	0.44
1:A:586:PHE:HB3	1:A:587:PRO:HD3	1.99	0.44
1:A:516:ASP:O	1:A:520:GLU:HG3	2.17	0.43
1:A:721:LEU:HD23	1:A:721:LEU:HA	1.82	0.42
1:A:586:PHE:HE1	1:A:718:LEU:HD12	1.83	0.42
1:A:653:VAL:HG11	1:A:695:VAL:HG13	2.02	0.41
1:A:662:PHE:HB2	1:A:699:PHE:CZ	2.56	0.41
1:A:684:THR:HG22	1:A:711:PRO:HB3	2.02	0.40
2:A:801:FI3:C5	2:A:801:FI3:H24	2.51	0.40
1:A:729:ALA:HA	1:A:730:PRO:HD3	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/323 (82%)	259 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/276 (83%)	224 (98%)	4 (2%)	51	66

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	482	VAL
1	A	527	ILE
1	A	550	LEU
1	A	674	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FI3	A	801	1	52,52,52	2.60	22 (42%)	66,72,72	2.65	28 (42%)
3	SO4	A	803	-	4,4,4	0.24	0	6,6,6	0.09	0
3	SO4	A	802	-	4,4,4	0.23	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FI3	A	801	1	-	2/34/44/44	0/5/5/5

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	FI3	CAZ-NCA	-7.57	1.33	1.46
2	A	801	FI3	CBA-NCA	-6.96	1.34	1.46
2	A	801	FI3	CBL-NBF	6.17	1.48	1.35
2	A	801	FI3	CAY-NBZ	-5.32	1.33	1.46
2	A	801	FI3	CAX-NBZ	-4.78	1.35	1.46
2	A	801	FI3	OAG-CBM	-3.92	1.16	1.23
2	A	801	FI3	CAS-CBR	-3.76	1.32	1.39
2	A	801	FI3	CAC-NBZ	-3.23	1.39	1.46
2	A	801	FI3	OBJ-CBW	3.04	1.42	1.37
2	A	801	FI3	CBM-NBG	2.88	1.42	1.37
2	A	801	FI3	CAX-CAZ	-2.83	1.41	1.51
2	A	801	FI3	CAP-CBP	-2.67	1.34	1.39
2	A	801	FI3	CBA-CAY	-2.67	1.41	1.51
2	A	801	FI3	CBT-CLH	-2.62	1.67	1.72
2	A	801	FI3	OBI-CAA	-2.57	1.35	1.42
2	A	801	FI3	CBR-NCA	2.45	1.45	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	FI3	CAN-CBO	-2.38	1.35	1.39
2	A	801	FI3	CBM-NCB	2.28	1.44	1.39
2	A	801	FI3	CBV-CBT	-2.15	1.37	1.40
2	A	801	FI3	OBI-CBV	2.14	1.40	1.37
2	A	801	FI3	C6-N1	-2.10	1.30	1.34
2	A	801	FI3	CBW-CBU	-2.10	1.37	1.40

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	FI3	CAZ-CAX-NBZ	6.71	121.64	110.86
2	A	801	FI3	CAY-NBZ-CAX	6.39	119.71	109.54
2	A	801	FI3	CBA-CAY-NBZ	5.39	119.52	110.86
2	A	801	FI3	CAC-NBZ-CAX	5.33	120.78	110.63
2	A	801	FI3	CBC-CBN-CAL	-4.80	111.91	120.75
2	A	801	FI3	CBN-CBC-NCB	4.78	121.23	113.46
2	A	801	FI3	CAC-NBZ-CAY	4.65	119.50	110.63
2	A	801	FI3	CBA-NCA-CAZ	4.45	121.58	111.57
2	A	801	FI3	CAY-CBA-NCA	4.24	119.72	110.78
2	A	801	FI3	NBG-CBM-NCB	4.13	120.13	115.94
2	A	801	FI3	OBI-CBV-CBT	4.08	119.91	115.44
2	A	801	FI3	CAS-CBR-NCA	-4.00	115.83	121.39
2	A	801	FI3	C6-C5-C4	3.46	119.41	116.64
2	A	801	FI3	CAX-CAZ-NCA	3.34	117.81	110.78
2	A	801	FI3	C5-C6-N1	-3.32	118.56	122.92
2	A	801	FI3	N3-C4-NCB	3.27	120.41	114.77
2	A	801	FI3	CAL-CBN-CAM	3.03	122.73	118.23
2	A	801	FI3	C2-N3-C4	2.97	117.45	114.95
2	A	801	FI3	CBP-NBH-C6	-2.97	121.12	128.75
2	A	801	FI3	CBV-CBT-CLH	2.82	125.47	120.27
2	A	801	FI3	CAU-CBV-CBT	-2.70	117.75	120.99
2	A	801	FI3	CBX-CBT-CLH	-2.68	112.83	118.24
2	A	801	FI3	CBX-CBU-CLI	2.63	123.54	118.24
2	A	801	FI3	CAO-CAM-CBN	-2.53	117.67	121.00
2	A	801	FI3	CBC-CBN-CAM	2.50	125.35	120.75
2	A	801	FI3	OBJ-CBW-CBU	2.47	118.15	115.44
2	A	801	FI3	CBW-CBU-CLI	-2.22	116.17	120.27
2	A	801	FI3	C5-C4-N3	-2.08	119.35	122.82

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	FI3	CBT-CBV-OBI-CAA
2	A	801	FI3	CAU-CBV-OBI-CAA

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	FI3	8	0
3	A	802	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/323 (84%)	0.16	24 (8%) 15 18	14, 28, 59, 84	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	495	ASP	4.4
1	A	662	PHE	4.0
1	A	476	GLY	4.0
1	A	652	PRO	3.9
1	A	750	VAL	3.5
1	A	450	VAL	3.2
1	A	449	LEU	3.2
1	A	497	ALA	2.9
1	A	582	GLY	2.8
1	A	749	ALA	2.6
1	A	478	PHE	2.5
1	A	496	GLN	2.5
1	A	713	HIS	2.4
1	A	634	ALA	2.3
1	A	489	MET	2.3
1	A	471	LYS	2.3
1	A	477	CYS	2.2
1	A	543	GLU	2.2
1	A	447	ALA	2.1
1	A	451	SER	2.1
1	A	712	PRO	2.1
1	A	475	GLU	2.0
1	A	443	ASP	2.0
1	A	479	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	803	5/5	0.60	0.16	124,124,125,125	0
2	FI3	A	801	48/48	0.90	0.13	32,40,59,61	0
3	SO4	A	802	5/5	0.93	0.09	57,58,59,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FI3 A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.