



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:42 AM UTC

PDB ID : 8R6L / pdb_00008r6l
Title : Crystal structure of Candida glabrata Bdf1 bromodomain 2 in the unbound state
Authors : Petosa, C.; Wei, K.; McKenna, C.E.; Govin, J.
Deposited on : 2023-11-22
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	118	3% 92% 5% ..
1	B	118	2% 96% ..
1	C	118	3% 95% ..
1	D	118	4% 88% 6% 6%
1	E	118	3% 93% 5% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	118	5% 92% 6%
1	G	118	5% 91% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Candida glabrata strain CBS138 chromosome C complete sequence.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	0	0
			950	609	156	176	9			
1	B	117	Total	C	N	O	S	0	0	0
			959	614	157	179	9			
1	C	116	Total	C	N	O	S	0	0	0
			950	609	156	176	9			
1	D	111	Total	C	N	O	S	0	0	0
			909	582	151	167	9			
1	E	116	Total	C	N	O	S	0	0	0
			950	609	156	176	9			
1	F	116	Total	C	N	O	S	0	0	0
			950	609	156	176	9			
1	G	116	Total	C	N	O	S	0	0	0
			950	609	156	176	9			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	301	ALA	-	expression tag	UNP Q6FWV7
A	302	MET	-	expression tag	UNP Q6FWV7
A	303	GLY	-	expression tag	UNP Q6FWV7
B	301	ALA	-	expression tag	UNP Q6FWV7
B	302	MET	-	expression tag	UNP Q6FWV7
B	303	GLY	-	expression tag	UNP Q6FWV7
C	301	ALA	-	expression tag	UNP Q6FWV7
C	302	MET	-	expression tag	UNP Q6FWV7
C	303	GLY	-	expression tag	UNP Q6FWV7
D	301	ALA	-	expression tag	UNP Q6FWV7
D	302	MET	-	expression tag	UNP Q6FWV7
D	303	GLY	-	expression tag	UNP Q6FWV7
E	301	ALA	-	expression tag	UNP Q6FWV7
E	302	MET	-	expression tag	UNP Q6FWV7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	303	GLY	-	expression tag	UNP Q6FWV7
F	301	ALA	-	expression tag	UNP Q6FWV7
F	302	MET	-	expression tag	UNP Q6FWV7
F	303	GLY	-	expression tag	UNP Q6FWV7
G	301	ALA	-	expression tag	UNP Q6FWV7
G	302	MET	-	expression tag	UNP Q6FWV7
G	303	GLY	-	expression tag	UNP Q6FWV7

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	6	2		
2	A	1	Total	C	O	0	0
			8	6	2		
2	B	1	Total	C	O	0	0
			8	6	2		
2	B	1	Total	C	O	0	0
			8	6	2		
2	C	1	Total	C	O	0	0
			8	6	2		
2	D	1	Total	C	O	0	0
			8	6	2		
2	D	1	Total	C	O	0	0
			8	6	2		
2	E	1	Total	C	O	0	0
			8	6	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	63	Total	O	0	0
			63	63		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	38	Total 38	O 38	0	0
5	C	30	Total 30	O 30	0	0
5	D	25	Total 25	O 25	0	0
5	E	22	Total 22	O 22	0	0
5	G	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence




- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence




- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence




- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence





- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence



- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence



- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.29Å 67.69Å 96.62Å 90.00° 95.52° 90.00°	Depositor
Resolution (Å)	44.37 – 1.96 44.37 – 1.96	Depositor EDS
% Data completeness (in resolution range)	98.9 (44.37-1.96) 99.0 (44.37-1.96)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.68 (at 1.73Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.200 , 0.227 0.200 , 0.228	Depositor DCC
R_{free} test set	4347 reflections (4.06%)	wwPDB-VP
Wilson B-factor (Å ²)	40.5	Xtriage
Anisotropy	0.392	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6973	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, ACT, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/976	0.69	0/1322
1	B	0.57	0/985	0.62	0/1334
1	C	0.37	0/976	0.55	0/1322
1	D	0.43	0/934	0.53	0/1264
1	E	0.31	0/976	0.46	0/1322
1	F	0.20	0/976	0.37	0/1322
1	G	0.21	0/976	0.40	0/1322
All	All	0.44	0/6799	0.53	0/9208

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	950	0	909	4	0
1	B	959	0	915	2	0
1	C	950	0	909	2	0
1	D	909	0	871	4	0
1	E	950	0	909	5	0
1	F	950	0	909	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	950	0	909	6	0
2	A	16	0	28	0	0
2	B	16	0	28	0	0
2	C	8	0	14	0	0
2	D	16	0	28	2	0
2	E	8	0	14	1	0
3	A	30	0	39	0	0
3	B	18	0	23	0	0
3	C	12	0	16	0	0
3	D	6	0	8	0	0
3	E	18	0	24	1	0
4	A	8	0	6	1	0
4	B	8	0	6	0	0
4	C	4	0	3	0	0
4	E	4	0	3	0	0
4	F	4	0	3	0	0
5	A	63	0	0	0	0
5	B	38	0	0	0	0
5	C	30	0	0	0	0
5	D	25	0	0	0	0
5	E	22	0	0	0	0
5	G	1	0	0	1	0
All	All	6973	0	6574	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:398:ARG:HH11	2:D:902:MPD:H31	1.43	0.81
1:D:398:ARG:NH1	2:D:902:MPD:H31	2.14	0.61
1:G:412:LEU:O	5:G:501:HOH:O	2.18	0.56
1:A:348:LYS:NZ	4:A:904:ACT:OXT	2.39	0.56
1:E:310:CYS:HB3	1:E:360:LEU:HD21	1.91	0.51
1:E:338:LEU:HD13	2:E:901:MPD:HM2	1.95	0.48
1:E:377:LEU:HD11	3:E:903:GOL:H12	1.96	0.47
1:A:310:CYS:HB3	1:A:360:LEU:HD21	1.96	0.46
1:B:333:VAL:HG13	1:B:338:LEU:HD12	1.98	0.45
1:G:322:TYR:HB3	1:G:325:PHE:CZ	2.53	0.44
1:A:349:GLU:HG2	1:A:377:LEU:HD21	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:369:ASP:OD1	1:C:410:ARG:NH1	2.52	0.43
1:F:390:THR:HG22	1:G:336:VAL:O	2.19	0.43
1:C:327:TYR:CG	1:C:328:PRO:HD3	2.53	0.43
1:G:327:TYR:CG	1:G:328:PRO:HD3	2.54	0.43
1:D:333:VAL:HG13	1:D:338:LEU:HD12	2.00	0.42
1:F:304:GLN:OE1	1:F:308:LYS:HE3	2.19	0.42
1:B:322:TYR:CE1	1:B:402:ILE:HD13	2.55	0.42
1:A:372:GLU:HB2	1:A:407:TRP:CZ2	2.54	0.41
1:E:376:ARG:HD3	1:E:376:ARG:HA	1.90	0.41
1:F:327:TYR:CG	1:F:328:PRO:HD3	2.55	0.41
1:G:305:ARG:HB3	1:G:415:ASP:CG	2.47	0.40
1:E:348:LYS:HA	1:E:348:LYS:HD3	1.93	0.40
1:D:314:VAL:O	1:D:318:MET:HG3	2.22	0.40
1:F:391:ILE:HD12	1:G:338:LEU:HD23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/118 (97%)	113 (99%)	1 (1%)	0	100	100
1	B	115/118 (98%)	113 (98%)	2 (2%)	0	100	100
1	C	114/118 (97%)	112 (98%)	2 (2%)	0	100	100
1	D	109/118 (92%)	108 (99%)	1 (1%)	0	100	100
1	E	114/118 (97%)	112 (98%)	2 (2%)	0	100	100
1	F	114/118 (97%)	113 (99%)	1 (1%)	0	100	100
1	G	114/118 (97%)	113 (99%)	1 (1%)	0	100	100
All	All	794/826 (96%)	784 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/106 (98%)	101 (97%)	3 (3%)	37	29
1	B	105/106 (99%)	105 (100%)	0	100	100
1	C	104/106 (98%)	104 (100%)	0	100	100
1	D	99/106 (93%)	97 (98%)	2 (2%)	48	43
1	E	104/106 (98%)	104 (100%)	0	100	100
1	F	104/106 (98%)	103 (99%)	1 (1%)	68	67
1	G	104/106 (98%)	104 (100%)	0	100	100
All	All	724/742 (98%)	718 (99%)	6 (1%)	73	74

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	321	LYS
1	A	360	LEU
1	A	377	LEU
1	D	377	LEU
1	D	388	ASP
1	F	374	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	373	HIS
1	A	386	ASN
1	B	373	HIS
1	C	362	ASN
1	C	386	ASN
1	D	380	GLN
1	D	386	ASN
1	D	393	ASN
1	F	362	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	381	ASN
1	F	397	HIS
1	G	362	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MPD	E	901	-	7,7,7	0.40	0	9,10,10	0.40	0
4	ACT	B	507	-	3,3,3	1.62	1 (33%)	3,3,3	1.43	1 (33%)
3	GOL	B	506	-	5,5,5	0.94	0	5,5,5	1.14	1 (20%)
3	GOL	E	903	-	5,5,5	1.22	1 (20%)	5,5,5	1.26	0
2	MPD	A	902	-	7,7,7	0.29	0	9,10,10	0.69	0
3	GOL	A	909	-	5,5,5	1.19	1 (20%)	5,5,5	0.83	0
2	MPD	D	902	-	7,7,7	0.38	0	9,10,10	0.61	0
4	ACT	F	501	-	3,3,3	1.37	0	3,3,3	1.50	1 (33%)
4	ACT	A	904	-	3,3,3	2.25	1 (33%)	3,3,3	1.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MPD	C	901	-	7,7,7	0.32	0	9,10,10	0.24	0
2	MPD	A	901	-	7,7,7	0.40	0	9,10,10	0.37	0
3	GOL	A	905	-	5,5,5	1.21	1 (20%)	5,5,5	1.19	0
3	GOL	A	906	-	5,5,5	1.35	1 (20%)	5,5,5	1.30	1 (20%)
3	GOL	E	904	-	5,5,5	1.02	0	5,5,5	0.99	0
4	ACT	B	504	-	3,3,3	1.43	1 (33%)	3,3,3	1.38	0
2	MPD	B	503	-	7,7,7	0.29	0	9,10,10	0.35	0
3	GOL	D	903	-	5,5,5	0.93	0	5,5,5	1.16	0
4	ACT	A	907	-	3,3,3	1.26	0	3,3,3	1.21	0
4	ACT	C	904	-	3,3,3	1.39	1 (33%)	3,3,3	1.30	0
2	MPD	D	901	-	7,7,7	0.33	0	9,10,10	0.33	0
3	GOL	C	903	-	5,5,5	1.00	0	5,5,5	1.22	0
2	MPD	B	501	-	7,7,7	0.49	0	9,10,10	1.02	0
3	GOL	C	902	-	5,5,5	1.14	0	5,5,5	0.90	0
4	ACT	E	905	-	3,3,3	1.23	0	3,3,3	1.39	0
3	GOL	A	903	-	5,5,5	1.05	0	5,5,5	1.01	0
3	GOL	B	505	-	5,5,5	1.04	0	5,5,5	1.01	0
3	GOL	E	902	-	5,5,5	1.04	0	5,5,5	0.96	0
3	GOL	A	908	-	5,5,5	1.04	0	5,5,5	1.06	0
3	GOL	B	502	-	5,5,5	1.46	1 (20%)	5,5,5	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	E	901	-	-	1/5/5/5	-
3	GOL	B	506	-	-	0/4/4/4	-
3	GOL	E	903	-	-	3/4/4/4	-
2	MPD	A	902	-	-	0/5/5/5	-
3	GOL	A	909	-	-	0/4/4/4	-
2	MPD	D	902	-	-	3/5/5/5	-
2	MPD	C	901	-	-	0/5/5/5	-
2	MPD	A	901	-	-	1/5/5/5	-
3	GOL	A	905	-	-	3/4/4/4	-
3	GOL	A	906	-	-	0/4/4/4	-
3	GOL	E	904	-	-	2/4/4/4	-
2	MPD	B	503	-	-	3/5/5/5	-
3	GOL	D	903	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	D	901	-	-	0/5/5/5	-
3	GOL	C	903	-	-	1/4/4/4	-
2	MPD	B	501	-	-	0/5/5/5	-
3	GOL	C	902	-	-	4/4/4/4	-
3	GOL	A	903	-	-	2/4/4/4	-
3	GOL	B	505	-	-	0/4/4/4	-
3	GOL	E	902	-	-	0/4/4/4	-
3	GOL	A	908	-	-	2/4/4/4	-
3	GOL	B	502	-	-	2/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	904	ACT	CH3-C	3.57	1.63	1.49
3	B	502	GOL	O2-C2	-2.74	1.35	1.43
3	A	906	GOL	C1-C2	2.62	1.61	1.51
3	A	905	GOL	C3-C2	2.30	1.60	1.51
4	B	507	ACT	CH3-C	2.29	1.58	1.49
3	A	909	GOL	O2-C2	-2.25	1.36	1.43
3	E	903	GOL	C3-C2	2.24	1.60	1.51
4	B	504	ACT	CH3-C	2.12	1.57	1.49
4	C	904	ACT	CH3-C	2.06	1.57	1.49

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	906	GOL	C3-C2-C1	-2.52	102.56	111.80
3	B	506	GOL	C3-C2-C1	-2.06	104.22	111.80
4	F	501	ACT	OXT-C-O	2.04	129.58	122.03
4	B	507	ACT	OXT-C-O	2.00	129.46	122.03

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	903	GOL	C1-C2-C3-O3
3	E	904	GOL	O1-C1-C2-C3
3	E	904	GOL	O1-C1-C2-O2
3	A	903	GOL	O1-C1-C2-C3
3	A	905	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	908	GOL	C1-C2-C3-O3
3	B	502	GOL	O1-C1-C2-C3
3	C	902	GOL	O1-C1-C2-C3
3	C	902	GOL	C1-C2-C3-O3
3	D	903	GOL	O1-C1-C2-C3
3	E	903	GOL	O1-C1-C2-C3
3	C	902	GOL	O1-C1-C2-O2
3	D	903	GOL	O2-C2-C3-O3
3	D	903	GOL	O1-C1-C2-O2
3	A	903	GOL	O1-C1-C2-O2
3	A	905	GOL	O2-C2-C3-O3
3	A	908	GOL	O1-C1-C2-O2
3	C	903	GOL	O2-C2-C3-O3
3	E	903	GOL	O1-C1-C2-O2
3	C	902	GOL	O2-C2-C3-O3
2	B	503	MPD	C1-C2-C3-C4
3	E	903	GOL	O2-C2-C3-O3
3	A	905	GOL	O1-C1-C2-O2
2	D	902	MPD	O2-C2-C3-C4
2	A	901	MPD	C2-C3-C4-O4
2	B	503	MPD	C2-C3-C4-O4
2	D	902	MPD	C2-C3-C4-O4
2	E	901	MPD	C2-C3-C4-O4
2	B	503	MPD	CM-C2-C3-C4
2	D	902	MPD	C1-C2-C3-C4
3	B	502	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	901	MPD	1	0
3	E	903	GOL	1	0
2	D	902	MPD	2	0
4	A	904	ACT	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	116/118 (98%)	-0.02	4 (3%)	48	54	32, 44, 68, 73	0
1	B	117/118 (99%)	0.20	2 (1%)	69	76	33, 53, 82, 93	0
1	C	116/118 (98%)	0.33	4 (3%)	48	54	35, 70, 105, 120	0
1	D	111/118 (94%)	0.58	5 (4%)	38	44	44, 75, 120, 133	0
1	E	116/118 (98%)	0.56	4 (3%)	48	54	45, 85, 127, 137	0
1	F	116/118 (98%)	0.77	6 (5%)	33	38	76, 97, 115, 126	0
1	G	116/118 (98%)	0.69	6 (5%)	33	38	78, 102, 120, 127	0
All	All	808/826 (97%)	0.44	31 (3%)	44	50	32, 78, 118, 137	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	411	PRO	5.1
1	F	301	ALA	4.7
1	E	301	ALA	4.5
1	G	301	ALA	4.3
1	F	310	CYS	3.4
1	D	301	ALA	3.4
1	B	301	ALA	3.1
1	C	301	ALA	3.0
1	A	301	ALA	3.0
1	A	340	CYS	2.9
1	F	416	VAL	2.9
1	B	310	CYS	2.9
1	C	340	CYS	2.7
1	A	416	VAL	2.5
1	E	340	CYS	2.5
1	E	363	TRP	2.5
1	C	310	CYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	363	TRP	2.4
1	F	412	LEU	2.4
1	D	363	TRP	2.4
1	G	413	TYR	2.3
1	E	357	SER	2.3
1	A	363	TRP	2.2
1	F	363	TRP	2.2
1	D	360	LEU	2.2
1	D	368	LEU	2.1
1	G	416	VAL	2.1
1	G	391	ILE	2.0
1	C	412	LEU	2.0
1	F	389	GLY	2.0
1	G	392	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACT	B	504	4/4	0.73	0.17	87,89,89,89	0
4	ACT	A	904	4/4	0.74	0.24	62,66,68,69	0
4	ACT	C	904	4/4	0.82	0.18	93,93,94,94	0
4	ACT	F	501	4/4	0.82	0.13	83,85,86,88	0
3	GOL	A	905	6/6	0.83	0.14	51,55,55,55	0
3	GOL	B	506	6/6	0.83	0.16	72,73,74,75	0
3	GOL	E	904	6/6	0.84	0.10	98,99,99,99	0
3	GOL	D	903	6/6	0.85	0.11	52,55,57,59	0
4	ACT	B	507	4/4	0.85	0.18	71,77,77,79	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	908	6/6	0.85	0.15	85,88,88,89	0
2	MPD	A	902	8/8	0.85	0.22	84,87,90,90	0
3	GOL	A	903	6/6	0.86	0.14	92,94,96,98	0
3	GOL	E	903	6/6	0.87	0.10	49,53,57,57	0
3	GOL	A	909	6/6	0.87	0.16	72,78,79,80	0
3	GOL	E	902	6/6	0.87	0.14	97,98,98,99	0
4	ACT	A	907	4/4	0.87	0.14	46,55,55,62	0
2	MPD	B	503	8/8	0.88	0.17	85,87,90,91	0
4	ACT	E	905	4/4	0.88	0.17	100,101,102,102	0
3	GOL	C	903	6/6	0.88	0.11	43,53,55,55	0
3	GOL	B	502	6/6	0.89	0.18	67,75,77,81	0
3	GOL	C	902	6/6	0.89	0.14	77,78,79,80	0
3	GOL	A	906	6/6	0.91	0.12	51,52,54,54	0
2	MPD	D	902	8/8	0.91	0.21	92,94,95,96	0
3	GOL	B	505	6/6	0.93	0.09	74,80,83,86	0
2	MPD	D	901	8/8	0.93	0.13	64,71,74,76	0
2	MPD	E	901	8/8	0.93	0.12	69,77,81,83	0
2	MPD	B	501	8/8	0.94	0.10	38,45,48,50	0
2	MPD	A	901	8/8	0.94	0.12	48,56,63,66	0
2	MPD	C	901	8/8	0.95	0.12	61,66,69,72	0

6.5 Other polymers

There are no such residues in this entry.