



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:38 AM UTC

PDB ID : 2R71 / pdb_00002r71
Title : Crystal structure of the complex of bovine C-lobe with inositol at 2.1Å resolution
Authors : Mir, R.; Jain, R.; Kumar, S.; Sinha, M.; Singh, N.; Sharma, S.; Kaur, P.; Singh, T.P.
Deposited on : 2007-09-07
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	4% 85% 13% ..
2	B	5	40% 60%
3	C	2	100%
4	D	6	17% 83%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 3086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	341	2604	1622	454	507	21	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	565	LYS	ASN	SEE REMARK 999	UNP P24627
A	608	GLU	LYS	SEE REMARK 999	UNP P24627

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-[beta-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	5	61	34	2	25	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 5 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Fe	0	0
			1	1		

- Molecule 6 is CARBONATE ION (CCD ID: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Zn	0	0
			2	2		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is 1,2,3,4,5,6-HEXAHYDROXY-CYCLOHEXANE (CCD ID: INS) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			12	6	6		
9	A	1	Total	C	O	0	0
			12	6	6		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	280	Total	O	0	0
			280	280		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lactotransferrin



• Molecule 2: beta-D-mannopyranose-(1-4)-[beta-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.48Å 50.31Å 65.97Å 90.00° 107.84° 90.00°	Depositor
Resolution (Å)	62.75 – 2.07 62.75 – 2.07	Depositor EDS
% Data completeness (in resolution range)	97.9 (62.75-2.07) 97.9 (62.75-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.183 , 0.229 0.182 , 0.227	Depositor DCC
R_{free} test set	1210 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.311	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.041 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3086	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, INS, NAG, BMA, FE, SO4, MAN, CO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2652	0.87	0/3591

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2604	0	2519	28	0
2	B	61	0	52	0	0
3	C	28	0	25	0	0
4	D	72	0	61	0	0
5	A	1	0	0	0	0
6	A	4	0	0	0	0
7	A	2	0	0	0	0
8	A	10	0	0	0	0
9	A	24	0	24	0	0
10	A	280	0	0	7	0
All	All	3086	0	2681	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:638:ASN:HD22	1:A:643:ASP:H	1.26	0.80
1:A:565:LYS:HD3	1:A:567:GLU:H	1.49	0.78
1:A:416:LYS:HG2	10:A:153:HOH:O	1.94	0.67
1:A:638:ASN:ND2	1:A:643:ASP:H	1.95	0.64
1:A:559:ASP:HA	1:A:562:LYS:HD3	1.80	0.62
1:A:471:MET:HE3	1:A:483:PHE:HB2	1.86	0.58
1:A:626:PRO:HA	1:A:630:CYS:SG	2.45	0.56
1:A:452:LYS:HE2	1:A:486:PHE:HA	1.88	0.54
1:A:514:LYS:HD3	10:A:272:HOH:O	2.07	0.54
1:A:635:GLU:O	1:A:636:THR:HG22	2.07	0.54
1:A:606:HIS:HB2	10:A:235:HOH:O	2.08	0.52
1:A:460:ALA:HB3	1:A:463:ARG:HD3	1.92	0.51
1:A:344:ARG:HG3	10:A:276:HOH:O	2.11	0.50
1:A:527:THR:HG21	1:A:636:THR:O	2.12	0.49
1:A:415:ARG:NH2	1:A:432:GLY:O	2.41	0.49
1:A:665:TYR:CZ	1:A:669:ILE:HD11	2.49	0.48
1:A:490:SER:HB2	1:A:501:LEU:HA	1.96	0.47
1:A:430:THR:HB	1:A:594:ASN:ND2	2.32	0.45
1:A:395:ASP:HA	1:A:595:HIS:CD2	2.51	0.45
1:A:669:ILE:O	1:A:673:LYS:HG2	2.18	0.42
1:A:632:PHE:O	1:A:633:LYS:HD2	2.20	0.42
1:A:511:GLY:O	10:A:97:HOH:O	2.22	0.42
1:A:606:HIS:CD2	10:A:235:HOH:O	2.72	0.42
1:A:565:LYS:HD3	1:A:567:GLU:N	2.28	0.41
1:A:446:LEU:HD11	1:A:451:LEU:HD23	2.02	0.41
1:A:520:LYS:HD2	1:A:520:LYS:HA	1.65	0.41
1:A:653:GLY:O	1:A:654:ARG:C	2.64	0.41
1:A:636:THR:HG21	10:A:94:HOH:O	2.19	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	337/345 (98%)	320 (95%)	14 (4%)	3 (1%)	14 7

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	684	CYS
1	A	634	SER
1	A	683	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	282/286 (99%)	271 (96%)	11 (4%)	28 23

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	367	GLN
1	A	414	ASN
1	A	417	SER
1	A	419	LYS
1	A	452	LYS
1	A	559	ASP
1	A	565	LYS
1	A	567	GLU
1	A	627	ASP
1	A	631	LEU
1	A	636	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	393	ASN
1	A	414	ASN
1	A	585	GLN
1	A	594	ASN
1	A	606	HIS
1	A	621	ASN
1	A	638	ASN
1	A	642	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.55	0	17,19,21	0.77	1 (5%)
2	NAG	B	2	2	14,14,15	0.45	0	17,19,21	1.07	1 (5%)
2	MAN	B	3	2	11,11,12	0.54	0	15,15,17	0.84	1 (6%)
2	BMA	B	4	2	11,11,12	0.59	0	15,15,17	0.79	0
2	BMA	B	5	2	11,11,12	0.58	0	15,15,17	0.56	0
3	NAG	C	1	1,3	14,14,15	0.61	0	17,19,21	1.67	4 (23%)
3	NAG	C	2	3	14,14,15	0.62	0	17,19,21	1.21	1 (5%)
4	NAG	D	1	1,4	14,14,15	0.61	0	17,19,21	1.38	1 (5%)
4	NAG	D	2	4	14,14,15	0.49	0	17,19,21	1.05	1 (5%)
4	BMA	D	3	4	11,11,12	0.80	0	15,15,17	1.62	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BMA	D	4	4	11,11,12	0.43	0	15,15,17	0.86	1 (6%)
4	BMA	D	5	4	11,11,12	0.67	0	15,15,17	1.11	1 (6%)
4	BMA	D	6	4	11,11,12	0.58	0	15,15,17	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	MAN	B	3	2	-	0/2/19/22	0/1/1/1
2	BMA	B	4	2	-	2/2/19/22	0/1/1/1
2	BMA	B	5	2	-	0/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	C	2	3	-	3/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
4	BMA	D	4	4	-	0/2/19/22	0/1/1/1
4	BMA	D	5	4	-	0/2/19/22	0/1/1/1
4	BMA	D	6	4	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	3	BMA	C1-C2-C3	4.84	116.69	109.64
4	D	1	NAG	C1-O5-C5	4.67	118.44	112.19
3	C	1	NAG	C4-C3-C2	3.92	116.77	111.02
2	B	2	NAG	C1-O5-C5	3.63	117.05	112.19
3	C	2	NAG	C4-C3-C2	3.39	115.98	111.02
3	C	1	NAG	C1-O5-C5	3.37	116.70	112.19
4	D	2	NAG	C1-O5-C5	2.81	115.95	112.19
4	D	3	BMA	C2-C3-C4	2.67	115.55	110.86
4	D	5	BMA	C3-C4-C5	2.60	114.95	110.23
3	C	1	NAG	O5-C1-C2	-2.51	107.41	111.29
2	B	1	NAG	C1-O5-C5	2.37	115.36	112.19
2	B	3	MAN	C1-O5-C5	2.24	115.19	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	4	BMA	C3-C4-C5	2.18	114.18	110.23
3	C	1	NAG	C3-C4-C5	2.10	114.04	110.23

There are no chirality outliers.


All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	4	BMA	O5-C5-C6-O6
4	D	6	BMA	O5-C5-C6-O6
3	C	2	NAG	C8-C7-N2-C2
2	B	4	BMA	C4-C5-C6-O6
3	C	2	NAG	O7-C7-N2-C2
4	D	6	BMA	C4-C5-C6-O6
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
3	C	2	NAG	C3-C2-N2-C7
3	C	1	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	CO3	A	695	5	3,3,3	0.84	0	2,3,3	1.08	0
9	INS	A	788	-	12,12,12	0.39	0	18,18,18	0.61	0
8	SO4	A	2001	-	4,4,4	0.23	0	6,6,6	0.11	0
8	SO4	A	2002	-	4,4,4	0.23	0	6,6,6	0.13	0
9	INS	A	787	-	12,12,12	0.43	0	18,18,18	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	INS	A	788	-	-	-	0/1/1/1
9	INS	A	787	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/345 (98%)	-0.02	14 (4%) 41 42	16, 28, 48, 70	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	342	TYR	4.8
1	A	681	LEU	4.0
1	A	682	GLU	3.9
1	A	683	ALA	3.8
1	A	418	SER	3.8
1	A	420	HIS	3.7
1	A	685	ALA	3.5
1	A	684	CYS	2.6
1	A	422	SER	2.4
1	A	419	LYS	2.4
1	A	343	THR	2.3
1	A	417	SER	2.3
1	A	627	ASP	2.2
1	A	445	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	B	3	11/12	0.26	0.23	64,65,65,66	11
2	BMA	B	5	11/12	0.34	0.19	66,66,67,67	11
4	BMA	D	4	11/12	0.42	0.15	65,66,67,68	11
4	BMA	D	5	11/12	0.43	0.20	70,70,71,71	11
2	BMA	B	4	11/12	0.47	0.16	65,66,66,66	11
4	BMA	D	6	11/12	0.51	0.17	71,72,72,72	11
4	BMA	D	3	11/12	0.53	0.15	59,60,61,63	11
3	NAG	C	2	14/15	0.59	0.17	61,62,63,63	14
3	NAG	C	1	14/15	0.77	0.13	52,55,58,60	0
2	NAG	B	2	14/15	0.79	0.14	59,61,63,63	0
4	NAG	D	2	14/15	0.80	0.14	49,52,55,57	0
2	NAG	B	1	14/15	0.86	0.12	49,51,55,56	0
4	NAG	D	1	14/15	0.91	0.07	31,36,40,45	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	SO4	A	2002	5/5	0.64	0.16	67,67,67,67	5
9	INS	A	788	12/12	0.75	0.28	64,64,64,65	12
9	INS	A	787	12/12	0.76	0.35	104,104,104,104	12
8	SO4	A	2001	5/5	0.91	0.07	74,75,75,75	0
6	CO3	A	695	4/4	0.96	0.06	15,16,16,17	0
7	ZN	A	1002	1/1	0.98	0.04	34,34,34,34	0
7	ZN	A	1001	1/1	1.00	0.02	26,26,26,26	0
5	FE	A	694	1/1	1.00	0.02	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.