



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 05:53 PM UTC

PDB ID : 2R72 / pdb_00002r72
Title : Crystal structure of infectious bursal disease virus VP1 polymerase, incubated with Mg²⁺ ion.
Authors : Garriga, D.; Navarro, A.; Querol-Audi, J.; Abaitua, F.; Rodriguez, J.F.; Verdaguier, N.
Deposited on : 2007-09-07
Resolution : 3.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2361 (3.20-3.12)
Clashscore	190562	2486 (3.20-3.12)
Ramachandran outliers	187476	2405 (3.20-3.12)
Sidechain outliers	187428	2404 (3.20-3.12)
RSRZ outliers	180081	2361 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	852	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INFECTIOUS BURSAL DISEASE VIRUS VP1 POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	765	Total	C	N	O	S	0	0	0
			5915	3787	1007	1099	22			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Mg	0	0
			3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INFECTIOUS BURSAL DISEASE VIRUS VP1 POLYMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	122.84Å 122.84Å 354.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 3.15 20.00 – 3.15	Depositor EDS
% Data completeness (in resolution range)	97.5 (20.00-3.15) 97.2 (20.00-3.15)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 3.15Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.209 , 0.235 0.206 , 0.230	Depositor DCC
R_{free} test set	1393 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	69.8	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5921	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/6053	0.82	3/8231 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	634	TYR	CA-C-N	6.73	126.24	119.24
1	A	634	TYR	C-N-CA	6.73	126.24	119.24
1	A	762	PRO	N-CA-CB	6.62	110.28	103.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5915	0	5861	69	0
2	A	3	0	0	0	0
3	A	3	0	0	0	0
All	All	5921	0	5861	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:SER:HB2	1:A:279:ASN:HD21	1.22	1.04
1:A:230:SER:HB2	1:A:279:ASN:ND2	2.04	0.69
1:A:427:GLN:HA	1:A:430:GLN:HG2	1.76	0.68
1:A:162:LYS:HA	1:A:326:ASP:HA	1.75	0.67
1:A:65:ARG:H	1:A:476:GLN:HE22	1.46	0.64
1:A:170:MET:O	1:A:174:ASN:HB2	1.98	0.64
1:A:600:PRO:HB3	1:A:622:ARG:HD2	1.81	0.63
1:A:41:GLU:HG2	1:A:118:LYS:HD2	1.81	0.62
1:A:103:ASP:O	1:A:104:GLN:HB2	2.00	0.61
1:A:435:TYR:CE2	1:A:439:ARG:HG3	2.34	0.61
1:A:111:TYR:HE1	1:A:473:MET:HE2	1.65	0.60
1:A:375:PHE:CE1	1:A:385:ILE:HD13	2.37	0.60
1:A:459:ASN:O	1:A:797:THR:HG21	2.02	0.59
1:A:521:GLU:HB2	1:A:526:ILE:O	2.03	0.58
1:A:438:THR:O	1:A:442:SER:HB2	2.05	0.57
1:A:429:MET:HE2	1:A:433:MET:HE2	1.86	0.56
1:A:364:LEU:HD21	1:A:385:ILE:HD12	1.88	0.55
1:A:209:THR:O	1:A:371:SER:HB2	2.06	0.55
1:A:176:LEU:HD11	1:A:318:LEU:HD11	1.90	0.54
1:A:472:ILE:HG23	1:A:473:MET:H	1.72	0.53
1:A:230:SER:CB	1:A:279:ASN:HD21	2.09	0.53
1:A:318:LEU:HD22	1:A:336:ASN:HB3	1.91	0.53
1:A:65:ARG:H	1:A:476:GLN:NE2	2.05	0.53
1:A:44:LEU:HD12	1:A:170:MET:HA	1.91	0.53
1:A:303:LEU:HD21	1:A:744:LEU:HD13	1.92	0.52
1:A:170:MET:HG3	1:A:331:LEU:O	2.10	0.51
1:A:565:THR:HG22	1:A:576:THR:HB	1.92	0.51
1:A:793:ALA:O	1:A:797:THR:HG22	2.10	0.51
1:A:108:PRO:HB2	1:A:471:LEU:HD22	1.93	0.50
1:A:544:LEU:HD11	1:A:566:VAL:HG23	1.93	0.50
1:A:192:PRO:HA	1:A:195:LEU:HD12	1.95	0.49
1:A:418:GLU:HB2	1:A:527:ASN:HB2	1.95	0.47
1:A:303:LEU:HD11	1:A:744:LEU:HD22	1.95	0.47
1:A:220:LYS:N	1:A:221:PRO:HD3	2.29	0.47
1:A:346:LEU:HG	1:A:350:MET:HE2	1.98	0.46
1:A:254:LEU:HD22	1:A:271:ILE:HD11	1.97	0.46
1:A:84:TRP:C	1:A:86:ARG:H	2.24	0.45
1:A:548:ALA:HB3	1:A:579:LYS:HE3	1.98	0.45
1:A:82:LEU:HD22	1:A:85:MET:HE2	1.98	0.45
1:A:482:GLN:CD	1:A:489:THR:HG22	2.42	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:THR:HG22	1:A:450:ASN:HB3	2.00	0.44
1:A:569:ASP:OD2	1:A:592:ARG:NH1	2.50	0.44
1:A:407:VAL:HA	1:A:411:THR:O	2.17	0.44
1:A:65:ARG:NH1	1:A:476:GLN:HE21	2.16	0.44
1:A:425:THR:H	1:A:428:HIS:HD2	1.66	0.44
1:A:667:SER:O	1:A:673:GLY:HA3	2.18	0.44
1:A:288:LEU:HD13	1:A:303:LEU:HD12	2.00	0.43
1:A:412:TRP:O	1:A:533:SER:HA	2.18	0.43
1:A:67:LEU:HD22	1:A:108:PRO:HG3	1.99	0.43
1:A:289:LEU:HD22	1:A:747:ALA:HB3	2.00	0.43
1:A:162:LYS:HG2	1:A:326:ASP:HB3	2.01	0.43
1:A:698:PRO:HA	1:A:699:PRO:HD3	1.91	0.42
1:A:553:LEU:HD13	1:A:585:VAL:HG21	2.00	0.42
1:A:218:ASP:C	1:A:220:LYS:H	2.27	0.42
1:A:134:MET:HE1	1:A:151:GLU:HB3	2.00	0.42
1:A:391:ALA:HA	1:A:392:PRO:HD3	1.91	0.42
1:A:742:SER:HA	1:A:786:ALA:HA	2.01	0.42
1:A:65:ARG:HB2	1:A:476:GLN:HE22	1.85	0.42
1:A:318:LEU:HD23	1:A:318:LEU:HA	1.91	0.42
1:A:627:ARG:HB3	1:A:685:VAL:HG11	2.02	0.42
1:A:425:THR:H	1:A:428:HIS:CD2	2.38	0.41
1:A:511:ARG:HD3	1:A:513:ASP:OD1	2.19	0.41
1:A:94:LYS:HA	1:A:95:PRO:HD3	1.89	0.41
1:A:411:THR:HB	1:A:413:TYR:CE1	2.55	0.41
1:A:731:LEU:HD11	1:A:755:GLU:HG2	2.02	0.41
1:A:628:LEU:HD13	1:A:685:VAL:HG22	2.04	0.40
1:A:385:ILE:O	1:A:388:TRP:HB3	2.21	0.40
1:A:563:SER:HA	1:A:564:PRO:HD3	1.95	0.40
1:A:289:LEU:HD11	1:A:455:THR:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	755/852 (89%)	705 (93%)	37 (5%)	13 (2%)	7	30

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	30	GLU
1	A	767	ALA
1	A	221	PRO
1	A	473	MET
1	A	607	SER
1	A	609	LYS
1	A	94	LYS
1	A	472	ILE
1	A	605	ASN
1	A	239	VAL
1	A	420	GLY
1	A	84	TRP
1	A	630	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	626/730 (86%)	596 (95%)	30 (5%)	23	52

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	GLU
1	A	30	GLU
1	A	46	SER
1	A	65	ARG
1	A	145	ASN
1	A	180	LYS
1	A	235	LEU
1	A	253	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	296	LYS
1	A	299	ASN
1	A	352	THR
1	A	385	ILE
1	A	398	LEU
1	A	399	VAL
1	A	407	VAL
1	A	451	GLN
1	A	455	THR
1	A	465	VAL
1	A	501	LEU
1	A	529	LYS
1	A	560	GLU
1	A	606	LYS
1	A	668	GLU
1	A	684	THR
1	A	744	LEU
1	A	757	LEU
1	A	789	VAL
1	A	794	LEU
1	A	797	THR
1	A	802	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	ASN
1	A	157	GLN
1	A	172	GLN
1	A	204	GLN
1	A	291	GLN
1	A	299	ASN
1	A	336	ASN
1	A	358	ASN
1	A	376	ASN
1	A	423	ASN
1	A	428	HIS
1	A	476	GLN
1	A	482	GLN
1	A	494	HIS
1	A	503	GLN
1	A	616	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	765/852 (89%)	0.18	31 (4%) 41 24	51, 62, 80, 89	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	ASP	6.8
1	A	607	SER	5.4
1	A	766	ASP	4.6
1	A	608	LEU	4.5
1	A	767	ALA	4.4
1	A	219	ASP	4.3
1	A	803	ALA	3.7
1	A	239	VAL	3.7
1	A	221	PRO	3.7
1	A	486	ASN	3.5
1	A	487	ALA	3.5
1	A	762	PRO	3.5
1	A	609	LYS	3.4
1	A	218	ASP	3.4
1	A	604	GLU	3.4
1	A	235	LEU	3.4
1	A	28	VAL	3.3
1	A	488	ALA	3.3
1	A	612	VAL	3.3
1	A	87	GLN	3.0
1	A	86	ARG	3.0
1	A	220	LYS	3.0
1	A	771	GLU	3.0
1	A	216	GLY	3.0
1	A	237	GLY	3.0
1	A	605	ASN	2.8
1	A	88	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	238	ASP	2.5
1	A	448	MET	2.2
1	A	603	VAL	2.2
1	A	240	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	A	846	1/1	0.84	0.35	71,71,71,71	0
2	MG	A	847	1/1	0.87	0.21	103,103,103,103	0
2	MG	A	848	1/1	0.87	0.24	86,86,86,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.