



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:35 AM UTC

PDB ID : 3R76 / pdb_00003r76
Title : Crystal structure of 2-amino-2-desoxyisochorismate synthase (ADIC) synthase PhzE from Burkholderia lata 383 in complex with benzoate, pyruvate and glutamine
Authors : Li, Q.A.; Mavrodi, D.V.; Thomashow, L.S.; Roessle, M.; Blankenfeldt, W.
Deposited on : 2011-03-22
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	 83% 11% . 5%
1	B	645	 79% 15% . .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Anthranilate/para-aminobenzoate synthases component I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	612	Total	C	N	O	S	0	2	0
			4595	2887	822	863	23			
1	B	622	Total	C	N	O	S	0	1	0
			4692	2937	845	888	22			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q396C7
A	0	HIS	-	expression tag	UNP Q396C7
B	-1	GLY	-	expression tag	UNP Q396C7
B	0	HIS	-	expression tag	UNP Q396C7

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is BENZOIC ACID (CCD ID: BEZ) (formula: C₇H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			9	7	2		
3	B	1	Total	C	O	0	0
			9	7	2		

- Molecule 4 is PYRUVIC ACID (CCD ID: PYR) (formula: $C_3H_4O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	133	Total	O	0	0
			133	133		
6	B	110	Total	O	0	0
			110	110		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Anthranilate/para-aminobenzoate synthases component I



• Molecule 1: Anthranilate/para-aminobenzoate synthases component I



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	258.17Å 97.79Å 53.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.66 – 2.60 29.66 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.66-2.60) 99.7 (29.66-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.47 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.163 , 0.225 0.165 , 0.223	Depositor DCC
R_{free} test set	2227 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	42.8	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9572	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CYG, PO4, PYR, BEZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	1/4677 (0.0%)	0.96	1/6360 (0.0%)
1	B	0.94	0/4774	1.00	7/6490 (0.1%)
All	All	0.96	1/9451 (0.0%)	0.98	8/12850 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	366	VAL	CA-CB	-6.21	1.48	1.55

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	272	ARG	CA-CB-CG	-7.13	99.83	114.10
1	B	487	VAL	CB-CA-C	-7.05	98.51	110.71
1	B	429	GLY	N-CA-C	6.90	129.53	113.18
1	B	443	SER	N-CA-C	6.23	118.59	111.11
1	A	633	ARG	N-CA-C	6.18	118.01	111.28
1	B	541	VAL	CB-CA-C	-5.75	100.58	110.30
1	B	396	PRO	O-C-N	5.45	123.82	121.31
1	B	310	ILE	CB-CA-C	-5.08	105.47	111.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4595	0	4462	52	0
1	B	4692	0	4558	69	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	9	0	5	0	0
3	B	9	0	5	0	0
4	A	6	0	0	0	0
4	B	6	0	0	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	133	0	0	0	0
6	B	110	0	0	7	0
All	All	9572	0	9030	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:9:LEU:HD13	1:B:39:VAL:HG22	1.45	0.98
1:A:452:ALA:HB3	1:A:492:PRO:O	1.73	0.89
1:A:255:MET:HE3	1:A:299:THR:HB	1.55	0.86
1:A:213:MET:HE1	1:A:255:MET:HE2	1.57	0.83
1:A:310:ILE:O	1:A:310:ILE:HD12	1.78	0.81
1:B:292:VAL:HG11	1:B:343:ARG:HD2	1.64	0.80
1:A:9:LEU:HD13	1:A:39:VAL:HG12	1.64	0.79
1:B:608:MET:HE1	6:B:654:HOH:O	1.82	0.78
1:B:344:THR:HG23	6:B:657:HOH:O	1.82	0.78
1:A:175:LEU:HD13	1:A:393:ALA:CB	2.17	0.75
1:B:277:LEU:C	1:B:277:LEU:HD12	2.13	0.73
1:A:403:LEU:HD13	1:B:232:ILE:HG13	1.71	0.73
1:A:210[B]:CYS:SG	6:B:660:HOH:O	2.46	0.73
1:B:292:VAL:HG13	1:B:343:ARG:HG2	1.72	0.72
1:B:512:LEU:HD23	1:B:531:ILE:HG22	1.73	0.70
1:A:213:MET:CE	1:A:255:MET:HE2	2.21	0.70
1:B:75:ARG:HA	1:B:84:THR:HG21	1.74	0.69
1:A:255:MET:HE3	1:A:299:THR:CB	2.24	0.67
1:A:277:LEU:C	1:A:277:LEU:HD12	2.19	0.67
1:B:253:LEU:HD13	1:B:284:ILE:HD13	1.75	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:292:VAL:HG11	1:B:343:ARG:CD	2.26	0.64
1:A:175:LEU:HD13	1:A:393:ALA:HB2	1.80	0.63
1:A:523:MET:HE1	1:A:624:LEU:HD22	1.80	0.63
1:B:248:VAL:HG11	1:B:304:THR:HG22	1.79	0.63
1:B:608:MET:CE	6:B:654:HOH:O	2.44	0.62
1:A:403:LEU:HD22	1:B:229:ILE:HD13	1.80	0.62
1:B:495:GLY:O	1:B:526:CYG:N1	2.33	0.62
1:B:543:ARG:NH1	1:B:596:GLU:OE2	2.33	0.61
1:B:440:ALA:N	1:B:441:GLU:HA	2.15	0.60
1:B:69:LEU:HD13	1:B:334:LEU:HD13	1.83	0.60
1:B:452:ALA:HB3	1:B:492:PRO:O	2.02	0.59
1:A:547:ASN:ND2	1:A:550:ILE:HG21	2.18	0.58
1:B:344:THR:CG2	6:B:657:HOH:O	2.45	0.58
1:B:292:VAL:CG1	1:B:343:ARG:HG2	2.33	0.58
1:B:406:HIS:CD2	1:B:408:SER:OG	2.56	0.58
1:A:258:ARG:NH2	1:A:298:GLU:OE1	2.36	0.58
1:A:523:MET:HA	1:A:606:SER:O	2.04	0.58
1:B:248:VAL:CG1	1:B:304:THR:HG22	2.33	0.58
1:B:54:PRO:HD2	1:B:68:LEU:HD11	1.87	0.57
1:A:308:SER:HA	1:A:309:PRO:C	2.30	0.57
1:A:403:LEU:CD2	1:B:229:ILE:HD13	2.35	0.56
1:B:490:MET:HG3	1:B:512:LEU:CD1	2.36	0.56
1:A:310:ILE:HD12	1:A:310:ILE:C	2.29	0.56
1:B:175:LEU:HD13	1:B:393:ALA:HB2	1.86	0.56
1:B:480:VAL:HG12	1:B:481:ASP:N	2.21	0.56
1:A:229:ILE:HD11	1:B:406:HIS:CG	2.41	0.55
1:A:403:LEU:HD11	1:B:227:PRO:HB2	1.88	0.55
1:B:308:SER:HA	1:B:309:PRO:C	2.31	0.54
1:B:406:HIS:HD2	1:B:408:SER:OG	1.90	0.54
1:B:427:PRO:O	1:B:430:GLY:HA3	2.07	0.54
1:A:251:GLU:CD	1:A:308:SER:CB	2.80	0.54
1:B:523:MET:HA	1:B:606:SER:O	2.07	0.54
1:B:258:ARG:NH2	1:B:298:GLU:OE1	2.41	0.53
1:A:153:LYS:HE3	1:A:155:THR:OG1	2.09	0.53
1:B:251:GLU:CD	1:B:308:SER:CB	2.82	0.52
1:B:277:LEU:C	1:B:277:LEU:CD1	2.82	0.52
1:A:20:CYS:SG	1:A:44:VAL:HG22	2.50	0.52
1:B:12:ARG:NH2	1:B:98:GLU:OE2	2.42	0.52
1:B:277:LEU:HD12	1:B:277:LEU:O	2.09	0.52
1:B:202:ARG:HD2	6:B:689:HOH:O	2.11	0.51
1:A:403:LEU:HD13	1:B:232:ILE:CG1	2.40	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:615:VAL:HG22	1:B:250:ASP:OD1	2.11	0.50
1:A:9:LEU:HD13	1:A:39:VAL:CG1	2.36	0.50
1:A:547:ASN:HD22	1:A:550:ILE:HG21	1.76	0.50
1:A:193:ARG:HH11	1:A:355:GLU:HB3	1.77	0.50
1:B:476:VAL:HG23	1:B:477:HIS:CD2	2.47	0.49
1:B:496:ASP:O	1:B:505:ILE:HD13	2.13	0.49
1:A:315:ARG:NH2	1:B:453:GLU:OE1	2.45	0.48
1:A:516:ILE:HD11	1:A:532:LEU:CD1	2.44	0.48
1:B:292:VAL:CG1	1:B:343:ARG:CG	2.91	0.48
1:B:561:GLU:OE2	1:B:622:ARG:NH2	2.46	0.48
1:B:71:MET:HB2	1:B:92:VAL:HG13	1.96	0.47
1:B:619:ASP:OD2	1:B:622:ARG:NH1	2.47	0.47
1:B:512:LEU:HD23	1:B:531:ILE:CG2	2.44	0.46
1:B:44:VAL:HG21	1:B:190:THR:O	2.15	0.46
1:A:187:VAL:HA	1:A:195:PHE:O	2.16	0.46
1:B:189:HIS:HD2	1:B:194:THR:OG1	1.99	0.46
1:A:571:ALA:HB3	1:A:597:VAL:HB	1.99	0.45
1:B:532:LEU:O	1:B:536:LEU:HG	2.18	0.44
1:B:270:HIS:HE1	6:B:731:HOH:O	1.99	0.44
1:A:175:LEU:CD1	1:A:393:ALA:HB2	2.45	0.44
1:A:462:GLN:NE2	1:B:261:PRO:O	2.51	0.44
1:A:496:ASP:O	1:A:505:ILE:HD13	2.17	0.44
1:B:494:PRO:O	1:B:504:ARG:NH2	2.37	0.44
1:A:494:PRO:O	1:A:504:ARG:NH2	2.39	0.44
1:A:532:LEU:O	1:A:536:LEU:HG	2.17	0.44
1:A:561:GLU:OE2	1:A:622:ARG:NH2	2.50	0.43
1:B:429:GLY:N	1:B:430:GLY:HA3	2.33	0.43
1:B:187:VAL:HA	1:B:195:PHE:O	2.18	0.43
1:A:251:GLU:CD	1:A:308:SER:HB2	2.44	0.43
1:B:251:GLU:CD	1:B:308:SER:HB2	2.43	0.43
1:A:69:LEU:HD13	1:A:334:LEU:HD13	2.00	0.43
1:B:493:GLY:O	1:B:527:LEU:HD22	2.19	0.43
1:B:440:ALA:N	1:B:441:GLU:CA	2.82	0.42
1:A:372:ARG:NH2	1:B:550:ILE:HD12	2.35	0.42
1:A:55:LEU:HA	1:A:55:LEU:HD23	1.84	0.42
1:A:117:ALA:HA	1:A:395:ASP:OD1	2.20	0.41
1:A:373:HIS:ND1	1:A:373:HIS:N	2.68	0.41
1:A:440:ALA:O	1:A:441:GLU:HB2	2.20	0.41
1:B:476:VAL:HG11	1:B:508:LEU:CD1	2.50	0.41
1:A:547:ASN:HD22	1:A:547:ASN:HA	1.62	0.41
1:B:124:ASP:OD1	1:B:125:ILE:N	2.52	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:317:ILE:HG23	1:B:321:GLU:CD	2.45	0.41
1:A:60:ALA:HB2	1:A:340:ARG:C	2.46	0.41
1:B:159:ASP:OD1	1:B:361:HIS:HD2	2.03	0.41
1:B:202:ARG:HD3	1:B:214:ASN:HB2	2.02	0.41
1:A:124:ASP:OD1	1:A:125:ILE:N	2.52	0.41
1:A:229:ILE:HD11	1:B:406:HIS:CD2	2.55	0.40
1:A:550:ILE:HD12	1:B:372:ARG:NH2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	605/645 (94%)	588 (97%)	16 (3%)	1 (0%)	43 66
1	B	618/645 (96%)	598 (97%)	17 (3%)	3 (0%)	24 46
All	All	1223/1290 (95%)	1186 (97%)	33 (3%)	4 (0%)	36 58

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	28	THR
1	B	29	GLY
1	A	429	GLY
1	B	31	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/506 (91%)	438 (95%)	24 (5%)	21	44
1	B	473/506 (94%)	439 (93%)	34 (7%)	13	30
All	All	935/1012 (92%)	877 (94%)	58 (6%)	16	36

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	14	LEU
1	A	44	VAL
1	A	75	ARG
1	A	103	SER
1	A	106	LEU
1	A	118	LEU
1	A	161	ASP
1	A	169	LEU
1	A	175	LEU
1	A	177	ARG
1	A	187	VAL
1	A	250	ASP
1	A	255	MET
1	A	310	ILE
1	A	334	LEU
1	A	350	LEU
1	A	373	HIS
1	A	377	VAL
1	A	442	LEU
1	A	456	PHE
1	A	465	SER
1	A	547	ASN
1	A	608	MET
1	B	8	ASN
1	B	9	LEU
1	B	14	LEU
1	B	39	VAL
1	B	84	THR
1	B	92	VAL
1	B	103	SER
1	B	118	LEU
1	B	161	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	169	LEU
1	B	175	LEU
1	B	217	SER
1	B	224	GLN
1	B	248	VAL
1	B	253	LEU
1	B	271	LEU
1	B	334	LEU
1	B	344	THR
1	B	350	LEU
1	B	373	HIS
1	B	375	ASP
1	B	387	VAL
1	B	405	GLN
1	B	442	LEU
1	B	453	GLU
1	B	456	PHE
1	B	487	VAL
1	B	508	LEU
1	B	512	LEU
1	B	541	VAL
1	B	547	ASN
1	B	560	ARG
1	B	578	MET
1	B	608	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	189	HIS
1	A	392	ASN
1	A	410	GLN
1	A	533	ASN
1	A	547	ASN
1	B	8	ASN
1	B	17	GLN
1	B	189	HIS
1	B	224	GLN
1	B	361	HIS
1	B	384	HIS
1	B	406	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	417	ASN
1	B	462	GLN
1	B	463	GLN
1	B	477	HIS
1	B	533	ASN
1	B	547	ASN
1	B	572	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CYG	B	526	1	12,14,15	1.96	4 (33%)	10,17,19	1.64	2 (20%)
1	CYG	A	526	1	12,14,15	1.77	3 (25%)	10,17,19	1.62	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CYG	B	526	1	-	0/14/16/18	-
1	CYG	A	526	1	-	1/14/16/18	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	526	CYG	CG1-CD1	4.39	1.55	1.50
1	A	526	CYG	CG1-CD1	3.98	1.54	1.50
1	B	526	CYG	OE2-CD1	3.12	1.25	1.21
1	A	526	CYG	OE2-CD1	2.84	1.25	1.21
1	B	526	CYG	CD1-SG	2.52	1.82	1.76
1	A	526	CYG	CD1-SG	2.31	1.81	1.76
1	B	526	CYG	CB-SG	2.06	1.86	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	526	CYG	CG1-CD1-SG	3.38	117.43	113.40
1	A	526	CYG	OE2-CD1-CG1	-3.35	120.12	123.98
1	B	526	CYG	OE2-CD1-CG1	-3.20	120.29	123.98
1	B	526	CYG	CB1-CA1-N1	2.29	116.09	110.12

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	526	CYG	N-CA-CB-SG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	526	CYG	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	PYR	B	702	-	5,5,5	1.64	2 (40%)	3,6,6	1.29	0
3	BEZ	A	701	2	9,9,9	0.88	0	11,11,11	1.23	2 (18%)
5	PO4	A	900	-	4,4,4	1.33	1 (25%)	6,6,6	0.98	1 (16%)
3	BEZ	B	701	2	9,9,9	0.85	0	11,11,11	0.93	1 (9%)
4	PYR	A	702	-	5,5,5	1.57	1 (20%)	3,6,6	1.58	1 (33%)
5	PO4	B	900	-	4,4,4	0.65	0	6,6,6	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PYR	A	702	-	-	0/4/4/4	-
4	PYR	B	702	-	-	0/4/4/4	-
3	BEZ	B	701	2	-	0/4/4/4	0/1/1/1
3	BEZ	A	701	2	-	0/4/4/4	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	702	PYR	CA-C	-2.70	1.45	1.54
4	B	702	PYR	CA-C	-2.48	1.46	1.54
4	B	702	PYR	OXT-C	-2.29	1.24	1.30
5	A	900	PO4	P-O4	-2.13	1.48	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	BEZ	O2-C-O1	-2.81	117.32	123.35
3	A	701	BEZ	O2-C-C1	2.44	121.09	114.84
3	B	701	BEZ	O2-C-C1	2.37	120.93	114.84
4	A	702	PYR	OXT-C-CA	2.26	119.85	113.59
5	A	900	PO4	O4-P-O3	2.04	114.27	107.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	611/645 (94%)	-0.55	2 (0%) 90 88	23, 50, 80, 127	2 (0%)
1	B	621/645 (96%)	-0.56	3 (0%) 87 85	32, 50, 78, 121	1 (0%)
All	All	1232/1290 (95%)	-0.56	5 (0%) 88 86	23, 50, 79, 127	3 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	636	LYS	2.8
1	B	30	SER	2.6
1	A	402	ALA	2.5
1	A	440	ALA	2.2
1	B	440	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CYG	A	526	15/16	0.94	0.09	47,67,80,83	0
1	CYG	B	526	15/16	0.96	0.09	32,39,45,47	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PO4	B	900	5/5	0.82	0.20	71,96,103,106	0
5	PO4	A	900	5/5	0.90	0.25	20,22,22,23	5
4	PYR	B	702	6/6	0.94	0.09	43,52,58,59	0
3	BEZ	A	701	9/9	0.98	0.04	33,35,37,38	0
3	BEZ	B	701	9/9	0.98	0.06	38,40,41,42	0
4	PYR	A	702	6/6	0.98	0.05	38,42,44,49	0
2	MG	A	700	1/1	1.00	0.03	45,45,45,45	0
2	MG	B	700	1/1	1.00	0.01	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.