



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:04 PM UTC

PDB ID : 8R71 / pdb_00008r71
Title : Polysaccharide lyase BtPL33HA (BT4410) Y291A with HAdp4 collected at 1.22 Å
Authors : Cartmell, A.
Deposited on : 2023-11-23
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	644	 89% 7% . .
1	B	644	 86% 6% 6%
2	C	3	 100%
2	D	3	 33% 67%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20415 atoms, of which 9780 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heparinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	618	Total	C	H	N	O	S	125	6	0
			9958	3197	4930	862	935	34			
1	B	603	Total	C	H	N	O	S	119	2	0
			9628	3093	4764	829	908	34			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP Q89ZG7
A	-6	GLY	-	expression tag	UNP Q89ZG7
A	-5	SER	-	expression tag	UNP Q89ZG7
A	-4	SER	-	expression tag	UNP Q89ZG7
A	-3	HIS	-	expression tag	UNP Q89ZG7
A	-2	HIS	-	expression tag	UNP Q89ZG7
A	-1	HIS	-	expression tag	UNP Q89ZG7
A	0	HIS	-	expression tag	UNP Q89ZG7
A	1	HIS	-	expression tag	UNP Q89ZG7
A	2	HIS	-	expression tag	UNP Q89ZG7
A	3	SER	-	expression tag	UNP Q89ZG7
A	4	SER	-	expression tag	UNP Q89ZG7
A	5	GLY	-	expression tag	UNP Q89ZG7
A	6	LEU	-	expression tag	UNP Q89ZG7
A	7	VAL	-	expression tag	UNP Q89ZG7
A	8	PRO	-	expression tag	UNP Q89ZG7
A	9	ARG	-	expression tag	UNP Q89ZG7
A	10	GLY	-	expression tag	UNP Q89ZG7
A	11	SER	-	expression tag	UNP Q89ZG7
A	12	HIS	-	expression tag	UNP Q89ZG7
A	13	MET	-	expression tag	UNP Q89ZG7
A	14	ALA	-	expression tag	UNP Q89ZG7
A	15	SER	-	expression tag	UNP Q89ZG7
A	283	ALA	TYR	engineered mutation	UNP Q89ZG7
B	-7	MET	-	initiating methionine	UNP Q89ZG7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	GLY	-	expression tag	UNP Q89ZG7
B	-5	SER	-	expression tag	UNP Q89ZG7
B	-4	SER	-	expression tag	UNP Q89ZG7
B	-3	HIS	-	expression tag	UNP Q89ZG7
B	-2	HIS	-	expression tag	UNP Q89ZG7
B	-1	HIS	-	expression tag	UNP Q89ZG7
B	0	HIS	-	expression tag	UNP Q89ZG7
B	1	HIS	-	expression tag	UNP Q89ZG7
B	2	HIS	-	expression tag	UNP Q89ZG7
B	3	SER	-	expression tag	UNP Q89ZG7
B	4	SER	-	expression tag	UNP Q89ZG7
B	5	GLY	-	expression tag	UNP Q89ZG7
B	6	LEU	-	expression tag	UNP Q89ZG7
B	7	VAL	-	expression tag	UNP Q89ZG7
B	8	PRO	-	expression tag	UNP Q89ZG7
B	9	ARG	-	expression tag	UNP Q89ZG7
B	10	GLY	-	expression tag	UNP Q89ZG7
B	11	SER	-	expression tag	UNP Q89ZG7
B	12	HIS	-	expression tag	UNP Q89ZG7
B	13	MET	-	expression tag	UNP Q89ZG7
B	14	ALA	-	expression tag	UNP Q89ZG7
B	15	SER	-	expression tag	UNP Q89ZG7
B	283	ALA	TYR	engineered mutation	UNP Q89ZG7

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	3	Total	C	H	N	O	6	0	0
			71	22	30	2	17			
2	D	3	Total	C	H	N	O	6	0	0
			71	22	30	2	17			

- Molecule 3 is polyethylene glycol (CCD ID: P4K) (formula: C₃₀H₆₂O₁₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	1	0
			25	7	14	4		
3	B	1	Total	C	H	O	0	0
			22	6	12	4		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	408	Total	O	0	0
			408	408		
5	B	230	Total	O	0	0
			230	230		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heparinase



- Molecule 1: Heparinase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

33%

67%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.63Å 137.35Å 202.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.66 – 1.92 60.66 – 1.92	Depositor EDS
% Data completeness (in resolution range)	100.0 (60.66-1.92) 100.0 (60.66-1.92)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.179 , 0.208 0.179 , 0.209	Depositor DCC
R_{free} test set	5759 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	31.8	Xtriage
Anisotropy	0.252	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	20415	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: P4K, BDP, ZN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/5150	1.09	11/6969 (0.2%)
1	B	0.62	0/4981	1.05	9/6742 (0.1%)
All	All	0.65	0/10131	1.07	20/13711 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	5
All	All	0	9

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	286	HIS	CA-CB-CG	-7.40	106.40	113.80
1	A	113	MET	CG-SD-CE	-6.87	85.79	100.90
1	B	52	ARG	CD-NE-CZ	6.41	133.38	124.40
1	A	411	GLU	CB-CG-CD	6.36	123.42	112.60
1	A	218	MET	CG-SD-CE	-5.99	87.73	100.90
1	B	502	ASN	CA-CB-CG	-5.94	106.66	112.60
1	A	246	MET	CG-SD-CE	5.80	113.67	100.90
1	B	274	ASP	CA-CB-CG	5.79	118.39	112.60
1	A	212	ASP	CA-CB-CG	5.70	118.30	112.60
1	B	622	LYS	N-CA-CB	-5.58	101.38	110.59
1	B	351	PHE	CA-CB-CG	5.50	119.30	113.80
1	A	600	ASP	CB-CA-C	-5.42	100.26	110.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	259	ARG	CB-CG-CD	-5.42	98.83	111.30
1	B	212	ASP	CA-CB-CG	5.33	117.93	112.60
1	A	507	GLU	CB-CG-CD	5.30	121.60	112.60
1	A	397	GLU	CB-CA-C	-5.29	100.61	109.65
1	A	351	PHE	CA-CB-CG	5.24	119.04	113.80
1	A	414	PHE	CA-CB-CG	5.24	119.03	113.80
1	B	246	MET	CG-SD-CE	5.17	112.26	100.90
1	B	507	GLU	CB-CG-CD	5.02	121.13	112.60

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	106	ARG	Sidechain
1	A	252	ARG	Sidechain
1	A	316[A]	ARG	Sidechain
1	A	510	ARG	Sidechain
1	B	106	ARG	Sidechain
1	B	21	ARG	Sidechain
1	B	259	ARG	Sidechain
1	B	316	ARG	Sidechain
1	B	531	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5028	4930	4916	35	1
1	B	4864	4764	4753	28	0
2	C	41	30	33	0	0
2	D	41	30	33	2	0
3	A	11	14	0	1	0
3	B	10	12	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	408	0	0	12	0
5	B	230	0	0	7	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	10635	9780	9735	64	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:358:ASN:HB3	5:B:913:HOH:O	1.51	1.10
1:A:35:ASN:HB3	5:A:1071:HOH:O	1.55	1.04
1:B:113:MET:CE	1:B:181:PHE:HD2	1.76	0.97
1:B:113:MET:CE	1:B:181:PHE:CD2	2.54	0.90
1:B:23:MET:HE3	1:B:308:LEU:HD13	1.55	0.88
1:B:113:MET:HE2	1:B:181:PHE:CD2	2.12	0.84
1:B:113:MET:HE1	1:B:177:TRP:HB3	1.61	0.82
1:A:600:ASP:OD2	5:A:801:HOH:O	1.98	0.81
1:A:246:MET:HE1	5:A:1061:HOH:O	1.84	0.77
1:A:245:LEU:HB3	1:A:246:MET:HE2	1.67	0.76
1:A:23:MET:HE3	1:A:308:LEU:HD13	1.68	0.75
1:B:246:MET:HE1	5:B:922:HOH:O	1.85	0.75
1:A:591:LYS:HE3	5:A:1143:HOH:O	1.86	0.74
1:B:532[C]:SER:HG	1:B:543:THR:HG1	1.29	0.73
1:A:245:LEU:HB3	1:A:246:MET:CE	2.20	0.71
1:B:245:LEU:HB3	1:B:246:MET:HE2	1.74	0.69
1:A:356:ASN:HB3	5:A:817:HOH:O	1.92	0.68
1:B:113:MET:HE3	1:B:181:PHE:HD2	1.57	0.67
1:A:451:ILE:HD11	5:A:819:HOH:O	1.95	0.67
1:A:35:ASN:CB	5:A:1071:HOH:O	2.28	0.67
1:A:564:ASN:ND2	5:A:804:HOH:O	2.28	0.66
1:B:245:LEU:HB3	1:B:246:MET:CE	2.27	0.65
1:B:316:ARG:NH2	5:B:803:HOH:O	2.29	0.63
1:A:277:CYS:SG	5:A:1140:HOH:O	2.54	0.62
1:A:23:MET:CE	1:A:308:LEU:HD13	2.32	0.59
1:B:88:GLU:OE1	5:B:801:HOH:O	2.16	0.59
1:A:246:MET:CE	5:A:1061:HOH:O	2.43	0.59
1:A:356:ASN:CB	5:A:817:HOH:O	2.47	0.59
1:B:113:MET:HE2	1:B:181:PHE:CE2	2.39	0.57
1:A:113:MET:HE1	1:A:178:VAL:N	2.21	0.54
1:B:23:MET:CE	1:B:308:LEU:HD13	2.35	0.54
1:B:230:ASN:HB2	1:B:286:HIS:O	2.08	0.54
1:B:246:MET:CE	5:B:922:HOH:O	2.50	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:490:ILE:HD13	1:B:523:GLU:HG2	1.90	0.54
1:A:113:MET:HE3	1:A:178:VAL:HG22	1.89	0.53
5:B:947:HOH:O	2:D:3:NAG:H61	2.08	0.53
1:A:113:MET:HE1	1:A:177:TRP:C	2.32	0.53
1:B:490:ILE:HD12	1:B:524:ALA:HA	1.93	0.51
1:A:461:THR:OG1	5:A:802:HOH:O	2.18	0.49
1:A:469:LYS:O	1:A:470:ASP:HB2	2.13	0.49
1:B:176:ALA:HB1	1:B:246:MET:HE3	1.96	0.48
1:A:21:ARG:O	1:A:26:LYS:HE2	2.13	0.47
1:A:147:LEU:O	1:A:150:GLN:HG2	2.14	0.47
1:B:551:ALA:HB1	1:B:622:LYS:HG2	1.96	0.47
1:A:113:MET:CE	1:A:178:VAL:HA	2.45	0.46
1:B:119:GLU:HG2	1:B:121:LYS:HG3	1.97	0.46
1:B:149:ARG:NH2	2:D:1:NAG:O7	2.41	0.46
1:A:230:ASN:HB2	1:A:286:HIS:O	2.17	0.45
1:A:584:LEU:HD23	1:A:584:LEU:C	2.42	0.45
1:A:147:LEU:N	1:A:148:PRO:CD	2.81	0.44
1:A:176:ALA:HB1	1:A:246:MET:HE3	2.00	0.44
1:B:584:LEU:C	1:B:584:LEU:HD23	2.43	0.44
1:A:551:ALA:HB1	1:A:622:LYS:HG2	1.99	0.43
1:B:453:VAL:HG11	1:B:582:VAL:HG21	1.99	0.43
1:B:147:LEU:N	1:B:148:PRO:CD	2.81	0.43
1:A:245:LEU:HB3	1:A:246:MET:HE3	1.98	0.43
1:A:319:GLU:OE1	3:A:701:P4K:C12	2.67	0.43
1:B:392:ASN:OD1	1:B:392:ASN:N	2.51	0.42
1:A:312:PRO:HB2	1:A:316[A]:ARG:NH2	2.35	0.42
1:A:33[A]:LEU:HD11	1:A:261:MET:SD	2.60	0.41
1:B:307:SER:HA	5:B:927:HOH:O	2.20	0.41
1:A:245:LEU:C	1:A:246:MET:HE2	2.46	0.41
1:A:469:LYS:O	1:A:470:ASP:CB	2.66	0.41
1:A:33[A]:LEU:HD23	1:A:306:ILE:HD12	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:564:ASN:HB3	5:B:802:HOH:O[3_654]	1.45	0.15

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	622/644 (97%)	609 (98%)	13 (2%)	0	100	100
1	B	602/644 (94%)	585 (97%)	17 (3%)	0	100	100
All	All	1224/1288 (95%)	1194 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	536/551 (97%)	533 (99%)	3 (1%)	78	75
1	B	520/551 (94%)	512 (98%)	8 (2%)	57	47
All	All	1056/1102 (96%)	1045 (99%)	11 (1%)	65	64

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	278	GLU
1	A	403	MET
1	A	411	GLU
1	B	65	ARG
1	B	215	MET
1	B	250	LYS
1	B	391	CYS
1	B	392	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	411	GLU
1	B	502	ASN
1	B	570	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	248	ASN
1	A	286	HIS
1	A	433	ASN
1	A	496	GLN
1	A	634	GLN
1	B	41	GLN
1	B	107	GLN
1	B	248	ASN
1	B	386	GLN
1	B	401	HIS
1	B	436	HIS
1	B	478	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2	15,15,15	1.24	1 (6%)	21,21,21	1.27	3 (14%)
2	BDP	C	2	2	12,12,13	1.28	1 (8%)	14,17,19	1.69	3 (21%)
2	NAG	C	3	2	14,14,15	2.74	4 (28%)	17,19,21	2.33	6 (35%)
2	NAG	D	1	2	15,15,15	0.83	0	21,21,21	1.49	2 (9%)
2	BDP	D	2	2	12,12,13	1.12	0	14,17,19	1.39	3 (21%)
2	NAG	D	3	2	14,14,15	2.68	2 (14%)	17,19,21	2.04	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2	-	0/6/26/26	0/1/1/1
2	BDP	C	2	2	-	0/4/21/24	0/1/1/1
2	NAG	C	3	2	-	3/6/23/26	0/1/1/1
2	NAG	D	1	2	-	0/6/26/26	0/1/1/1
2	BDP	D	2	2	-	0/4/21/24	0/1/1/1
2	NAG	D	3	2	-	2/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	NAG	C8-C7	-8.40	1.33	1.50
2	C	3	NAG	C8-C7	-7.97	1.34	1.50
2	D	3	NAG	O7-C7	4.56	1.33	1.23
2	C	3	NAG	O7-C7	4.10	1.32	1.23
2	C	1	NAG	C2-N2	3.22	1.50	1.45
2	C	3	NAG	C1-C2	2.49	1.55	1.52
2	C	3	NAG	O5-C5	2.15	1.47	1.43
2	C	2	BDP	O4-C4	2.04	1.48	1.43

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	NAG	C8-C7-N2	5.65	125.49	116.12
2	D	1	NAG	C4-C3-C2	-4.71	103.54	110.40
2	D	3	NAG	C8-C7-N2	4.35	123.33	116.12
2	C	3	NAG	O7-C7-N2	-3.99	114.92	121.98
2	C	2	BDP	C1-C2-C3	3.98	115.43	109.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	NAG	C1-O5-C5	3.68	117.12	112.19
2	D	3	NAG	O5-C1-C2	-3.57	105.77	111.29
2	C	3	NAG	O5-C1-C2	-3.16	106.41	111.29
2	C	2	BDP	C3-C4-C5	2.95	114.38	109.30
2	D	3	NAG	C4-C3-C2	-2.94	106.71	111.02
2	C	1	NAG	C1-O5-C5	2.86	119.19	113.65
2	D	2	BDP	C3-C4-C5	2.59	113.75	109.30
2	C	1	NAG	C2-N2-C7	-2.58	117.07	123.11
2	D	3	NAG	C2-N2-C7	2.53	126.29	122.90
2	C	3	NAG	O5-C5-C6	2.47	112.47	107.66
2	D	3	NAG	O3-C3-C2	2.41	114.40	109.40
2	D	2	BDP	C1-C2-C3	2.41	113.15	109.64
2	C	2	BDP	O6B-C6-C5	2.38	122.22	113.64
2	D	2	BDP	O6B-C6-C5	2.37	122.20	113.64
2	C	1	NAG	O1-C1-C2	2.35	114.09	109.22
2	D	1	NAG	C1-O5-C5	-2.16	109.47	113.65
2	C	3	NAG	C3-C4-C5	2.15	114.13	110.23
2	D	3	NAG	O7-C7-N2	-2.11	118.25	121.98
2	D	3	NAG	O7-C7-C8	-2.05	118.40	122.05

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	C	3	NAG	C8-C7-N2-C2
2	C	3	NAG	O7-C7-N2-C2
2	D	3	NAG	C8-C7-N2-C2
2	D	3	NAG	O7-C7-N2-C2
2	C	3	NAG	C1-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0
2	D	3	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	P4K	A	701	-	10,10,44	0.30	0	9,9,43	0.26	0
3	P4K	B	701	-	9,9,44	0.25	0	8,8,43	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	P4K	A	701	-	-	4/8/8/42	-
3	P4K	B	701	-	-	0/7/7/42	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	701	P4K	C16-C15-O8-C14
3	A	701	P4K	O9-C17-C18-O10
3	A	701	P4K	O7-C13-C14-O8
3	A	701	P4K	O8-C15-C16-O9

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	701	P4K	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	618/644 (95%)	-0.33	8 (1%) 75 82	15, 30, 54, 77	6 (0%)
1	B	603/644 (93%)	0.03	11 (1%) 67 74	8, 39, 63, 104	2 (0%)
All	All	1221/1288 (94%)	-0.15	19 (1%) 70 77	8, 35, 60, 104	8 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	585[A]	ASP	3.9
1	B	222	TRP	3.3
1	B	538	ARG	2.9
1	B	459	VAL	2.8
1	B	227	ILE	2.7
1	B	474	ILE	2.7
1	B	53	ALA	2.6
1	A	280	GLY	2.6
1	B	395	ALA	2.5
1	A	283	ALA	2.5
1	B	508	LYS	2.4
1	B	47	PRO	2.3
1	A	227	ILE	2.2
1	A	494	TYR	2.2
1	A	277	CYS	2.1
1	B	591	LYS	2.1
1	A	279	GLU	2.1
1	B	220	PHE	2.0
1	A	551	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	C	3	14/15	0.77	0.14	37,54,61,76	2
2	NAG	D	3	14/15	0.78	0.17	37,49,56,59	25
2	BDP	D	2	12/13	0.86	0.13	33,42,49,56	19
2	NAG	C	1	15/15	0.89	0.10	36,44,54,60	2
2	NAG	D	1	15/15	0.90	0.14	37,47,51,54	27
2	BDP	C	2	12/13	0.91	0.09	28,40,55,65	2

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	P4K	A	701	11/45	0.94	0.08	25,35,47,56	1
3	P4K	B	701	10/45	0.96	0.08	42,46,53,57	0
4	ZN	B	702	1/1	0.99	0.03	34,34,34,34	0
4	ZN	A	702	1/1	1.00	0.02	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around P4K A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.