



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 10:27 PM UTC

PDB ID : 1R8Y / pdb_00001r8y
Title : Crystal Structure of Mouse Glycine N-Methyltransferase (Monoclinic Form)
Authors : Pakhomova, S.; Luka, Z.; Wagner, C.; Newcomer, M.E.
Deposited on : 2003-10-28
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	292	
1	B	292	
1	C	292	
1	D	292	
1	E	292	
1	F	292	
1	G	292	
1	H	292	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BME	B	2057	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 17606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glycine N-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2192	1391	380	408	13			
1	B	282	Total	C	N	O	S	0	0	0
			2201	1399	382	407	13			
1	C	283	Total	C	N	O	S	0	0	0
			2175	1382	378	402	13			
1	D	286	Total	C	N	O	S	0	0	0
			2226	1417	382	414	13			
1	E	286	Total	C	N	O	S	0	0	0
			2216	1408	385	410	13			
1	F	282	Total	C	N	O	S	0	0	0
			2137	1357	370	397	13			
1	G	283	Total	C	N	O	S	0	0	0
			2217	1409	385	410	13			
1	H	282	Total	C	N	O	S	0	0	0
			2188	1391	379	405	13			

- Molecule 2 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			4	2	1	1		
2	A	1	Total	C	O	S	0	0
			4	2	1	1		
2	B	1	Total	C	O	S	0	0
			4	2	1	1		
2	D	1	Total	C	O	S	0	0
			4	2	1	1		
2	D	1	Total	C	O	S	0	0
			4	2	1	1		
2	D	1	Total	C	O	S	0	0
			4	2	1	1		
2	E	1	Total	C	O	S	0	0
			4	2	1	1		
2	F	1	Total	C	O	S	0	0
			4	2	1	1		
2	G	1	Total	C	O	S	0	0
			4	2	1	1		
2	G	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	4	Total	O	0	0
			4	4		
3	D	3	Total	O	0	0
			3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	2	Total	O	0	0
			2	2		
3	F	3	Total	O	0	0
			3	3		
3	G	2	Total	O	0	0
			2	2		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: glycine N-methyltransferase




- Molecule 1: glycine N-methyltransferase




- Molecule 1: glycine N-methyltransferase






- Molecule 1: glycine N-methyltransferase

Chain D: 63% 28% 7% ..



- Molecule 1: glycine N-methyltransferase

Chain E: 66% 24% 8% ..



- Molecule 1: glycine N-methyltransferase


Chain F: 66% 24% 6% ..





- Molecule 1: glycine N-methyltransferase

Chain G: 62% 27% 7% ..





• Molecule 1: glycine N-methyltransferase



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	107.98Å 108.30Å 119.01Å 90.00° 93.71° 90.00°	Depositor
Resolution (Å)	13.00 – 3.00	Depositor
% Data completeness (in resolution range)	74.2 (13.00-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.222 , 0.285	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	17606	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.03	5/2244 (0.2%)	1.20	9/3047 (0.3%)
1	B	1.00	3/2253 (0.1%)	1.16	9/3058 (0.3%)
1	C	0.93	2/2228 (0.1%)	1.10	4/3029 (0.1%)
1	D	1.02	0/2280	1.16	7/3096 (0.2%)
1	E	0.95	0/2269	1.14	7/3081 (0.2%)
1	F	0.96	2/2189 (0.1%)	1.14	7/2978 (0.2%)
1	G	1.05	3/2269 (0.1%)	1.20	9/3076 (0.3%)
1	H	0.95	1/2240 (0.0%)	1.16	8/3043 (0.3%)
All	All	0.99	16/17972 (0.1%)	1.16	60/24408 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	2
1	E	0	1
1	G	0	1
1	H	0	1
All	All	0	6

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	38	ARG	CA-C	8.86	1.64	1.52
1	B	37	THR	CA-CB	8.07	1.67	1.53
1	G	38	ARG	CA-CB	6.66	1.58	1.53
1	G	266	VAL	CA-CB	6.46	1.61	1.54
1	F	4	VAL	CA-C	6.35	1.60	1.52

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	198	LEU	N-CA-C	-9.77	93.71	108.46
1	C	38	ARG	N-CA-C	8.82	123.65	111.74
1	C	2	ASP	N-CA-C	7.57	122.71	107.41
1	A	37	THR	CB-CA-C	7.17	122.31	111.70
1	E	2	ASP	N-CA-C	7.03	120.82	107.75

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	38	ARG	Peptide
1	C	36	ASP	Peptide
1	C	38	ARG	Peptide
1	E	1	VAL	Peptide
1	G	38	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2192	0	2106	80	0
1	B	2201	0	2121	89	0
1	C	2175	0	2063	47	0
1	D	2226	0	2156	67	0
1	E	2216	0	2134	64	0
1	F	2137	0	1985	52	0
1	G	2217	0	2158	52	0
1	H	2188	0	2103	58	0
2	A	8	0	10	3	0
2	B	4	0	5	5	0
2	D	12	0	15	1	0
2	E	4	0	5	0	0
2	F	4	0	5	0	0
2	G	8	0	10	2	0
3	C	4	0	0	1	0
3	D	3	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	2	0	0	3	0
3	F	3	0	0	1	0
3	G	2	0	0	0	0
All	All	17606	0	16876	449	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 449 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:4:VAL:HG12	1:H:5:TYR:H	1.15	1.09
1:A:264:HIS:ND1	1:A:288:LEU:HD21	1.71	1.05
1:A:37:THR:O	1:A:38:ARG:HG3	1.62	0.99
1:C:20:GLN:HG2	3:C:293:HOH:O	1.66	0.96
1:E:20:GLN:HG2	3:E:5247:HOH:O	1.67	0.94

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/292 (95%)	250 (90%)	22 (8%)	5 (2%)	6	31
1	B	278/292 (95%)	250 (90%)	23 (8%)	5 (2%)	6	31
1	C	279/292 (96%)	253 (91%)	20 (7%)	6 (2%)	5	26
1	D	282/292 (97%)	258 (92%)	21 (7%)	3 (1%)	11	43
1	E	282/292 (97%)	253 (90%)	24 (8%)	5 (2%)	6	31
1	F	278/292 (95%)	245 (88%)	24 (9%)	9 (3%)	3	18
1	G	279/292 (96%)	250 (90%)	24 (9%)	5 (2%)	6	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	278/292 (95%)	252 (91%)	22 (8%)	4 (1%)	9	36
All	All	2233/2336 (96%)	2011 (90%)	180 (8%)	42 (2%)	6	30

5 of 42 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	ARG
1	A	20	GLN
1	B	20	GLN
1	B	39	SER
1	C	5	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/246 (94%)	198 (86%)	32 (14%)	3	17
1	B	230/246 (94%)	193 (84%)	37 (16%)	2	12
1	C	222/246 (90%)	193 (87%)	29 (13%)	4	19
1	D	236/246 (96%)	194 (82%)	42 (18%)	2	10
1	E	231/246 (94%)	191 (83%)	40 (17%)	2	10
1	F	211/246 (86%)	179 (85%)	32 (15%)	3	14
1	G	235/246 (96%)	197 (84%)	38 (16%)	2	12
1	H	228/246 (93%)	191 (84%)	37 (16%)	2	12
All	All	1823/1968 (93%)	1536 (84%)	287 (16%)	2	13

5 of 287 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	162	SER
1	H	291	THR
1	G	205	SER
1	H	69	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	287	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	245	HIS
1	H	159	ASN
1	F	116	ASN
1	H	245	HIS
1	G	277	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BME	G	7282	1	3,3,3	0.48	0	2,2,2	0.83	0
2	BME	A	1262	1	3,3,3	0.56	0	2,2,2	0.49	0
2	BME	F	6282	1	3,3,3	0.35	0	2,2,2	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BME	D	4262	1	3,3,3	0.32	0	2,2,2	0.25	0
2	BME	E	5246	1	3,3,3	0.37	0	2,2,2	0.54	0
2	BME	D	4185	1	3,3,3	0.79	0	2,2,2	1.48	1 (50%)
2	BME	D	4246	1	3,3,3	0.27	0	2,2,2	0.73	0
2	BME	B	2057	1	3,3,3	0.47	0	2,2,2	0.45	0
2	BME	G	7246	1	3,3,3	0.76	0	2,2,2	0.92	0
2	BME	A	1282	1	3,3,3	0.91	0	2,2,2	1.78	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BME	G	7282	1	-	0/1/1/1	-
2	BME	A	1262	1	-	1/1/1/1	-
2	BME	F	6282	1	-	0/1/1/1	-
2	BME	D	4262	1	-	0/1/1/1	-
2	BME	E	5246	1	-	1/1/1/1	-
2	BME	D	4185	1	-	1/1/1/1	-
2	BME	D	4246	1	-	0/1/1/1	-
2	BME	B	2057	1	-	0/1/1/1	-
2	BME	G	7246	1	-	1/1/1/1	-
2	BME	A	1282	1	-	1/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1282	BME	O1-C1-C2	2.04	118.80	110.82
2	D	4185	BME	O1-C1-C2	2.03	118.74	110.82

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1262	BME	O1-C1-C2-S2
2	D	4185	BME	O1-C1-C2-S2
2	E	5246	BME	O1-C1-C2-S2
2	A	1282	BME	O1-C1-C2-S2
2	G	7246	BME	O1-C1-C2-S2

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	7282	BME	2	0
2	A	1262	BME	2	0
2	D	4185	BME	1	0
2	B	2057	BME	5	0
2	A	1282	BME	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.