



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 11:03 PM UTC

PDB ID : 1R9N / pdb_00001r9n
Title : Crystal Structure of human dipeptidyl peptidase IV in complex with a decapeptide (tNPY) at 2.3 Ang. Resolution
Authors : Aertgeerts, K.; Ye, S.; Tennant, M.G.; Collins, B.; Rogers, J.; Sang, B.-C.; Skene, R.; Webb, D.R.; Prasad, G.S.
Deposited on : 2003-10-30
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	739	
1	B	739	
1	C	739	
1	D	739	
2	E	10	
2	F	10	
2	G	10	
2	H	10	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	2	 100%
3	J	2	 50% 50%
3	K	2	 100%
3	L	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	2811	X	-	-	-
4	NAG	A	6851	X	-	-	-
4	NAG	D	2811	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 25627 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	728	Total	C	N	O	S	0	0	0
			5963	3827	982	1128	26			
1	B	731	Total	C	N	O	S	0	0	0
			5993	3845	991	1131	26			
1	C	727	Total	C	N	O	S	0	0	0
			5957	3824	981	1126	26			
1	D	727	Total	C	N	O	S	0	0	0
			5957	3824	981	1126	26			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	ALA	-	cloning artifact	UNP P27487
A	29	ASP	-	cloning artifact	UNP P27487
A	30	PRO	-	cloning artifact	UNP P27487
A	31	GLY	-	cloning artifact	UNP P27487
A	32	GLY	-	cloning artifact	UNP P27487
A	33	SER	-	cloning artifact	UNP P27487
A	34	HIS	-	expression tag	UNP P27487
A	35	HIS	-	expression tag	UNP P27487
A	36	HIS	-	expression tag	UNP P27487
A	37	HIS	-	expression tag	UNP P27487
A	38	HIS	-	expression tag	UNP P27487
B	28	ALA	-	cloning artifact	UNP P27487
B	29	ASP	-	cloning artifact	UNP P27487
B	30	PRO	-	cloning artifact	UNP P27487
B	31	GLY	-	cloning artifact	UNP P27487
B	32	GLY	-	cloning artifact	UNP P27487
B	33	SER	-	cloning artifact	UNP P27487
B	34	HIS	-	expression tag	UNP P27487
B	35	HIS	-	expression tag	UNP P27487
B	36	HIS	-	expression tag	UNP P27487
B	37	HIS	-	expression tag	UNP P27487

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	38	HIS	-	expression tag	UNP P27487
C	28	ALA	-	cloning artifact	UNP P27487
C	29	ASP	-	cloning artifact	UNP P27487
C	30	PRO	-	cloning artifact	UNP P27487
C	31	GLY	-	cloning artifact	UNP P27487
C	32	GLY	-	cloning artifact	UNP P27487
C	33	SER	-	cloning artifact	UNP P27487
C	34	HIS	-	expression tag	UNP P27487
C	35	HIS	-	expression tag	UNP P27487
C	36	HIS	-	expression tag	UNP P27487
C	37	HIS	-	expression tag	UNP P27487
C	38	HIS	-	expression tag	UNP P27487
D	28	ALA	-	cloning artifact	UNP P27487
D	29	ASP	-	cloning artifact	UNP P27487
D	30	PRO	-	cloning artifact	UNP P27487
D	31	GLY	-	cloning artifact	UNP P27487
D	32	GLY	-	cloning artifact	UNP P27487
D	33	SER	-	cloning artifact	UNP P27487
D	34	HIS	-	expression tag	UNP P27487
D	35	HIS	-	expression tag	UNP P27487
D	36	HIS	-	expression tag	UNP P27487
D	37	HIS	-	expression tag	UNP P27487
D	38	HIS	-	expression tag	UNP P27487

- Molecule 2 is a protein called Neuropeptide Y.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	4	Total	C	N	O	0	0	0
			34	23	5	6			
2	F	6	Total	C	N	O	0	0	0
			49	32	7	10			
2	G	4	Total	C	N	O	0	0	0
			34	23	5	6			
2	H	5	Total	C	N	O	0	0	0
			41	28	6	7			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	J	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	K	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	L	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	342	Total	O	0	0
			342	342		
5	E	3	Total	O	0	0
			3	3		
5	B	361	Total	O	0	0
			361	361		
5	F	4	Total	O	0	0
			4	4		
5	C	277	Total	O	0	0
			277	277		
5	G	6	Total	O	0	0
			6	6		
5	D	254	Total	O	0	0
			254	254		
5	H	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: Dipeptidyl peptidase IV


Chain A: 



• Molecule 1: Dipeptidyl peptidase IV


Chain B: 






• Molecule 1: Dipeptidyl peptidase IV

Chain C: 86% 11% ..




• Molecule 1: Dipeptidyl peptidase IV

Chain D: 77% 19% ..



• Molecule 2: Neuropeptide Y

Chain E: 40% 60%



• Molecule 2: Neuropeptide Y


Chain F: 20% 40% 40%



● Molecule 2: Neuropeptide Y



● Molecule 2: Neuropeptide Y



● Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




● Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



● Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



● Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	122.61Å 122.70Å 145.41Å 90.00° 114.88° 90.00°	Depositor
Resolution (Å)	41.17 – 2.30	Depositor
% Data completeness (in resolution range)	100.0 (41.17-2.30)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	0.05	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.211 , 0.251	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	25627	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.63	3/6135 (0.0%)	0.85	3/8344 (0.0%)
1	B	0.61	0/6168	0.85	3/8389 (0.0%)
1	C	0.61	3/6129 (0.0%)	0.83	1/8336 (0.0%)
1	D	0.60	2/6129 (0.0%)	0.82	3/8336 (0.0%)
2	E	0.77	0/35	0.79	0/46
2	F	0.70	0/51	1.00	0/69
2	G	0.55	0/35	0.91	0/46
2	H	0.57	0/43	0.74	0/58
All	All	0.61	8/24725 (0.0%)	0.84	10/33624 (0.0%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	262	VAL	CA-CB	6.24	1.61	1.54
1	A	262	VAL	CA-CB	5.99	1.60	1.54
1	C	594	ILE	CA-CB	5.92	1.62	1.54
1	D	728	VAL	CA-CB	5.36	1.60	1.54
1	A	688	VAL	CA-CB	5.36	1.61	1.54

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	202	VAL	CB-CA-C	-6.66	103.02	112.22
1	A	202	VAL	CB-CA-C	-6.56	103.28	112.14
1	D	297	ASP	N-CA-CB	6.42	119.54	110.36
1	D	454	CYS	N-CA-C	6.08	118.84	108.02
1	D	202	VAL	N-CA-C	5.83	116.56	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5963	0	5677	87	0
1	B	5993	0	5700	74	0
1	C	5957	0	5674	48	0
1	D	5957	0	5677	75	0
2	E	34	0	33	8	0
2	F	49	0	44	5	0
2	G	34	0	33	5	0
2	H	41	0	40	3	0
3	I	28	0	25	0	0
3	J	28	0	25	0	0
3	K	28	0	25	0	0
3	L	28	0	25	0	0
4	A	84	0	78	3	0
4	B	70	0	65	1	0
4	C	56	0	52	0	0
4	D	28	0	26	0	0
5	A	342	0	0	0	0
5	B	361	0	0	0	0
5	C	277	0	0	2	0
5	D	254	0	0	0	0
5	E	3	0	0	0	0
5	F	4	0	0	0	0
5	G	6	0	0	0	0
5	H	2	0	0	0	0
All	All	25627	0	23199	286	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 286 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:ARG:HH21	1:B:253:ARG:HH21	1.08	0.97
1:C:711:VAL:HG13	1:C:740:HIS:CE1	2.03	0.93
1:A:711:VAL:HG13	1:A:740:HIS:CE1	2.04	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:589:LYS:HD3	5:C:5477:HOH:O	1.70	0.90
1:A:74:ASN:HD21	1:A:92:ASN:HD22	1.17	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	726/739 (98%)	695 (96%)	29 (4%)	2 (0%)	36	46
1	B	729/739 (99%)	695 (95%)	34 (5%)	0	100	100
1	C	725/739 (98%)	688 (95%)	33 (5%)	4 (1%)	21	27
1	D	725/739 (98%)	683 (94%)	41 (6%)	1 (0%)	48	60
2	E	2/10 (20%)	2 (100%)	0	0	100	100
2	F	4/10 (40%)	3 (75%)	1 (25%)	0	100	100
2	G	2/10 (20%)	1 (50%)	1 (50%)	0	100	100
2	H	3/10 (30%)	3 (100%)	0	0	100	100
All	All	2916/2996 (97%)	2770 (95%)	139 (5%)	7 (0%)	43	55

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	C	85	ASN
1	A	630	SER
1	C	74	ASN
1	D	630	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	653/661 (99%)	609 (93%)	44 (7%)	15	21
1	B	656/661 (99%)	611 (93%)	45 (7%)	14	20
1	C	652/661 (99%)	616 (94%)	36 (6%)	19	28
1	D	652/661 (99%)	600 (92%)	52 (8%)	11	15
2	E	4/9 (44%)	4 (100%)	0	100	100
2	F	6/9 (67%)	6 (100%)	0	100	100
2	G	4/9 (44%)	4 (100%)	0	100	100
2	H	5/9 (56%)	5 (100%)	0	100	100
All	All	2632/2680 (98%)	2455 (93%)	177 (7%)	15	21

5 of 177 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	594	ILE
1	D	327	ILE
1	C	673	LEU
1	D	90	LEU
1	D	377	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 55 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	757	HIS
1	C	344	GLN
1	D	612	GLN
1	D	314	GLN
1	C	51	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	I	1	1,3	14,14,15	0.65	0	17,19,21	1.29	2 (11%)
3	NAG	I	2	3	14,14,15	0.54	0	17,19,21	1.33	3 (17%)
3	NAG	J	1	1,3	14,14,15	0.60	0	17,19,21	1.11	1 (5%)
3	NAG	J	2	3	14,14,15	0.51	0	17,19,21	0.92	0
3	NAG	K	1	1,3	14,14,15	0.63	0	17,19,21	1.01	1 (5%)
3	NAG	K	2	3	14,14,15	0.59	0	17,19,21	1.78	5 (29%)
3	NAG	L	1	1,3	14,14,15	0.54	0	17,19,21	1.16	2 (11%)
3	NAG	L	2	3	14,14,15	0.50	0	17,19,21	1.36	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	I	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	I	2	3	-	2/6/23/26	0/1/1/1
3	NAG	J	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	J	2	3	-	4/6/23/26	0/1/1/1
3	NAG	K	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	K	2	3	-	2/6/23/26	0/1/1/1
3	NAG	L	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	L	2	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	K	2	NAG	C3-C4-C5	4.21	117.86	110.23
3	I	1	NAG	C1-O5-C5	3.35	116.67	112.19
3	I	2	NAG	C1-O5-C5	3.03	116.24	112.19
3	K	2	NAG	C8-C7-N2	2.82	120.79	116.12
3	L	2	NAG	C2-N2-C7	2.65	126.45	122.90

There are no chirality outliers.


5 of 13 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	J	2	NAG	C8-C7-N2-C2
3	J	2	NAG	O7-C7-N2-C2
3	I	1	NAG	C4-C5-C6-O6
3	I	1	NAG	O5-C5-C6-O6
3	K	2	NAG	C8-C7-N2-C2


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	2811	1	14,14,15	0.96	1 (7%)	17,19,21	1.13	3 (17%)
4	NAG	C	3211	1	14,14,15	0.52	0	17,19,21	2.02	4 (23%)
4	NAG	A	1501	1	14,14,15	0.70	0	17,19,21	2.28	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	1501	1	14,14,15	0.52	0	17,19,21	2.13	5 (29%)
4	NAG	D	2811	1	14,14,15	0.51	0	17,19,21	1.47	3 (17%)
4	NAG	A	2811	1	14,14,15	0.57	0	17,19,21	2.16	2 (11%)
4	NAG	A	3211	1	14,14,15	0.58	0	17,19,21	1.02	1 (5%)
4	NAG	D	2191	1	14,14,15	0.55	0	17,19,21	0.95	1 (5%)
4	NAG	C	2191	1	14,14,15	0.71	0	17,19,21	1.14	1 (5%)
4	NAG	B	2811	1	14,14,15	0.47	0	17,19,21	1.18	1 (5%)
4	NAG	B	851	1	14,14,15	0.63	0	17,19,21	1.93	2 (11%)
4	NAG	C	5201	1	14,14,15	0.64	0	17,19,21	1.75	2 (11%)
4	NAG	A	6851	1	14,14,15	0.73	1 (7%)	17,19,21	2.30	3 (17%)
4	NAG	A	2191	1	14,14,15	0.55	0	17,19,21	1.07	1 (5%)
4	NAG	A	851	1	14,14,15	0.51	0	17,19,21	1.15	1 (5%)
4	NAG	B	3211	1	14,14,15	0.51	0	17,19,21	1.61	1 (5%)
4	NAG	B	2191	1	14,14,15	0.54	0	17,19,21	0.97	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	2811	1	-	2/6/23/26	0/1/1/1
4	NAG	C	3211	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1501	1	-	3/6/23/26	0/1/1/1
4	NAG	D	2811	1	1/1/5/7	4/6/23/26	0/1/1/1
4	NAG	B	1501	1	-	2/6/23/26	0/1/1/1
4	NAG	A	2811	1	1/1/5/7	5/6/23/26	0/1/1/1
4	NAG	A	3211	1	-	2/6/23/26	0/1/1/1
4	NAG	D	2191	1	-	2/6/23/26	0/1/1/1
4	NAG	C	2191	1	-	2/6/23/26	0/1/1/1
4	NAG	B	2811	1	-	3/6/23/26	0/1/1/1
4	NAG	B	851	1	-	2/6/23/26	0/1/1/1
4	NAG	C	5201	1	-	4/6/23/26	0/1/1/1
4	NAG	A	6851	1	1/1/5/7	3/6/23/26	0/1/1/1
4	NAG	A	2191	1	-	0/6/23/26	0/1/1/1
4	NAG	A	851	1	-	3/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	3211	1	-	0/6/23/26	0/1/1/1
4	NAG	B	2191	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	2811	NAG	O6-C6	2.51	1.53	1.42
4	A	6851	NAG	C1-C2	2.14	1.55	1.52

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2811	NAG	C1-O5-C5	7.59	122.36	112.19
4	B	851	NAG	C1-O5-C5	6.16	120.45	112.19
4	A	6851	NAG	C1-O5-C5	6.08	120.34	112.19
4	C	5201	NAG	C1-O5-C5	6.05	120.30	112.19
4	C	3211	NAG	C1-O5-C5	5.84	120.01	112.19

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	2811	NAG	C1
4	A	6851	NAG	C1
4	D	2811	NAG	C1

5 of 41 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	6851	NAG	C3-C2-N2-C7
4	B	851	NAG	C8-C7-N2-C2
4	B	851	NAG	O7-C7-N2-C2
4	B	2811	NAG	C8-C7-N2-C2
4	B	2811	NAG	O7-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	851	NAG	1	0
4	A	6851	NAG	2	0
4	A	851	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.