



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:07 AM UTC

PDB ID : 5R9Z / pdb_00005r9z
Title : PanDDA analysis group deposition Form1 MAP kinase p38-alpha – Fragment
N13502a in complex with MAP kinase p38-alpha
Authors : De Nicola, G.F.; Nichols, C.E.
Deposited on : 2020-03-04
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	30% 89% 7%

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 3058 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	348	Total	C	N	O	S	0	4	0
			2781	1781	469	516	15			

- Molecule 2 is 2-[(4-chlorophenyl)amino]benzamide (CCD ID: AYS) (formula: C₁₃H₁₁ClN₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			17	13	1	2	1		


- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	Cl	0	0
			5	5		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	240	Total	O	0	0
			240	240		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 14



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.81Å 86.51Å 127.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.83 – 1.66 29.83 – 1.66	Depositor EDS
% Data completeness (in resolution range)	98.7 (29.83-1.66) 98.8 (29.83-1.66)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 1.66Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.178 , 0.197 0.227 , 0.227	Depositor DCC
R_{free} test set	3019 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3058	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AYS, SO4, DMS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.31	6/2847 (0.2%)	1.22	3/3873 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	116	ILE	C-N	8.67	1.44	1.33
1	A	102	VAL	C-O	5.41	1.29	1.24
1	A	227	ASP	C-O	5.38	1.29	1.23
1	A	234	LEU	C-O	5.37	1.30	1.24
1	A	228	HIS	CE1-NE2	5.29	1.37	1.32
1	A	128	GLN	C-O	5.06	1.29	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	294	ASP	CA-CB-CG	5.32	117.92	112.60
1	A	264	GLN	CB-CA-C	5.31	118.73	109.65
1	A	223	PHE	CA-CB-CG	5.10	118.90	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2781	0	2702	6	0
2	A	17	0	0	0	0
3	A	5	0	0	0	0
4	A	1	0	0	0	0
5	A	10	0	0	0	0
6	A	4	0	6	0	0
7	A	240	0	0	0	0
All	All	3058	0	2708	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ILE:HD13	1:A:109:MET:HE3	1.64	0.80
1:A:84:ILE:CD1	1:A:109:MET:HE3	2.29	0.62
1:A:201:ASN:ND2	1:A:203:THR:OG1	2.44	0.48
1:A:78[B]:MET:HE3	1:A:169:PHE:CZ	2.54	0.43
1:A:147:ILE:HG22	1:A:149[B]:ARG:HG3	2.02	0.41
1:A:203:THR:OG1	1:A:293:SER:HB2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/360 (97%)	345 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	295/319 (92%)	283 (96%)	12 (4%)	27 6

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	30	VAL
1	A	37	SER
1	A	63	ILE
1	A	113	LEU
1	A	116	ILE
1	A	167	LEU
1	A	193	ILE
1	A	201	ASN
1	A	247	LEU
1	A	254	SER
1	A	352	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	A	115	ASN
1	A	128	GLN
1	A	174	HIS
1	A	201	ASN
1	A	260	GLN
1	A	272	ASN
1	A	310	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	409	-	4,4,4	0.35	0	6,6,6	0.05	0
6	DMS	A	410	-	3,3,3	0.21	0	3,3,3	0.10	0
5	SO4	A	408	-	4,4,4	0.26	0	6,6,6	0.28	0
2	AYS	A	401	-	18,18,18	1.47	4 (22%)	24,24,24	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AYS	A	401	-	-	0/8/8/8	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	AYS	C7-N1	3.47	1.39	1.33
2	A	401	AYS	C3-C4	3.02	1.44	1.39
2	A	401	AYS	C9-C8	2.13	1.42	1.39
2	A	401	AYS	C13-C8	2.02	1.42	1.39

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	348/360 (96%)	2.20	108 (31%) 1 1	4, 22, 41, 63	78 (22%)

All (108) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	117	VAL	15.1
1	A	34	ALA	13.4
1	A	116	ILE	13.4
1	A	30	VAL	13.3
1	A	239	VAL	12.2
1	A	255	ALA	11.7
1	A	172	ALA	11.1
1	A	250	ILE	11.1
1	A	258	TYR	10.7
1	A	122	LEU	10.6
1	A	35	TYR	10.6
1	A	193	ILE	10.4
1	A	311	TYR	10.4
1	A	254	SER	10.3
1	A	175	THR	10.2
1	A	244	ALA	10.2
1	A	241	THR	9.9
1	A	195	LEU	9.8
1	A	171	LEU	9.6
1	A	252	SER	9.5
1	A	50	VAL	9.4
1	A	242	PRO	8.9
1	A	33	GLY	8.8
1	A	119	CYS	8.8
1	A	115	ASN	8.7
1	A	190	ALA	8.7
1	A	31	GLY	8.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	185	THR	8.6
1	A	240	GLY	8.5
1	A	247	LEU	8.4
1	A	197	TRP	8.3
1	A	246	LEU	8.3
1	A	118	LYS	8.3
1	A	79	LYS	8.1
1	A	243	GLY	8.1
1	A	78[A]	MET	8.0
1	A	313	ASP	7.9
1	A	45	LYS	7.9
1	A	312	HIS	7.8
1	A	196	ASN	7.7
1	A	201	ASN	7.7
1	A	32	SER	7.7
1	A	180	THR	7.7
1	A	82	ASN	7.7
1	A	80	HIS	7.7
1	A	191	PRO	7.4
1	A	245	GLU	7.4
1	A	120	GLN	7.3
1	A	257	ASN	7.2
1	A	177	ASP	7.1
1	A	114	ASN	7.1
1	A	121	LYS	7.0
1	A	249	LYS	6.9
1	A	194	MET	6.9
1	A	125[A]	ASP	6.9
1	A	149[A]	ARG	6.7
1	A	288[A]	MET	6.6
1	A	36	GLY	6.6
1	A	256	ARG	6.5
1	A	176	ASP	6.4
1	A	200	TYR	6.3
1	A	37	SER	6.3
1	A	173	ARG	6.3
1	A	202	GLN	6.0
1	A	81	GLU	6.0
1	A	251	SER	5.9
1	A	248	LYS	5.9
1	A	179	MET	5.5
1	A	253	GLU	5.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	174	HIS	5.5
1	A	310	GLN	5.4
1	A	189	ARG	5.4
1	A	192	GLU	5.3
1	A	186	ARG	5.1
1	A	170	GLY	5.0
1	A	198	MET	4.9
1	A	199	HIS	4.7
1	A	56	SER	4.4
1	A	178	GLU	4.0
1	A	162	CYS	4.0
1	A	48	HIS	3.9
1	A	16	THR	3.7
1	A	352	PRO	3.5
1	A	13	LEU	3.2
1	A	64	HIS	3.2
1	A	60	GLN	3.2
1	A	159	ASN	3.1
1	A	98	GLU	2.8
1	A	97	GLU	2.8
1	A	100	ASN	2.7
1	A	18	TRP	2.6
1	A	160	GLU	2.5
1	A	161	ASP	2.5
1	A	62	ILE	2.5
1	A	14	ASN	2.4
1	A	111	ALA	2.3
1	A	63	ILE	2.3
1	A	42	PHE	2.2
1	A	46	THR	2.2
1	A	17	ILE	2.1
1	A	346	ILE	2.1
1	A	331	ASP	2.1
1	A	334	ILE	2.1
1	A	109	MET	2.0
1	A	260	GLN	2.0
1	A	113	LEU	2.0
1	A	107	HIS	2.0
1	A	57	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	AYS	A	401	17/17	0.66	0.21	22,23,25,29	17
5	SO4	A	408	5/5	0.79	0.19	53,62,68,72	0
6	DMS	A	410	4/4	0.80	0.24	67,69,75,81	0
3	CL	A	404	1/1	0.92	0.12	48,48,48,48	0
5	SO4	A	409	5/5	0.93	0.14	28,33,34,35	5
3	CL	A	403	1/1	0.95	0.15	43,43,43,43	0
3	CL	A	402	1/1	0.97	0.26	46,46,46,46	0
3	CL	A	405	1/1	0.98	0.21	39,39,39,39	0
3	CL	A	406	1/1	0.98	0.10	35,35,35,35	0
4	MG	A	407	1/1	0.98	0.17	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.