



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:39 PM UTC

PDB ID : 6R9Q / pdb_00006r9q
Title : Structure of *Saccharomyces cerevisiae* apo Pan2 pseudoubiquitin hydrolase-RNA exonuclease (UCH-Exo) module in complex with AACCAA RNA
Authors : Tang, T.T.L.; Stowell, J.A.W.; Hill, C.H.; Passmore, L.A.
Deposited on : 2019-04-03
Resolution : 3.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2010 (3.10-3.06)
Clashscore	190562	2102 (3.10-3.06)
Ramachandran outliers	187476	1982 (3.10-3.06)
Sidechain outliers	187428	1981 (3.10-3.06)
RSRZ outliers	180081	2010 (3.10-3.06)
RNA backbone	3983	1054 (3.30-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	672	
2	B	6	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAN2-PAN3 deadenylation complex catalytic subunit PAN2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	610	Total	C	N	O	S	0	0	0
			4590	2930	757	885	18			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	444	MET	-	initiating methionine	UNP P53010
A	445	HIS	-	expression tag	UNP P53010
A	446	HIS	-	expression tag	UNP P53010
A	447	HIS	-	expression tag	UNP P53010
A	448	HIS	-	expression tag	UNP P53010
A	449	HIS	-	expression tag	UNP P53010
A	450	HIS	-	expression tag	UNP P53010
A	451	HIS	-	expression tag	UNP P53010
A	452	HIS	-	expression tag	UNP P53010
A	453	LEU	-	expression tag	UNP P53010
A	454	GLU	-	expression tag	UNP P53010
A	455	VAL	-	expression tag	UNP P53010
A	456	LEU	-	expression tag	UNP P53010
A	457	PHE	-	expression tag	UNP P53010
A	458	GLN	-	expression tag	UNP P53010
A	459	GLY	-	expression tag	UNP P53010
A	460	PRO	-	expression tag	UNP P53010
A	912	ALA	GLU	engineered mutation	UNP P53010


- Molecule 2 is a RNA chain called AACCAA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	4	Total	C	N	O	P	0	0	0
			81	38	16	24	3			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PAN2-PAN3 deadenylation complex catalytic subunit PAN2



- Molecule 2: AACCAA



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.19Å 117.32Å 256.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.49 – 3.08 72.49 – 3.08	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.49-3.08) 99.7 (72.49-3.08)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 3.07Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.261 , 0.297 (Not available) , 0.283	Depositor DCC
R_{free} test set	1298 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	107.4	Xtriage
Anisotropy	0.460	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 93.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4671	wwPDB-VP
Average B, all atoms (Å ²)	125.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.09	0/4684	0.27	0/6400
2	B	0.37	0/90	0.59	0/138
All	All	0.10	0/4774	0.28	0/6538

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4590	0	4177	63	0
2	B	81	0	46	5	0
All	All	4671	0	4223	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:975:TYR:CE1	2:B:6:A:N6	2.43	0.86
1:A:983:LEU:HG	1:A:985:PRO:HD2	1.69	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:866:ASP:OD2	1:A:1007:GLN:NE2	2.25	0.70
1:A:956:PRO:HG3	1:A:1069:ILE:HD11	1.76	0.67
1:A:537:LEU:O	1:A:1000:ARG:NH1	2.29	0.66
1:A:855:ALA:HA	1:A:859:ARG:HH21	1.60	0.66
1:A:874:ILE:HD13	1:A:1000:ARG:HD3	1.78	0.65
1:A:756:LEU:HD21	1:A:789:LEU:HD11	1.78	0.65
1:A:465:GLN:HG2	1:A:828:LEU:HD21	1.82	0.62
1:A:569:ASN:H	1:A:1025:ASN:HD22	1.46	0.61
1:A:527:PRO:O	1:A:531:ASN:ND2	2.33	0.61
1:A:1013:VAL:HG12	1:A:1034:ARG:HB2	1.83	0.60
2:B:3:C:H6	2:B:3:C:HO5'	1.51	0.58
1:A:523:TYR:OH	1:A:623:ASN:OD1	2.23	0.56
1:A:894:LEU:O	1:A:946:ARG:NH2	2.39	0.56
1:A:1015:HIS:HD2	1:A:1049:LEU:HB2	1.70	0.56
1:A:963:VAL:HG22	1:A:994:ARG:HA	1.89	0.55
1:A:858:LEU:HD23	1:A:858:LEU:O	2.08	0.54
1:A:962:VAL:HG22	1:A:998:VAL:HG11	1.91	0.53
1:A:929:ARG:HH12	1:A:933:ARG:HA	1.72	0.53
1:A:569:ASN:H	1:A:1025:ASN:ND2	2.05	0.53
1:A:513:ASN:OD1	1:A:571:GLN:NE2	2.33	0.52
1:A:707:THR:HG23	1:A:724:GLN:HB3	1.92	0.52
1:A:825:ASN:HB3	1:A:828:LEU:HB3	1.92	0.52
1:A:941:ARG:NH1	1:A:961:TYR:OH	2.43	0.52
1:A:756:LEU:HD13	1:A:840:MET:HE1	1.92	0.50
1:A:919:GLU:HA	1:A:934:PRO:HB3	1.94	0.50
1:A:635:ALA:O	1:A:640:HIS:ND1	2.37	0.50
1:A:1015:HIS:CD2	1:A:1049:LEU:HB2	2.47	0.50
1:A:479:TYR:HD1	1:A:806:VAL:HG11	1.77	0.49
1:A:736:SER:HA	1:A:851:ILE:HD13	1.93	0.49
1:A:895:LEU:HD23	1:A:946:ARG:HH21	1.78	0.49
1:A:639:ASN:O	1:A:639:ASN:ND2	2.46	0.48
1:A:945:ILE:HA	1:A:956:PRO:HA	1.95	0.47
1:A:522:LEU:HD22	1:A:793:VAL:HG23	1.96	0.47
1:A:749:ILE:HD11	1:A:755:TRP:NE1	2.30	0.47
1:A:1042:GLN:HG2	1:A:1099:TYR:HE2	1.79	0.46
1:A:624:ARG:NH1	1:A:738:GLU:OE1	2.48	0.46
1:A:476:PRO:HG2	1:A:479:TYR:CD2	2.51	0.45
1:A:574:LEU:HD11	1:A:625:PHE:HE2	1.82	0.45
1:A:471:ASN:HD21	1:A:473:TYR:HB2	1.82	0.45
1:A:476:PRO:HG2	1:A:479:TYR:HD2	1.80	0.45
1:A:549:ASP:OD2	1:A:573:SER:OG	2.25	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1096:ASN:HA	1:A:1099:TYR:CD1	2.51	0.45
1:A:975:TYR:CZ	2:B:6:A:N6	2.83	0.45
1:A:979:LEU:O	1:A:981:GLY:N	2.46	0.45
1:A:855:ALA:CA	1:A:859:ARG:HH21	2.29	0.44
1:A:855:ALA:O	1:A:859:ARG:NE	2.43	0.44
2:B:3:C:H6	2:B:3:C:O5'	1.98	0.44
1:A:908:ALA:HB3	1:A:945:ILE:HB	2.00	0.44
1:A:578:THR:HA	1:A:592:LEU:HD12	1.99	0.43
1:A:708:GLN:HG3	1:A:723:THR:HG22	2.01	0.43
1:A:1099:TYR:O	1:A:1103:ARG:HG2	2.18	0.43
1:A:935:LYS:HE2	1:A:935:LYS:HB2	1.89	0.42
1:A:647:CYS:O	1:A:732:PRO:HB2	2.19	0.42
1:A:529:MET:O	1:A:533:VAL:HG12	2.20	0.42
1:A:542:PHE:HZ	1:A:874:ILE:HG12	1.84	0.42
1:A:872:TYR:CD2	1:A:1004:LEU:HD22	2.54	0.42
1:A:507:LEU:HG	1:A:521:GLN:HE21	1.85	0.42
1:A:696:LEU:HD23	1:A:771:LEU:HB3	2.01	0.42
1:A:975:TYR:CD1	2:B:6:A:N6	2.88	0.41
1:A:941:ARG:HD2	1:A:961:TYR:CZ	2.56	0.41
1:A:922:GLU:O	1:A:929:ARG:HA	2.20	0.41
1:A:810:LYS:HG3	1:A:821:TRP:CE2	2.56	0.41
1:A:696:LEU:HD12	1:A:696:LEU:H	1.85	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	594/672 (88%)	563 (95%)	31 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	460/617 (75%)	460 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	641	ASN
1	A	800	ASN
1	A	1032	GLN
1	A	1096	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	3/6 (50%)	1 (33%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	4	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	610/672 (90%)	0.62	55 (9%) 15 8	63, 122, 176, 208	0
2	B	4/6 (66%)	0.85	0 100 100	241, 259, 280, 298	0
All	All	614/678 (90%)	0.62	55 (8%) 15 8	63, 122, 179, 298	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	601	ILE	5.4
1	A	891	GLU	4.2
1	A	1111	GLU	3.6
1	A	682	VAL	3.4
1	A	595	TYR	3.3
1	A	1109	VAL	3.3
1	A	1059	MET	3.2
1	A	1056	LEU	3.2
1	A	883	GLY	3.0
1	A	856	GLU	3.0
1	A	765	ILE	2.9
1	A	587	GLU	2.9
1	A	612	GLU	2.9
1	A	802	GLU	2.8
1	A	1063	GLU	2.8
1	A	676	SER	2.7
1	A	760	PHE	2.6
1	A	468	THR	2.6
1	A	515	TYR	2.6
1	A	674	SER	2.5
1	A	979	LEU	2.5
1	A	1031	ASN	2.5
1	A	677	GLY	2.5
1	A	513	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	581	ARG	2.5
1	A	1076	LEU	2.5
1	A	611	SER	2.4
1	A	1038	ILE	2.4
1	A	589	GLN	2.4
1	A	680	LYS	2.4
1	A	588	PRO	2.4
1	A	719	THR	2.3
1	A	958	VAL	2.3
1	A	658	CYS	2.3
1	A	686	LEU	2.3
1	A	614	ILE	2.3
1	A	781	THR	2.3
1	A	967	HIS	2.3
1	A	935	LYS	2.2
1	A	484	LEU	2.2
1	A	980	PRO	2.2
1	A	878	ASP	2.2
1	A	1098	VAL	2.2
1	A	668	LYS	2.1
1	A	779	LYS	2.1
1	A	988	SER	2.1
1	A	803	THR	2.1
1	A	928	ILE	2.1
1	A	1110	PRO	2.1
1	A	464	ILE	2.0
1	A	493	ASN	2.0
1	A	1039	TYR	2.0
1	A	788	GLU	2.0
1	A	841	THR	2.0
1	A	982	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.