



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 07:28 PM UTC

PDB ID : 3RA0 / pdb_00003ra0
Title : Crystal Structure of a StWhy2 K67A-dT32 complex
Authors : Cappadocia, L.; Brisson, N.; Sygusch, J.
Deposited on : 2011-03-26
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	
2	B	32	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Why2 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	161	Total	C	N	O	S	0	0	0
			1262	803	225	229	5			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	MET	-	initiating methionine	UNP D9J034
A	67	ALA	LYS	engineered mutation	UNP D9J034
A	217	LEU	-	expression tag	UNP D9J034
A	218	GLU	-	expression tag	UNP D9J034
A	219	HIS	-	expression tag	UNP D9J034
A	220	HIS	-	expression tag	UNP D9J034
A	221	HIS	-	expression tag	UNP D9J034
A	222	HIS	-	expression tag	UNP D9J034
A	223	HIS	-	expression tag	UNP D9J034
A	224	HIS	-	expression tag	UNP D9J034


- Molecule 2 is a DNA chain called DNA 32-mer dT32.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	9	Total	C	N	O	P	0	0	0
			180	90	18	63	9			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total	O	0	0
			27	27		
3	B	3	Total	O	0	0
			3	3		

- Molecule 1: Why2 protein



4 Data and refinement statistics

Property	Value	Source
Space group	F 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	166.47Å 166.47Å 166.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.43 – 2.45 29.43 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.8 (29.43-2.45) 96.7 (29.43-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.45Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.250 , 0.267 0.248 , 0.272	Depositor DCC
R_{free} test set	606 reflections (7.89%)	wwPDB-VP
Wilson B-factor (Å ²)	67.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1472	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1293	0.70	0/1748
2	B	0.14	0/197	0.65	0/302
All	All	0.27	0/1490	0.69	0/2050

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1262	0	1249	33	0
2	B	180	0	109	13	0
3	A	27	0	0	0	0
3	B	3	0	0	1	0
All	All	1472	0	1358	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:ARG:HH11	1:A:81:ARG:HG3	1.01	1.08

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:ARG:HG3	1:A:81:ARG:NH1	1.82	0.89
1:A:81:ARG:HH11	1:A:81:ARG:CG	1.88	0.85
1:A:91:ARG:NH1	2:B:2:DT:H3	1.82	0.78
1:A:65:LYS:HG3	1:A:186:PRO:HD2	1.68	0.74
2:B:7:DT:H2''	2:B:8:DT:C5'	2.17	0.74
1:A:215:ARG:HA	1:A:215:ARG:NH1	2.03	0.73
2:B:8:DT:C6	2:B:8:DT:H5''	2.26	0.71
2:B:7:DT:H2''	2:B:8:DT:H5''	1.75	0.67
1:A:91:ARG:HH12	2:B:2:DT:H3	1.48	0.62
1:A:133:ASP:O	1:A:159:PRO:HD3	2.00	0.62
1:A:101:PRO:HG3	1:A:183:PHE:HE1	1.64	0.61
1:A:81:ARG:HG2	1:A:87:VAL:HG22	1.82	0.61
2:B:7:DT:H2''	2:B:8:DT:O5'	2.02	0.60
2:B:8:DT:H2'	2:B:9:DT:C6	2.39	0.57
1:A:77:PRO:HG3	1:A:206:ILE:O	2.06	0.55
1:A:160:ASN:HB3	1:A:167:PHE:CE1	2.42	0.54
2:B:8:DT:C5'	2:B:8:DT:H6	2.19	0.54
1:A:214:ASN:O	1:A:215:ARG:HB3	2.08	0.54
2:B:8:DT:H2'	2:B:9:DT:C5	2.44	0.53
2:B:8:DT:C2	2:B:9:DT:N3	2.77	0.53
1:A:56:ARG:H	1:A:56:ARG:HD2	1.73	0.52
1:A:70:LEU:HD13	1:A:99:PHE:CZ	2.44	0.52
1:A:215:ARG:HA	1:A:215:ARG:HH11	1.75	0.52
1:A:203:LEU:HB3	1:A:204:PRO:HD3	1.91	0.52
2:B:8:DT:C5'	2:B:8:DT:C6	2.90	0.52
1:A:56:ARG:HD2	1:A:56:ARG:N	2.26	0.50
1:A:102:SER:HA	1:A:108:TYR:HA	1.94	0.48
1:A:153:LYS:HG2	1:A:172:VAL:HG22	1.96	0.47
2:B:1:DT:H71	3:B:35:HOH:O	2.16	0.45
1:A:99:PHE:O	1:A:113:ARG:HA	2.17	0.45
1:A:79:PHE:HB3	1:A:87:VAL:CG1	2.47	0.44
1:A:137:PHE:O	1:A:154:SER:HA	2.18	0.44
1:A:114:GLN:HG3	1:A:183:PHE:HE2	1.83	0.43
1:A:143:MET:HE3	2:B:3:DT:O2	2.18	0.43
1:A:150:GLN:HB3	1:A:176:ASN:OD1	2.19	0.42
1:A:73:GLU:HA	1:A:74:PRO:HD3	1.85	0.42
1:A:134:SER:HB3	1:A:159:PRO:HD2	2.01	0.42
1:A:132:ARG:HA	1:A:132:ARG:HE	1.84	0.42
1:A:214:ASN:O	1:A:215:ARG:CB	2.68	0.42
1:A:170:LEU:O	1:A:182:ARG:HA	2.21	0.41
1:A:103:VAL:HG23	1:A:107:LYS:HE2	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:SER:HA	1:A:183:PHE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/178 (89%)	147 (92%)	12 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/150 (91%)	130 (96%)	6 (4%)	25	37

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	ARG
1	A	81	ARG
1	A	132	ARG
1	A	144	LEU
1	A	158	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	162	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	161/178 (90%)	0.37	3 (1%) 66 67	49, 66, 90, 99	0
2	B	9/32 (28%)	0.95	0 100 100	45, 53, 64, 65	9 (100%)
All	All	170/210 (80%)	0.40	3 (1%) 67 69	45, 66, 90, 99	9 (5%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	103	VAL	2.6
1	A	143	MET	2.6
1	A	106	ARG	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.