



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 08:06 AM UTC

PDB ID : 3RAE / pdb_00003rae
Title : Quinolone(Levofloxacin)-DNA cleavage complex of type IV topoisomerase from *S. pneumoniae*
Authors : Laponogov, I.; Pan, X.-S.; Veselkov, D.A.; McAuley, K.E.; Fisher, L.M.; Sanderson, M.R.
Deposited on : 2011-03-28
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	 68% 25% . .
1	B	496	 68% 27% . .
2	C	268	 9% 53% 21% . 22%
2	D	268	 5% 59% 16% . 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	7	
4	F	11	
5	G	7	
6	H	11	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	MG	C	701	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 11234 atoms, of which 38 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase 4 subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	482	Total	C	N	O	S	0	1	0
			3736	2366	648	709	13			
1	B	482	Total	C	N	O	S	0	1	0
			3717	2359	640	705	13			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	257	THR	ILE	SEE REMARK 999	UNP P72525
A	489	LEU	-	expression tag	UNP P72525
A	490	GLU	-	expression tag	UNP P72525
A	491	HIS	-	expression tag	UNP P72525
A	492	HIS	-	expression tag	UNP P72525
A	493	HIS	-	expression tag	UNP P72525
A	494	HIS	-	expression tag	UNP P72525
A	495	HIS	-	expression tag	UNP P72525
A	496	HIS	-	expression tag	UNP P72525
B	257	THR	ILE	SEE REMARK 999	UNP P72525
B	489	LEU	-	expression tag	UNP P72525
B	490	GLU	-	expression tag	UNP P72525
B	491	HIS	-	expression tag	UNP P72525
B	492	HIS	-	expression tag	UNP P72525
B	493	HIS	-	expression tag	UNP P72525
B	494	HIS	-	expression tag	UNP P72525
B	495	HIS	-	expression tag	UNP P72525
B	496	HIS	-	expression tag	UNP P72525

- Molecule 2 is a protein called DNA topoisomerase 4 subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	208	Total	C	N	O	S	0	0	0
			1445	917	254	268	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	209	Total	C	N	O	S	0	0	0
			1456	924	253	273	6			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	380	MET	-	expression tag	UNP Q59961
C	381	GLY	-	expression tag	UNP Q59961
C	382	HIS	-	expression tag	UNP Q59961
C	383	HIS	-	expression tag	UNP Q59961
C	384	HIS	-	expression tag	UNP Q59961
C	385	HIS	-	expression tag	UNP Q59961
C	386	HIS	-	expression tag	UNP Q59961
C	387	HIS	-	expression tag	UNP Q59961
C	388	HIS	-	expression tag	UNP Q59961
C	389	HIS	-	expression tag	UNP Q59961
C	390	HIS	-	expression tag	UNP Q59961
C	391	HIS	-	expression tag	UNP Q59961
C	392	SER	-	expression tag	UNP Q59961
C	393	SER	-	expression tag	UNP Q59961
C	394	GLY	-	expression tag	UNP Q59961
C	395	HIS	-	expression tag	UNP Q59961
C	396	ILE	-	expression tag	UNP Q59961
C	397	ASP	-	expression tag	UNP Q59961
C	398	ASP	-	expression tag	UNP Q59961
C	399	ASP	-	expression tag	UNP Q59961
C	400	ASP	-	expression tag	UNP Q59961
C	401	LYS	-	expression tag	UNP Q59961
C	402	HIS	-	expression tag	UNP Q59961
C	403	MET	-	expression tag	UNP Q59961
C	460	ILE	VAL	SEE REMARK 999	UNP Q59961
C	644	ALA	THR	SEE REMARK 999	UNP Q59961
D	380	MET	-	expression tag	UNP Q59961
D	381	GLY	-	expression tag	UNP Q59961
D	382	HIS	-	expression tag	UNP Q59961
D	383	HIS	-	expression tag	UNP Q59961
D	384	HIS	-	expression tag	UNP Q59961
D	385	HIS	-	expression tag	UNP Q59961
D	386	HIS	-	expression tag	UNP Q59961
D	387	HIS	-	expression tag	UNP Q59961
D	388	HIS	-	expression tag	UNP Q59961
D	389	HIS	-	expression tag	UNP Q59961

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	390	HIS	-	expression tag	UNP Q59961
D	391	HIS	-	expression tag	UNP Q59961
D	392	SER	-	expression tag	UNP Q59961
D	393	SER	-	expression tag	UNP Q59961
D	394	GLY	-	expression tag	UNP Q59961
D	395	HIS	-	expression tag	UNP Q59961
D	396	ILE	-	expression tag	UNP Q59961
D	397	ASP	-	expression tag	UNP Q59961
D	398	ASP	-	expression tag	UNP Q59961
D	399	ASP	-	expression tag	UNP Q59961
D	400	ASP	-	expression tag	UNP Q59961
D	401	LYS	-	expression tag	UNP Q59961
D	402	HIS	-	expression tag	UNP Q59961
D	403	MET	-	expression tag	UNP Q59961
D	460	ILE	VAL	SEE REMARK 999	UNP Q59961
D	644	ALA	THR	SEE REMARK 999	UNP Q59961

- Molecule 3 is a DNA chain called 5'-D(*CP*AP*TP*GP*AP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	7	Total	C	N	O	P	0	0	0
			140	69	27	38	6			

- Molecule 4 is a DNA chain called 5'-D(P*AP*GP*TP*CP*AP*TP*TP*CP*AP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	11	Total	C	N	O	P	0	0	0
			225	108	39	67	11			

- Molecule 5 is a DNA chain called 5'-D(*CP*GP*TP*GP*CP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	G	7	Total	C	N	O	P	0	0	0
			139	68	25	40	6			


- Molecule 6 is a DNA chain called 5'-D(P*GP*AP*CP*TP*AP*TP*GP*CP*AP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	H	11	Total	C	N	O	P	0	0	0
			226	107	43	65	11			

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	2	Total Mg 2 2	0	0
7	B	1	Total Mg 1 1	0	0
7	C	1	Total Mg 1 1	0	0
7	D	1	Total Mg 1 1	0	0
7	G	1	Total Mg 1 1	0	0

- Molecule 8 is (3S)-9-fluoro-3-methyl-10-(4-methylpiperazin-1-yl)-7-oxo-2,3-dihydro-7H-[1,4]oxazino[2,3,4-ij]quinoline-6-carboxylic acid (CCD ID: LFX) (formula: C₁₈H₂₀FN₃O₄).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	F	1	Total 45	C 18	F 1	H 19	N 3	O 4	0	0
8	H	1	Total 45	C 18	F 1	H 19	N 3	O 4	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	23	Total O 23 23	0	0
9	B	21	Total O 21 21	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	C	1	Total 1	O 1	0	0
9	D	3	Total 3	O 3	0	0
9	E	4	Total 4	O 4	0	0
9	G	2	Total 2	O 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA topoisomerase 4 subunit A



• Molecule 1: DNA topoisomerase 4 subunit A



Chain C:

Amino Acid	Percentage
T640	9%
LEU	53%
GLU	21%
GLU	22%
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	
GLU	
ALA	
ALA	
THR	
VAL	
PHE	
T640	
LEU	
GLU	

Chain D:

5%

59%

16%

22%

MET GLY HIS HIS HIS HIS HIS HIS HIS HIS ASP ASP ASP LYS MET LYS ASN LYS LYS ASP LYS LEU LEU SER SER GLY K415 S421 N427 E428 L429 Y430 L431 R445 D446 R447 K448 F449 L453 V459 I460 T462 A463 K464 D465 I470 E474 E475 T478 R479 T482 D490 F491 S492 K500 I501 T509 D510 H513 T516 L517 L518 L519 T520 F521 F522 T523 G524 Y525 M526 E527 P528 Y536 M545 SER LYS GLY LYS GLY LYS GLU GLU VAL VAL ALA Y557 Q570 PHE GLY LYS GLY ALA THR

Chain E:


Chain F: 45% 55%

Chain G: 71% 29%



- Molecule 6: 5'-D(P*GP*AP*CP*TP*AP*TP*GP*CP*AP*CP*G)-3'

Chain H:  36% 64%



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	157.83Å 157.83Å 211.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.83 – 2.90 41.83 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (41.83-2.90) 99.4 (41.83-2.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.186 , 0.226 0.181 , 0.226	Depositor DCC
R_{free} test set	9357 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	61.9	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 61.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.048 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11234	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, LFX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/3800	0.95	9/5146 (0.2%)
1	B	0.58	0/3781	0.89	3/5126 (0.1%)
2	C	0.50	0/1471	0.85	0/2008
2	D	0.50	0/1482	0.86	0/2022
3	E	0.37	0/157	1.07	0/241
4	F	0.33	0/251	1.10	0/385
5	G	0.36	0/155	0.94	0/238
6	H	0.32	0/253	0.98	0/388
All	All	0.55	0/11350	0.92	12/15554 (0.1%)

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	171	SER	N-CA-C	6.37	114.87	108.75
1	A	170	ILE	CB-CA-C	-6.24	103.30	110.41
1	A	159	PHE	CA-C-N	5.93	127.25	119.84
1	A	159	PHE	C-N-CA	5.93	127.25	119.84
1	A	213	PHE	CA-C-N	5.61	125.28	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3736	0	3650	122	0
1	B	3717	0	3630	121	0
2	C	1445	0	1242	60	0
2	D	1456	0	1243	53	0
3	E	140	0	78	2	0
4	F	225	0	126	5	0
5	G	139	0	78	2	0
6	H	226	0	124	9	0
7	A	2	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
7	G	1	0	0	0	0
8	F	26	19	19	0	0
8	H	26	19	19	1	0
9	A	23	0	0	1	0
9	B	21	0	0	0	0
9	C	1	0	0	0	0
9	D	3	0	0	0	0
9	E	4	0	0	0	0
9	G	2	0	0	0	0
All	All	11196	38	10209	352	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 352 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:ARG:HH11	1:A:60:ARG:HB2	1.07	1.19
1:A:28:ARG:HG3	1:A:28:ARG:HH11	1.05	1.18
2:D:583:LEU:HD23	2:D:586:MET:HE1	1.26	1.18
1:B:307:ASN:HD22	1:B:310:LEU:HB2	0.97	1.11
2:D:520:THR:HG21	2:D:622:MET:HG3	1.34	1.07

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/496 (97%)	454 (94%)	22 (5%)	5 (1%)	12	39
1	B	481/496 (97%)	447 (93%)	29 (6%)	5 (1%)	12	39
2	C	202/268 (75%)	186 (92%)	15 (7%)	1 (0%)	24	54
2	D	203/268 (76%)	191 (94%)	11 (5%)	1 (0%)	24	54
All	All	1367/1528 (90%)	1278 (94%)	77 (6%)	12 (1%)	14	41

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	PHE
1	A	329	MET
1	B	168	THR
1	B	293	ARG
2	C	496	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/431 (88%)	341 (90%)	37 (10%)	7	25
1	B	376/431 (87%)	346 (92%)	30 (8%)	11	34
2	C	111/224 (50%)	97 (87%)	14 (13%)	4	14
2	D	111/224 (50%)	99 (89%)	12 (11%)	6	21
All	All	976/1310 (74%)	883 (90%)	93 (10%)	8	26

5 of 93 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	338	ARG
2	C	477	ASN
1	B	372	LEU
1	B	462	LEU
2	C	524	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	307	ASN
1	B	326	ASN
2	C	477	ASN
1	B	324	ASN
1	B	328	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	LFX	F	101	7	29,29,29	0.83	1 (3%)	44,44,44	1.45	9 (20%)
8	LFX	H	101	7	29,29,29	0.83	1 (3%)	44,44,44	1.27	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	LFX	F	101	7	-	0/8/27/27	0/4/4/4
8	LFX	H	101	7	-	0/8/27/27	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	F	101	LFX	O02-C15	3.11	1.29	1.23
8	H	101	LFX	O02-C15	2.73	1.28	1.23

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	F	101	LFX	O-C-C17	-3.82	114.76	122.67
8	H	101	LFX	O-C-C17	-2.84	116.80	122.67
8	H	101	LFX	O01-C06-C04	-2.71	115.06	117.90
8	F	101	LFX	C10-C15-C17	-2.40	112.62	115.64
8	F	101	LFX	O01-C06-C04	-2.35	115.44	117.90

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	H	101	LFX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	482/496 (97%)	-0.46	1 (0%) 91 89	36, 63, 101, 138	1 (0%)
1	B	482/496 (97%)	-0.44	2 (0%) 88 85	31, 63, 95, 135	1 (0%)
2	C	208/268 (77%)	0.52	24 (11%) 9 8	67, 105, 148, 170	0
2	D	209/268 (77%)	0.43	14 (6%) 24 19	68, 99, 133, 167	0
3	E	7/7 (100%)	0.04	1 (14%) 6 5	62, 70, 102, 151	0
4	F	11/11 (100%)	-0.32	0 100 100	71, 91, 121, 146	0
5	G	7/7 (100%)	-0.39	0 100 100	57, 70, 105, 139	0
6	H	11/11 (100%)	-0.33	0 100 100	67, 86, 111, 122	0
All	All	1417/1564 (90%)	-0.17	42 (2%) 52 43	31, 71, 127, 170	2 (0%)

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	589	ASP	4.5
2	C	469	ASP	4.4
1	B	3	ASN	4.1
2	C	565	GLU	4.0
2	C	495	ASP	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	MG	C	701	1/1	0.53	0.40	98,98,98,98	0
7	MG	G	101	1/1	0.82	0.23	103,103,103,103	0
7	MG	A	502	1/1	0.83	0.18	89,89,89,89	0
7	MG	D	701	1/1	0.84	0.27	72,72,72,72	0
7	MG	A	501	1/1	0.85	0.11	72,72,72,72	0
7	MG	B	700	1/1	0.91	0.09	70,70,70,70	0
8	LFX	F	101	26/26	0.95	0.10	78,96,121,127	0
8	LFX	H	101	26/26	0.96	0.09	85,102,118,125	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LFX F 101:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFX H 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.