



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 05:15 PM UTC

PDB ID : 6RAV / pdb_00006rav
Title : Complement factor B protease domain in complex with the reversible inhibitor 4-((2S,4S)-4-ethoxy-1-((5-methoxy-7-methyl-1H-indol-4-yl)methyl)piperidin-2-yl)benzoic acid
Authors : Adams, C.M.; Sellner, H.; Ehara, T.; Mac Sweeney, A.; Crowley, M.; Anderson, K.; Karki, R.; Mainolfi, N.; Valeur, E.; Sirockin, F.; Gerhartz, B.; Erbel, P.; Hughes, N.; Smith, T.M.; Cumin, F.; Argikar, U.; Mogi, M.; Sedrani, R.; Wiesmann, C.; Jaffee, B.; Maibaum, J.; Flohr, S.; Harrison, R.; Eder, J.
Deposited on : 2019-04-08
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	: 4-5-2 with Phenix2.0
Mogul	: 2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	: 2.0
EDS	: 3.0
Buster-report	: wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	: 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	: 9.0.010 (Gargrove)
Density-Fitness	: 1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	291	2% 90% 8% ..
1	BBB	291	3% 85% 9% 5%

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9377 atoms, of which 4455 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement factor B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	287	Total	C	H	N	O	S	105	3	0
			4548	1445	2273	397	421	12			
1	BBB	275	Total	C	H	N	O	S	99	7	0
			4374	1401	2182	370	409	12			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	O	S	0	0
			5	4	1		
2	AAA	1	Total	O	S	0	0
			5	4	1		
2	BBB	1	Total	O	S	0	0
			5	4	1		
2	BBB	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	BBB	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 4-[(2 {S},4 {S})-4-ethoxy-1-[(5-methoxy-7-methyl-1 {H}-indol-4-yl)methyl]piperidin-2-yl]benzoic acid (CCD ID: JGQ) (formula: C₂₅H₃₀N₂O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	N	O	0	0
			31	25	2	4		
3	BBB	1	Total	C	N	O	0	0
			31	25	2	4		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	BBB	2	Total	Zn	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	198	Total	O	0	0
			198	198		
5	BBB	168	Total	O	0	0
			168	168		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Complement factor B

Chain AAA:  2% 90% 8% ..



- Molecule 1: Complement factor B

Chain BBB:  3% 85% 9% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.46Å 97.45Å 61.81Å 90.00° 100.47° 90.00°	Depositor
Resolution (Å)	48.72 – 1.70 48.72 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.5 (48.72-1.70) 98.5 (48.72-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0238 2018/15/10	Depositor
R, R_{free}	0.195 , 0.227 (Not available) , 0.227	Depositor DCC
R_{free} test set	3435 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 27.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9377	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JGQ, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.01	0/2334	1.15	0/3157
1	BBB	1.00	0/2256	1.15	1/3056 (0.0%)
All	All	1.01	0/4590	1.15	1/6213 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	164(B)	ASP	CB-CA-C	-6.05	100.39	110.68

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2275	2273	2257	13	2
1	BBB	2192	2182	2163	17	0
2	AAA	10	0	0	1	0
2	BBB	15	0	0	0	0
3	AAA	31	0	0	0	0
3	BBB	31	0	0	0	0
4	BBB	2	0	0	0	2
5	AAA	198	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	BBB	168	0	0	0	0
All	All	4922	4455	4420	29	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:30[A]:GLN:HE21	1:BBB:43:MET:HE3	1.13	1.11
1:BBB:30[A]:GLN:NE2	1:BBB:43:MET:HE3	1.91	0.85
1:BBB:91:HIS:HD2	1:BBB:93:ASN:H	1.36	0.72
1:BBB:30[A]:GLN:HE21	1:BBB:43:MET:CE	1.98	0.68
1:BBB:27:GLN:NE2	1:BBB:29:TRP:HE1	1.95	0.65
1:AAA:1(A):LEU:HB3	1:AAA:3:MET:HE3	1.87	0.56
1:AAA:208:ILE:HD12	1:AAA:208:ILE:N	2.20	0.56
1:AAA:97(D):GLU:HG3	1:BBB:250:LEU:OXT	2.07	0.53
1:BBB:192[B]:ARG:HG3	1:BBB:218(A):ASP:OD2	2.08	0.53
1:AAA:135:LYS:HE2	1:AAA:159:GLU:OE1	2.11	0.51
1:BBB:49[B]:GLU:H	1:BBB:49[B]:GLU:CD	2.19	0.50
1:BBB:129:LEU:HA	1:BBB:134:ILE:HD13	1.96	0.48
1:AAA:129:LEU:HA	1:AAA:134:ILE:HD13	1.97	0.47
1:BBB:125(H):LEU:O	1:BBB:204:ARG:NH2	2.48	0.47
1:BBB:175:VAL:HG23	1:BBB:176:VAL:HG23	1.99	0.45
1:AAA:97(A):GLY:HA3	5:AAA:583:HOH:O	2.15	0.45
1:AAA:79:GLU:HB3	2:AAA:302:SO4:O2	2.17	0.45
1:BBB:91:HIS:CD2	1:BBB:93:ASN:H	2.26	0.44
1:AAA:125(F):ARG:NH2	5:AAA:401:HOH:O	2.29	0.44
1:AAA:172(L):ILE:O	1:AAA:175:VAL:HG22	2.19	0.43
1:BBB:30[A]:GLN:CG	1:BBB:43:MET:HE3	2.49	0.42
1:AAA:89:LEU:HD13	1:AAA:241:LYS:HD2	2.00	0.42
1:BBB:1(A):LEU:HD12	1:BBB:1(A):LEU:HA	1.84	0.42
1:AAA:175:VAL:HG23	1:AAA:176:VAL:HG23	2.01	0.42
1:AAA:192:ARG:NE	5:AAA:402:HOH:O	2.38	0.41
1:BBB:124:PRO:HB3	1:BBB:232:PHE:CZ	2.54	0.41
1:BBB:172(L):ILE:HD12	1:BBB:172(L):ILE:HA	2.00	0.41
1:BBB:201:VAL:HG12	1:BBB:203:LYS:HG3	2.02	0.41
1:AAA:124:PRO:HB3	1:AAA:232:PHE:CZ	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:62(D):HIS:HE2	4:BBB:305:ZN:ZN[1_556]	1.16	0.44
1:AAA:223:HIS:HD1	4:BBB:306:ZN:ZN[1_655]	1.20	0.40

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	249/256 (97%)	243 (98%)	6 (2%)	43	26
1	BBB	241/256 (94%)	235 (98%)	6 (2%)	42	24
All	All	490/512 (96%)	478 (98%)	12 (2%)	44	26

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	1(A)	LEU
1	AAA	3	MET
1	AAA	81	ARG
1	AAA	97(C)	LYS
1	AAA	119	ARG
1	AAA	125(T)	LYS
1	BBB	1(A)	LEU
1	BBB	40	GLU
1	BBB	41[A]	SER
1	BBB	41[B]	SER
1	BBB	154	LYS
1	BBB	172(G)	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	AAA	302	-	4,4,4	0.27	0	6,6,6	0.12	0
2	SO4	AAA	301	-	4,4,4	0.31	0	6,6,6	0.13	0
3	JGQ	BBB	304	-	34,34,34	1.74	7 (20%)	41,48,48	1.97	12 (29%)
2	SO4	BBB	303	-	4,4,4	0.30	0	6,6,6	0.09	0
2	SO4	BBB	301	-	4,4,4	0.32	0	6,6,6	0.13	0
2	SO4	BBB	302	-	4,4,4	0.28	0	6,6,6	0.12	0
3	JGQ	AAA	303	-	34,34,34	1.98	11 (32%)	41,48,48	2.10	11 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JGQ	BBB	304	-	-	0/17/30/30	0/4/4/4
3	JGQ	AAA	303	-	-	2/17/30/30	0/4/4/4

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	303	JGQ	C12-N13	4.79	1.55	1.47
3	BBB	304	JGQ	O10-C5	4.22	1.44	1.37
3	AAA	303	JGQ	C18-N13	3.99	1.56	1.47
3	BBB	304	JGQ	C18-N13	3.59	1.56	1.47
3	BBB	304	JGQ	O27-C25	3.22	1.32	1.22
3	BBB	304	JGQ	C4-C5	3.10	1.44	1.38
3	AAA	303	JGQ	C12-C6	3.02	1.56	1.51
3	BBB	304	JGQ	C12-N13	2.93	1.52	1.47
3	AAA	303	JGQ	O26-C25	-2.89	1.21	1.30
3	AAA	303	JGQ	O27-C25	2.88	1.31	1.22
3	BBB	304	JGQ	C8-N9	-2.53	1.33	1.38
3	AAA	303	JGQ	C15-C16	2.52	1.57	1.51
3	AAA	303	JGQ	C1-C7	2.46	1.53	1.44
3	AAA	303	JGQ	C8-N9	-2.29	1.33	1.38
3	BBB	304	JGQ	O26-C25	-2.23	1.23	1.30
3	AAA	303	JGQ	C24-C23	2.19	1.42	1.38
3	AAA	303	JGQ	C2-C3	2.14	1.43	1.40
3	AAA	303	JGQ	C21-C20	2.08	1.42	1.38

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	BBB	304	JGQ	O10-C5-C6	6.31	121.81	115.30
3	AAA	303	JGQ	C1-C6-C5	-5.14	114.79	118.36
3	AAA	303	JGQ	C31-O10-C5	4.41	123.98	117.51
3	AAA	303	JGQ	C12-N13-C18	4.35	118.44	111.72
3	AAA	303	JGQ	O26-C25-C22	4.26	125.77	114.84
3	AAA	303	JGQ	C4-C3-C2	-3.82	113.88	118.25
3	BBB	304	JGQ	C2-C1-C7	-3.31	102.12	106.72
3	BBB	304	JGQ	O26-C25-C22	3.20	123.05	114.84
3	BBB	304	JGQ	O28-C16-C17	-3.19	99.61	110.00
3	BBB	304	JGQ	O10-C5-C4	-2.87	119.13	124.08
3	BBB	304	JGQ	C4-C5-C6	-2.78	118.95	122.06
3	BBB	304	JGQ	O27-C25-C22	-2.76	114.22	121.46
3	AAA	303	JGQ	C1-C7-C8	-2.72	103.82	106.64
3	BBB	304	JGQ	C31-O10-C5	2.72	121.50	117.51
3	BBB	304	JGQ	C12-N13-C18	2.72	115.92	111.72
3	BBB	304	JGQ	C14-N13-C18	-2.70	105.85	111.21
3	AAA	303	JGQ	O26-C25-O27	-2.47	118.04	123.35
3	BBB	304	JGQ	C17-C18-C19	-2.31	105.69	112.22
3	AAA	303	JGQ	O10-C5-C4	-2.28	120.14	124.08
3	AAA	303	JGQ	C23-C22-C25	-2.16	116.10	120.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	303	JGQ	C2-N9-C8	2.09	110.36	108.53
3	BBB	304	JGQ	C21-C20-C19	-2.07	119.12	121.18
3	AAA	303	JGQ	O27-C25-C22	-2.01	116.18	121.46

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	AAA	303	JGQ	C15-C16-O28-C29
3	AAA	303	JGQ	C17-C16-O28-C29


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	302	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	287/291 (98%)	0.05	5 (1%) 69 73	15, 29, 54, 75	3 (1%)
1	BBB	275/291 (94%)	0.26	9 (3%) 49 53	13, 30, 60, 106	7 (2%)
All	All	562/582 (96%)	0.15	14 (2%) 58 62	13, 29, 56, 106	10 (1%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	5	TRP	9.3
1	BBB	172(L)	ILE	3.5
1	AAA	5	TRP	3.5
1	BBB	172(I)	VAL	3.3
1	BBB	61	VAL	2.7
1	BBB	172(J)	LYS	2.7
1	AAA	172(L)	ILE	2.7
1	BBB	172(G)	ASP	2.6
1	AAA	82	ASP	2.5
1	BBB	1(A)	LEU	2.5
1	BBB	172(H)	LYS	2.3
1	AAA	20	LYS	2.1
1	AAA	220(G)	LYS	2.1
1	BBB	173	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	AAA	302	5/5	0.68	0.14	74,76,90,90	0
2	SO4	BBB	302	5/5	0.75	0.12	63,72,78,82	0
2	SO4	BBB	303	5/5	0.82	0.12	57,93,99,102	0
2	SO4	AAA	301	5/5	0.83	0.10	78,79,82,87	0
2	SO4	BBB	301	5/5	0.86	0.10	56,62,68,72	0
3	JGQ	BBB	304	31/31	0.94	0.08	24,29,37,43	0
3	JGQ	AAA	303	31/31	0.95	0.07	22,27,35,39	0
4	ZN	BBB	306	1/1	0.99	0.02	29,29,29,29	0
4	ZN	BBB	305	1/1	1.00	0.01	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around JGQ BBB 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around JGQ AAA 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.